

The Role of Rainstorm Properties on Crop-Land Soil Erosion: Coupling Event-Scale Modeling with a Stochastic Rainfall Generator for Estimating Erosion Risks

Yuval Shmilovitz¹, Francesco Marra¹, Haiyan Wei², Eli Argaman³, Mark Nearing⁴, David Goodrich⁴, Shmuel Assouline⁵, and Efrat Morin¹

¹The Hebrew University of Jerusalem

²University of Arizona

³Ministry of Agriculture and Rural Development

⁴USDA-ARS

⁵Agricultural Research Organization Volcani Center

November 22, 2022

Abstract

Soil erosion is a worldwide agricultural and environmental problem that threatens food security and ecosystem viability. In arable environments, the primary cause of soil loss is short and intense storms that are characterized with high spatiotemporal variability. The complex nature of these erosive events imposes a great challenge for erosion modeling and risk analysis. Accurate high-resolution measurement of rain intensity is often lacking or sparsely available. As a result, many studies rely on coarser-resolution rainfall data that often fail to address the impact of intra-storm properties. In this study, based on a novel statistical method, we quantify the discrete and cumulative multiannual impact of rainstorm regime on runoff and soil erosion to better understand the most important rainstorm properties on erosion rate and amount, and, to provide storm-scale risk analyses. Central to our analyses is the coupling of a processes-based crop-land erosion model, Dynamic Water Erosion Prediction Project (DWEPP), with a stochastic rainfall generator that produces localized rainfall statistics at 5-min resolution (CoSMoS). To our knowledge, this is the first study that calibrated DWEPP runoff and sediment at the plot-scale on cropland. The model had an acceptable fit with measured event runoff and sediment data collected in northern Israel ($NSE = 0.79 - 0.82$). We then generated 300-year stochastic simulations of event-based runoff and sediment yield and used them to estimate erosion risk and calibrate a state-of-the-art frequency analysis method that explicitly accounts for rainstorms occurrence and properties. Results indicate that in the study area, high erosion rate events are characterized by intense rain bursts of short duration (shorter than the usually adopted erosivity index of 30-min), and not necessary by events of large volume accumulation or long duration. On these bases, we proposed an optimal rainfall erosivity index that combines intra-storm properties for the study area. As changes in rainstorm properties are expected under a changing climate, we expect our methodology to be a valuable tool for investigating the global concerns about future changes in soil erosion rate.

Hosted file

essoar.10506221.1.docx available at <https://authorea.com/users/540307/articles/600183-the-role-of-rainstorm-properties-on-crop-land-soil-erosion-coupling-event-scale-modeling-with-a-stochastic-rainfall-generator-for-estimating-erosion-risks>

The role of rainstorm properties on crop-land soil erosion: coupling event-scale modeling with a stochastic rainfall generator for estimating erosion risks

Yuval Shmilovitz¹, Francesco Marra^{1,6}, Haiyan Wei⁴, Eli Argaman², Mark Nearing⁵, David Goodrich⁵, Shmuel Assouline³, Efrat Morin¹

- 1- The Fredy & Nadine Herrmann Institute of Earth Sciences, The Hebrew University of Jerusalem, Jerusalem, Israel
- 2- Soil Erosion Research Station, Ministry of Agriculture and Rural Development, Rishon LeZion, Israel
- 3- Institute of Soil, Water and Environmental Sciences, Agricultural Research Organization, Volcani Center, Rishon LeZion, Israel
- 4- University of Arizona, School of Natural Resources and the Environment, Tucson, AZ, USA
- 5- USDA-ARS Southwest Watershed Research Center, Tucson, AZ, USA
- 6- Institute of Atmospheric Sciences and Climate, National Research Council of Italy, Bologna, Italy

Abstract

Soil erosion is a worldwide agricultural and environmental problem that threatens food security and ecosystem viability. In arable environments, the primary cause of soil loss is short and intense storms that are characterized with high spatiotemporal variability. The complex nature of these erosive events imposes a great challenge for erosion modeling and risk analysis. Accurate high-resolution measurement of rain intensity is often lacking or sparsely available. As a result, many studies rely on coarser-resolution rainfall data that often fail to address the impact of intra-storm properties. In this study, based on a novel statistical method, we quantify the discrete and cumulative multiannual impact of rainstorm regime on runoff and soil erosion to better understand the most important rainstorm properties on erosion rate and amount, and, to provide storm-scale risk analyses.

Central to our analyses is the coupling of a processes-based crop-land erosion model, Dynamic Water Erosion Prediction Project (DWEPP), with a stochastic rainfall generator that produces localized rainfall statistics at 5-min resolution (CoSMoS). To our knowledge, this is the first study that calibrated DWEPP runoff and sediment at the plot-scale on cropland. The model had an acceptable fit with measured event runoff and sediment data collected in northern Israel ($NSE = 0.79 - 0.82$). We then generated 300-year stochastic simulations of event-based runoff and sediment yield and used them to estimate erosion risk and calibrate a state-of-the-art frequency analysis method that explicitly accounts for rainstorms occurrence and properties. Results indicate that in the study area, high erosion rate events are characterized by intense rain bursts of short duration (shorter than the usually adopted erosivity index of 30-min), and not necessary by events of large volume accumulation or long duration. On these bases, we proposed an optimal rainfall erosivity index that combines intra-storm properties for the study area. As changes in rainstorm properties are expected under a changing climate, we expect our methodology to be a valuable tool for investigating the global concerns about future changes in soil erosion rate.