

# Vulnerability to Water Shortage Under Current and Future Water Supply-Demand Conditions Across U.S. River Basins

Hadi Heidari<sup>1</sup>, Mazdak Arabi<sup>2</sup>, and Travis Warziniack<sup>3</sup>

<sup>1</sup>Department of Civil and Environmental Engineering, Colorado State University

<sup>2</sup>Civil and Environmental Engineering Department, Colorado State University

<sup>3</sup>Rocky Mountain Research Station, U. S. Forest Service

November 22, 2022

## Abstract

Climate change, population growth, urbanization, and interactions thereof may alter the water supply-demand balance and lead to shifts in water shortage characteristics at different timescales. This study assesses the vulnerability of water supply systems to the interannual to the decadal water shortage events across the contiguous United States (CONUS) by characterizing shifts in intensity, duration, and frequency (IDF) from current (1986-2015) to future (2070-2099) periods. The water yield was estimated using the Variable Infiltration Capacity (VIC) model driven with the Multivariate Adaptive Constructed Analogs (MACA) climate model with RCP 4.5 and 8.5. The water demand was projected under the A1B population growth scenario. The Water Evaluation and Planning (WEAP) model was applied to determine water shortage conditions in which water demand exceeds water supply. Changes in characteristics of shortage events were assessed using the Mixture Gamma-GPD model. The results indicate that the frequency and intensity of over-year ( $D > 12$  months) events at the monthly scale and decadal ( $D > 10$  years) events at the annual scale tend to increase in the Southwest, Southern, middle Great Plain, and Great Lakes. Conversely, the frequency of interannual ( $D < 12$  months) events at the monthly scale and annual ( $D > 1$  year) and multi-year ( $D > 3$  years) events at the annual scale may increase in the West Coast. Basins with a higher rate of aridification may experience more frequent over-year events while basins with a decrease in aridification may undergo more frequent interannual events due to an increase in the variability of extreme weather anomalies within a year under climate change.



# Vulnerability to Water Shortage Under Current and Future Water Supply-Demand Conditions Across U.S. River Basins

Hadi Heidari, Mazdak Arabi, Travis Warziniack


Civil & Environmental Engineering, Colorado State University, Fort Collins, CO 80523

## Introduction

- Future changes in **climate** and **population** may cause a decrease in freshwater availability and an increase in water demand leading to more frequent **water shortage** conditions.
- The enhanced characterizations of changes in water shortage conditions from **interannual to decadal** events are requisite to the appropriate management and planning of future water resources, and improved implementation of regional adaptation and mitigation strategies.


## Objectives

- This study proposes an approach to improve the vulnerability assessments of U.S. river basins to the shortage at the interannual to decadal timescales by characterizing shifts in intensity, duration, and frequency (IDF) of water shortage events from current (1986-2015) to future (2070-2099) periods.




## (1) Climate Projection

- The **current** precipitation and temperature of U.S. river basins (1986-2015) were obtained from the Daymet dataset (1) and then biased corrected using the Parameter-elevation Regressions on Independent Slopes Model (PRISM) climate dataset (2) at the monthly scale.



- The **future** precipitation, temperature, and wind speed of U.S. river basins were obtained from the downscaled Multivariate Adaptive Constructed Analogs (MACA) datasets (3).


- We selected the **IPSL-CM5A-MR** model that represents the **worst-case** scenario for the future climate conditions (4).



## (2) Hydrological Projection


- The Variable Infiltration Capacity (VIC) hydrologic model was implemented in this study to evaluate the hydrologic response of the climate change emission scenarios.

- The model is set up, calibrated and evaluated by Oubeidillah et al. (2014) using matching the simulated monthly total streamflow with the observed monthly runoff from the USGS WaterWatch runoff dataset (5) for each HUC8 subbasin.



## (3) Water Demand Projection

- The monthly water demand of each HUC4 river basin was estimated by summing projections for six water use sectors including domestic and public, agricultural irrigation, thermoelectric, industrial, commercial, and mining, livestock, and aquaculture.
- The current water use data were obtained from the USGS water use circulars and for thermoelectric power water use from (Diehl & Harris, 2014). The future water withdrawal for each sector was estimated as the product of a water use driver such as population and irrigated area; and a water withdrawal rate such as domestic withdrawals per capita and irrigation withdrawal per unit area.
- The A1B scenario from the Intergovernmental Panel on Climate Change (IPCC) Fourth Assessment set of global socioeconomic scenarios was chosen to project future changes in population and income levels using the AIM global emissions model.




## (4) Water Supply Projection

- The Water Evaluation and Planning (WEAP) model was applied in this study to estimate the water supply allocated to each HUC4 river basin.
- The WEAP model used in this study was set up by Brown et al., (2019) (6) at HUC04 watersheds spatial scale. The WEAP model runs at the monthly time step for the period of 1985 to 2099 to calculate the past and future water supply of each HUC4 river basin.
- The water supply of a basin was defined as the amount of water available to meet demands for a given month and obtained from the sum of water yield, net trans-basin diversions, reservoir storage from the prior month, and inflow from upstream minus the sum of required instream flow release, reservoir evaporation, and any required release to satisfy downstream demands.

## (5) Water Shortage Projections

- The probabilistic approach developed by Heidari et al., (2020) (7) was used in this study to assess shifts in intensity, duration, and frequency (IDF) of water shortage events across the CONUS at the monthly and annual scales.
- The duration (D), intensity (I), and frequency (F) of a water shortage event were respectively defined as the number of consecutive months/years where water demand exceeds water supply, the cumulative water deficit divided by its duration and the number of times that a specific event occurs.
- The water shortage events were modeled using the mixture Gamma-GPD distribution.

$$F(x|r, a, \xi, \beta, u, \phi_u) = \begin{cases} (1 - \phi_u) \frac{G(x|r, a)}{G(u|r, a)} & x < u \\ (1 - \phi_u) + \phi_u g(x|\xi, \beta, u) & x \geq u \end{cases}$$



## References


- Thornton, P. E., Running, S. W., & White, M. A. (1997). Generating surfaces of daily meteorological variables over large regions of complex terrain. *Journal of Hydrology*, 190(3-4), 214-251. [https://doi.org/10.1016/S0022-1694\(96\)03128-9](https://doi.org/10.1016/S0022-1694(96)03128-9)
- Daly, C., Halbleib, M., Smith, J. I., Gibson, W. P., Doggett, M. K., Taylor, G. H., et al. (2008). Physiographically sensitive mapping of climatological temperature and precipitation across the conterminous United States. *International Journal of Climatology*, 28(15), 2031-2064. <https://doi.org/10.1002/joc.1688>
- Abatzoglou, J. T., & Brown, T. J. (2012). A comparison of statistical downscaling methods suited for wildfire applications. *International Journal of Climatology*, 32(5), 772-780. <https://doi.org/10.1002/joc.2312>
- Heidari, H., Arabi, M., Warziniack, T., & Kao, S. C. (2020). Assessing Shifts in Regional Hydroclimatic Conditions of U.S. River Basins in Response to Climate Change over the 21st Century. *Earth's Future*, 8(10), 1-14. <https://doi.org/10.1029/2020EF001657>
- Brakebill, J. W., Wolock, D. M., & Terziotti, S. E. (2011). Digital Hydrologic Networks Supporting Applications Related to Spatially Referenced Regression Modeling. *JAWRA Journal of the American Water Resources Association*, 47(5), 916-932. <https://doi.org/10.1111/j.1752-1688.2011.00578.x>
- Brown, T. C., Mahat, V., & Ramirez, J. A. (2019). Adaptation to Future Water Shortages in the United States Caused by Population Growth and Climate Change. *Earth's Future*, 7(3), 219-234. <https://doi.org/10.1029/2018EF001091>
- Heidari, H., Arabi, M., Ghanbari, M., & Warziniack, T. (2020). A Probabilistic Approach for Characterization of Sub-Annual Socioeconomic Drought Intensity- Duration-Frequency (IDF) Relationships in a Changing Environment. *Water*. <https://doi.org/10.3390/w12061522>
- Heidari, H., Arabi, M., Warziniack, T., & Kao, S. C. (2021). Vulnerability to Water Shortage Under Current and Future Water Supply-Demand Conditions Across U.S. River Basins. *Earth's Future*. <https://doi.org/10.1029/2021EF002278>

• This work was funded by NSF Sustainability Research Network (SRN) Cooperative Agreement 1444758 and a cooperative agreement with US Forest Service Research and Development, Rocky Mountain Research Station.

## Results and Conclusion

- The findings under the worst-case scenario at the conterminous scale indicates that the river basins located in the Southwest, Southern, and the middle Great Plain regions may experience more intense water shortage conditions at the end of the century.
- Conversely, the river basins located in the West Coast region are likely to experience a decrease in the intensity of water shortage conditions.
- The Southwest, Southern, and the middle Great Plain river basin are likely to experience less frequent interannual (D<12 months), annual (D>1 year), and multi-year (D>3 years) events in the future.
- However, the frequency of over-year (D>12 months) events at the sub-annual scale and decadal (D>10 years) events at the annual scale were projected to increase in the future in these regions.


### The Frequency Amplification Factor of Water Shortage Events At Monthly Scale




- Although the frequency of over-year (D>12 months), and decadal (D>10 years) events were estimated to decrease in the West Coast regions under the on average the driest climate change scenario for the CONUS, the interannual (D<12 months) events at the monthly scale and annual (D>1 year) and multi-year (D>3 years) events at the annual scale were projected to increase in the future in the West Coast regions.

- The intensity of water shortage events with longer duration and higher return periods is likely to be more affected in the future in response to climate changes and population growth.


### Intensity of Current and Future Water Shortage Events At monthly Scale



### Intensity of Current and Future Water Shortage Events At Annual Scale



### The Frequency Amplification Factor of Water Shortage Events At Annual Scale



### Basins that are more vulnerable to changes in demand (demand-based), supply (supply-based), or both (supply and demand based), or both

