

A unique case of venous anomalies in the situs solitus and normocardia

Ranko Zdravkovic¹, Aleksandar Redzek¹, Stamenko Susak¹, Jelena Vuckovic¹, Miodrag Golubovic¹, Dragan Lazarevic¹, Mirko Todic¹, and Lazar Velicki¹

¹Institute of Cardiovascular Diseases of Vojvodina

September 24, 2020

Abstract

A persistent left superior vena cava is relatively common venous anomaly. But, the presence of persistent left superior vena cava with the atresia of the right superior vena cava is a much rarer situation. The association of these anomalies with the drainage of the left hepatic vein into the right atrium is an extremely rare situation. Here we present one such case in the situs solitus and normocardia.

References

1. Shoiab I, Schaff H, Saran N, Charnin J, Ritter M. Recommendations for Perioperative Management in Patients With Absent Right Superior Vena Cava. *J Cardiothorac Vasc Anesth* 2019;33(6):1710-1713.
2. Uraz S, Duran C, Balci D, et al. A very rare venous anomaly in a living liver donor: left hepatic venous connection to the right atrium. *Transplant Proc* 2007;39(5):1714-5.

Conflict of Interest: The authors declare that there are no conflict of interests.

Ethics statement: Informed consent was obtained in writing form from the patient.

Hosted file

Figure.pdf available at <https://authorea.com/users/361483/articles/482882-a-unique-case-of-venous-anomalies-in-the-situs-solitus-and-normocardia>