

# Knowledge, Attitude and Practices among cardiac surgical patients towards Coronavirus Disease (COVID-19) at tertiary care center.

Danishwar Meena<sup>1</sup>, Pramod Chandolia<sup>1</sup>, Surendra Patel<sup>1</sup>, Prem Prakash Sharma<sup>1</sup>, Anupam Das<sup>1</sup>, Madhusudan Katti<sup>1</sup>, ANIRUDH MATHUR<sup>1</sup>, Devendra Gavade<sup>1</sup>, and Alok Sharma<sup>1</sup>

<sup>1</sup>All India Institute of Medical Sciences - Jodhpur

July 11, 2022

## Abstract

**Abstract: Purpose:** This study was conducted to assess the Knowledge, Attitude, and Practice (KAP) of cardiac surgical patients towards the ongoing Coronavirus Disease (COVID-19) pandemic, as they are among the high-risk group for morbidity and mortality associated with COVID-19. **Methods:** This cross-sectional survey was conducted from September 2021 to November 2021 at a tertiary care hospital in western Rajasthan. A total of 88 patients were enrolled, and a questionnaire was created with 20 questions to assess KAP towards COVID-19. The KAP of enrolled patients was compared according to age, gender, place of residence, and education status. **Results:** A total of 88 participants completed the survey questionnaire, of whom 41 were male, and 47 were female. The mean age of the participants was 41.5 years. The mean knowledge score of our study group was  $21.49 \pm 7.42$ , the mean attitude score was  $4.08 \pm 1.84$ , and the mean practice score was  $4.60 \pm 0.74$ . In the subgroup analysis, the difference in the knowledge score was statistically significant ( $p < 0.05$ ) in different age groups, however, the attitude and practice scores were comparable for all the age groups. There was no difference in KAP scores concerning the gender of the participants. KAP scores were significantly higher for the participants residing in urban areas than those from rural places. The KAP scores also showed a statistically significant relationship with the education status of the population ( $p < 0.001$ ). The knowledge score of the participants with graduation and above was  $32.50 \pm 2.31$ , whereas the illiterate subgroup scored  $13.72 \pm 3.15$ . Similar results were seen in the attitude and practice scores, with the mean values showing a gradual rise with the education status of the study population. **Conclusion:** This single-center KAP study on COVID-19 among cardiac patients who are considered high risk was able to postulate a comprehensive evaluation. The findings suggest that high-risk patients have decent knowledge scores and an overall optimistic outlook on surviving the pandemic. However, novel and far-reaching awareness programs are required for patients who are less educated and reside in rural areas as they are found to have lower KAP scores. **Keywords:** KAP, COVID 19, Cardiac surgery, Knowledge, Attitude, Practice

---

## Introduction

The COVID-19 outbreak first occurred at the Wuhan seafood market in early December 2019, where patients with respiratory illnesses were observed to have a causative association with seafood consumption. On 30th January 2020, the World Health Organization (WHO) declared it a Public Health Emergency of International Concern. The outbreak was officially linked to the coronavirus in February 2020 by WHO and termed COVID-19. India is the largest democracy in the world, with a largely rural and growing urban population base suffering a lot from the current pandemic situation. India recorded the first 1 lakh cases on 18th May 2020 and crossed the 8.5 lakh mark on 11th July. Despite the enforcement of a four-phased lockdown from 24th March to the end of May 2020, multiple waves of COVID-19 caused an enormous blow to the economy and health infrastructure<sup>1</sup>. A COVID-19 pandemic is an unprecedented event, resulting in uncertainty, job

losses & mental despair. It has also severely affected patients with pre-existing chronic diseases, who require regular medications and monitoring. The patients with pre-existing cardiac diseases were also severely affected by the pandemic.

As per "KAP theory," modification in human behavior involves three successive steps: knowledge acquisition and attitude generation, followed by the adoption of new practice. Various studies have proven that the KAP levels among individuals are paramount in preventing and effectively managing diseases<sup>2,3</sup>.

As the population's adherence to control measures during lockdown mainly depends on their knowledge, attitude, and practice towards COVID-19, it is assumed that KAP levels of cardiac patients will be the deciding factor in their combat against the current pandemic. Hence, this study assessed the knowledge, attitude, and practices about COVID-19 in cardiac surgical patients during the current pandemic era.

## Materials and Methods

The study was a cross-sectional survey done in the Department of Cardio-Thoracic and Vascular Surgery at the tertiary care hospital in western Rajasthan, India, from September 2021 to November 2021. The institute's ethical committee approved the study protocol, consent form, questionnaire, and information sheet, which followed the Declaration of Helsinki guidelines. All postoperative cardiac surgical patients with valvular heart disease, coronary artery disease, and congenital heart disease of age 18 years and above, surgically treated at the institute, were included after informed consent. The KAP questionnaire was designed based on the extensive literature review published on COVID-19 and guidelines issued by health ministry, Government of India. A questionnaire was created with 20 questions on the Knowledge, Attitude & Practices of cardiac surgical patients during this pandemic in Hindi and was validated by 2 Biostatisticians and two cardiac surgeons. The questionnaire comprised three parts: Demographic details, knowledge-attitude-practice (KAP), and access to medical facilities. The demographic and clinical details included age, gender, place of residence (rural vs. urban), educational status, surgical status, and current New York Heart Association (NYHA) grade. All the participants were evaluated during their follow-up visit by a single interviewer with the aid of a validated questionnaire at the outpatient clinic.

The KAP questionnaire consisted of a total of 20 questions. Fourteen questions were dedicated to assessing knowledge about the novel coronavirus disease, four questions pertaining to attitude towards pandemic, and finally, two questions related to practices towards COVID-19 (Table 1 – Questionnaire). These questions were based on public health information and Covid-19 guidelines from the Government of India. The questions assessed knowledge of clinical symptoms, spread, treatment, and prevention of COVID-19. Each correct response was awarded a score of 1 while do not know, and an incorrect response was given a zero score. Maximum scores were 43 for knowledge, 9 for attitude, and 8 for practice questions. The higher the total score, the more knowledgeable the participant was considered. The design, setting, analyses, and reporting of this study adhered to the STROBE guidelines for cross-sectional survey in epidemiology.

## Statistical Analysis

The statistical analysis was performed using the Statistical Package for Social Sciences (SPSS) 17.0 software program (SPSS Inc., Chicago, IL, USA). Descriptive data were depicted as numbers and percentages, while normally distributed data were presented as mean  $\pm$  standard deviation (SD). Comparisons between total knowledge score and categorical demographic variables (gender, educational status, residence) were made using the one-way analysis of variance (ANOVA) or Independent Student's t-test. A Pearson correlation coefficient analysis / Spearman's rho correlation was used to examine the association of two related variables. A chi-squared test was used to compare two attributes, and a p-value  $< 0.05$  was considered statistically significant.

## Results

Ninety-eight postoperative cardiac surgical patients were informed in detail and offered to enroll in the study. 88 patients agreed to participate after giving consent. Out of 88 participants who completed the study questionnaire, 41 were male, and 47 were female. The mean age of the study subjects was 41.5 years, and

most participants were from the Western region of India (96%). The demographic profile of the participants has been summarized in (Table 2).

The mean knowledge score of the study population was  $21.49 \pm 7.42$ , the mean attitude score was  $4.08 \pm 1.84$ , and the mean practice score was  $4.60 \pm 0.74$ . In the subgroup analysis, the difference in the knowledge score was statistically significant ( $p < 0.05$ ) in different age groups, with the mean knowledge score being higher in the age groups of 18-29 years ( $23.65 \pm 6.85$ ), 30-39 years ( $24.27 \pm 8.22$ ) and 40-49 years ( $22.47 \pm 7.50$ ) (Table 3). However, the attitude and practice scores were comparable for all the age groups. There was no difference in KAP scores concerning the gender of the participants (Table 4).

When compared to the place of residence, KAP scores were significantly higher for the participants residing in urban areas than those from rural areas. The mean knowledge score, attitude score, and practice score for the urban residents were  $26.86 \pm 6.9$ ,  $5.11 \pm 1.95$ , and  $4.94 \pm 0.86$ , respectively, and those for the rural areas were  $17.77 \pm 5.18$  ( $p < 0.001$ ),  $3.37 \pm 1.37$  ( $p < 0.001$ ) and  $4.37 \pm 0.53$  ( $p < 0.001$ ) (Table 5).

The KAP scores also showed a statistically significant relationship with the education status of the population ( $p < 0.001$ ). The knowledge score of the participants with graduation and above was  $32.50 \pm 2.31$ , whereas the illiterate subgroup scored  $13.72 \pm 3.15$ . Similar results were seen concerning the attitude and practice scores, with the mean values showing a gradual rise with the education status of the population from illiterate to graduation and above (Table 6). All these results were found to be statistically significant. The percentage score was computed in each subgroup of the education status category (knowledge, attitude, and practice). It also showed similar results, with all scores being higher in the participants who have at least graduated (Table 7).

## Discussion

The KAP plays a crucial role in dealing with major global health challenges like pandemics. India has suffered considerable losses in terms of human fatalities and economic growth due to the COVID-19 pandemic. As the scientific community attempts to find a cure to the disease, the population's knowledge, attitudes, and practices are of foremost importance when it comes to combatting the virus. Our study shows the trend of these variables in the general population attending a tertiary care hospital for various cardiac surgeries, which is considered a high-risk subgroup for coronavirus infection.

Our study reported significantly lower knowledge scores among the older population ( $> 50$  years) and illiterate individuals, similar to the study conducted by Christy et al. in South India<sup>4</sup>. The younger population probably has better social and print media access which may explain their better knowledge. The most common source of information about COVID-19 were televised programs, the world web, local press, and social media<sup>5,6</sup>. In recent times social media have emerged as the primary source of information, followed by web sources and scientific papers<sup>7,8</sup>. The above explains the higher scores of knowledge, attitudes, and practices in the educated subgroup.

Narayanaswamy et al.<sup>9</sup> studied the KAP scores among cardiac patients at a tertiary hospital in South India. They did not find any difference in the level of knowledge in the urban patients compared to the rural individuals. They also reported a lower level of practice in rural and urban participants, unlike in our study, where the level of knowledge, attitudes, and practice was significantly higher in urban patients. It may be explained by the inadequate reach of awareness in rural areas due to lesser access to newspapers, media, and healthcare.

Pal et al.<sup>10</sup> studied the KAP among another high-risk subgroup of Type 1 Diabetes mellitus. They also concluded that the level of knowledge was significantly higher in the educated and urban patients. Younger patients were found to have average knowledge, positive attitude, and healthier practices for preventing COVID-19. Less-educated individuals residing in rural areas generally tend to have wider gaps in KAP<sup>11,12</sup>. Differences in knowledge and behavior among urban and rural residents can result from socioeconomic incongruity between them<sup>13</sup>.

In a study conducted among medical students, Maheshwari et al.<sup>3</sup> reported appropriate knowledge, positive

attitude, and acceptable practice toward COVID-19. Even in this educated study population, the knowledge level was better in the younger age subgroup (21-23 years). Unlike our study, Ferdous et al. <sup>14</sup> found a better level of knowledge among the older participants in a study conducted in Bangladesh. However, when compared to the place of residence, the findings were similar to our study, wherein the urban individuals fared much better in terms of better knowledge, positive attitude, and healthier practices.

Our study shows that the rise in knowledge improves attitude and practices in the urban and educated subgroups of cardiac surgical patients. This is similar to the finding in the Korean study conducted by Lee et al. <sup>15</sup>.

A study by BS Tomar et al. <sup>16</sup> stated a strong relationship between gender and knowledge score towards COVID-19. This could be explained by underlying confounding factors such as education level and occupation, providing better information access. Contradictory to the above study, the KAP survey in the Saudi community by Al Hanawi et al. <sup>17</sup> showed better knowledge, positive attitude, and good practice among females toward non-pharmacological preventive measures. This could be explained by the assumption that women were more apprehensive about the adverse effects of the vaccine than contracting COVID-19 <sup>18</sup>; however, in our study, we were unable to find any disparity in KAP scores in relation to the gender of the participants.

Individuals with higher knowledge about the disease and modes of transmission are associated with a more positive attitude and perception <sup>19</sup>. Participants with better knowledge about disease tend to have a superior attitude reflected in their better perceptions of preventive actions, resulting in active engagement in positive practices. Several previous KAP surveys performed for various infectious diseases reported identical associations <sup>19,20,21</sup>. A Chinese study demonstrated that higher education corresponds to better knowledge scores, but a similarly designed population survey of the Iranian population had varied conclusions <sup>22,23</sup>.

According to Ntontis E et al. <sup>24</sup>, poor knowledge, improper information, and deceit can result in hysteria and may cause panic buying. Such hysterical buying may break health supply chains as a shortage of sanitizers, masks, and essential drugs <sup>25</sup>. However, patients in our study did not witness any shortage of cardiac medications at district-level pharmacies and were not involved in stockpiling.

Beliefs about COVID-19 are acquired from variable sources such as public discussions, knowledge about similar viral diseases, governmental outreach programs, social and print media, community experiences, and healthcare sources. The factuality of these beliefs and hence, knowledge determine the attitude and thus the practices for prevention of COVID-19 infection. It varies significantly in the population depending upon age, place of residence, and education, as shown by our study<sup>4</sup>.

The awareness campaign must be designed to reach people of all age groups equally and effectively, irrespective of their education status and place of residence. For instance, visual depiction of guidelines and awareness through audio campaigns may help target people with varying literacy levels. Social fabric among the community produces everlasting interpersonal bonds, which nurtures empathy and a sense of caring for others <sup>26</sup>.

The limitation of our study was a smaller sample size, which could be justified by the fact that the study group was exclusive, and the number of elective cardiac surgeries declined during the lockdown. Regression analysis was not done to establish whether the level of knowledge corroborated with the attitude and practices in each subgroup.

## Conclusion

This single-center KAP study on COVID-19 among cardiac patients who are considered high risk was able to postulate a comprehensive evaluation. The findings suggest that high-risk patients have decent knowledge scores and an overall optimistic outlook on surviving the pandemic. However, novel and far-reaching awareness programs are required for patients who are less educated and reside in rural areas as they are found to have lower KAP scores. Keeping in mind the probability of upcoming waves in the near

future, consistent enforcement of preventive measures from healthcare agencies is crucial to surviving this worldwide pandemic.

## References

1. Ghosh A, Nundy S, Mallick TK. How India is dealing with the COVID-19 pandemic. *Sensors International*. 2020;1(July):100021. *Sensors International* 1 (2020) 100021
2. Khalil A, Abdalrahim M. Knowledge, attitudes, and practices towards prevention and early detection of chronic kidney disease. *International Nursing Review* 2014. 61, 237–245
3. Maheshwari S, Gupta P, Sinha R, Rawat P. Knowledge, attitude, and practice towards coronavirus disease 2019 (COVID-19) among medical students: A cross-sectional study. *Journal of Acute Disease*. 2020;9(3):100.
4. Christy JS, Kaur K, Gurnani B, Hess OM, Narendran K, Venugopal A, et al. Knowledge, attitude and practice toward COVID-19 among patients presenting to five tertiary eye care hospitals in South India-A multicentre questionnaire-based survey. *Indian Journal of Ophthalmology*. 2020 Nov 1;68(11):2385–90.
5. Ali SH, Foreman J, Tozan Y, Capasso A, Jones AM, Diclemte RJ. Trends and Predictors of COVID-19 Information Sources and Their Relationship With Knowledge and Beliefs Related to the Pandemic : Nationwide Cross-Sectional Study. *JMIR Public Health Surveill* 2020;6(4):e21071
6. Geana M V, Allen W. Kansans in the Middle of the Pandemic: Risk Perception, Knowledge, Compliance with Preventive Measures, and Primary Sources of Information about COVID-19 [Internet]. *Kans J Med* 2020;13:160-164
7. Myers C, Hamilton JF, Myers C, Hamilton JF. Social Media as Primary Source The narrativization of twenty-first-century social movements. *Media History* 2014Vol. 20, No. 4, 431–444
8. Quintana-salcedo Á, Vargas-ortiz L, Severiche-sierra C, Castro-bocanegra V. Knowledge and Attitudes of Health Workers for the Care of Patients COVID-19 in some cities of the Colombian Caribbean. *Indian Journal of Public Health Research & Development*, August 2020, Vol. 11, No. 8
9. Narayanaswamy DA, Meenakshi DK, Porchelvan DS, Vinoth Kumar DJ, Malathi M v, Ragini ML, et al. Knowledge, Attitude And Practices of Cardiac patients during the COVID19 pandemic in a Tertiary Care Medical College Hospital. *Annals of Tropical Medicine & Public Health*. 2021;23(23).
10. Pal R, Yadav U, Grover S, Saboo B, Verma A, Bhadada SK. Knowledge, attitudes, and practices towards COVID-19 among young adults with Type 1 Diabetes Mellitus amid the nationwide lockdown in India: A cross-sectional survey. *Diabetes Research and Clinical Practice* [Internet]. 2020 Aug;166:108344.
11. Bates BR, Moncayo AL, Costales JA, Herrera CA, Grijalva MJ. Knowledge, Attitudes, and Practices Towards COVID-19 Among Ecuadorians During the Outbreak: An Online Cross-Sectional Survey. *Journal of Community Health* [Internet]. 2020;(0123456789).
12. Clements JM, Street E. Knowledge and Behaviors Toward COVID-19 Among US Residents During the Early Days of the Pandemic : Cross-Sectional Online Questionnaire *JMIR Public Health Surveill* 2020;6(2):e19161
13. Glenn ND, Hill L, Norval JR, Glenn D. Rural-Urban Differences in Attitudes and Behavior in the United States. *ANNALS, AAPSS*, 429. Jan 1977
14. Ferdous MZ, Islam MS, Sikder MT, Mosaddek ASM, Zegarra-Valdivia JA, Gozal D. Knowledge, attitude, and practice regarding COVID-19 outbreak in Bangladesh: An online-based cross-sectional study. *PLoS ONE* [Internet]. 2020;15(10 October):1–17.
15. Lee M, Kang B ah, You M. Knowledge, attitudes, and practices ( KAP ) toward COVID-19 : a cross-sectional study in South Korea. *BMC Public Health* (2021) 21:295

16. Tomar BS, Singh P, Suman S, Raj P, Nathiya D. Indian community ' s Knowledge, Attitude & Practice towards. MedRxiv 2020.05.05.20092122
17. Al-hanawi MK, Angawi K, Al shareef N, Qattan AMN, Helmy HZ, Abudawood Y et al. Knowledge, Attitude and Practice Toward COVID-19 Among the Public in the Kingdom of Saudi Arabia : A Cross-Sectional Study. Front. Public Health 2020, 8:217
18. Neumann S, Nirosha B, Varghese E, Sabat I, Pita P, Werner B. Once we have it, will we use it ? A European survey on willingness to be vaccinated against COVID-19. The European Journal of Health Economics [Internet]. 2020;21(7):977–82.
19. Papagiannis D, Malli F, Raptis DG, Papathanasiou I. Assessment of Knowledge, Attitudes, and Practices towards New Coronavirus ( SARS-CoV-2 ) of Health Care Professionals in Greece before the Outbreak Period. Int. J. Environ. Res. Public Health 2020,17, 4925
20. Haq N, Hassali MA, Shafie AA, Saleem F, Farooqui M. A cross-sectional assessment of knowledge, attitude and practice towards Hepatitis B among healthy population of Quetta, Pakistan. BMC Public Health [Internet]. 2012;12(1):1.
21. Ajilore, Kolade et al. “College students’ knowledge, attitudes and adherence to public service announcements on Ebola in Nigeria: Suggestions for improving future Ebola prevention education programmes.” Health Education Journal 76 (2017): 648 - 660.
22. Zhong BL, Luo W, Li HM, Zhang QQ, Liu XG, Li WT, et al. Knowledge, attitudes, and practices towards COVID-19 among Chinese residents during the rapid rise period of the COVID-19 outbreak: A quick online cross-sectional survey. International Journal of Biological Sciences. 2020;16(10):1745–52.
23. Erfani A, Shahriarirad R, Ranjbhar K, Mirahmadizadeh A, Moghadhami M. Knowledge, Attitude and Practice toward the Novel Coronavirus(COVID-19) outbreak: A population based survey in Iran: Bulletin of world health organization. 30 march 2020
24. Ntontis E, Vestergren S, Saavedra P, Neville F, Jurstakova K, Cocking C, et al. Is it really “panic buying”? Public perceptions and experiences of extra buying at the onset of the COVID-19 pandemic. PLoS ONE 17(2):e0264618.
25. Sharma A, Gupta P, Jha R. COVID-19 : Impact on Health Supply Chain and Lessons to Be Learnt. Journal of Health Management 22(2) 248–261, 2020
26. Quandt SA, Lamonto NJ, Mora DC, Talton JW, Laurienti PJ, Arcury TA. COVID-19 Pandemic among Latinx Farmworker and Non farmworker Families in North Carolina : Knowledge, Risk Perceptions, and Preventive Behaviors. Int. J. Environ. Res. Public Health 2020,17, 5786.

#### Hosted file

Table 1 Questionnaire.docx available at <https://authorea.com/users/494398/articles/576478-knowledge-attitude-and-practices-among-cardiac-surgical-patients-towards-coronavirus-disease-covid-19-at-tertiary-care-center>

#### Hosted file

Table 2 Demography.docx available at <https://authorea.com/users/494398/articles/576478-knowledge-attitude-and-practices-among-cardiac-surgical-patients-towards-coronavirus-disease-covid-19-at-tertiary-care-center>

#### Hosted file

Table 3.docx available at <https://authorea.com/users/494398/articles/576478-knowledge-attitude-and-practices-among-cardiac-surgical-patients-towards-coronavirus-disease-covid-19-at-tertiary-care-center>

### Hosted file

Table 4.docx available at <https://authorea.com/users/494398/articles/576478-knowledge-attitude-and-practices-among-cardiac-surgical-patients-towards-coronavirus-disease-covid-19-at-tertiary-care-center>

### Hosted file

Table 5.docx available at <https://authorea.com/users/494398/articles/576478-knowledge-attitude-and-practices-among-cardiac-surgical-patients-towards-coronavirus-disease-covid-19-at-tertiary-care-center>

### Hosted file

Table 6.docx available at <https://authorea.com/users/494398/articles/576478-knowledge-attitude-and-practices-among-cardiac-surgical-patients-towards-coronavirus-disease-covid-19-at-tertiary-care-center>

### Hosted file

Table 7.docx available at <https://authorea.com/users/494398/articles/576478-knowledge-attitude-and-practices-among-cardiac-surgical-patients-towards-coronavirus-disease-covid-19-at-tertiary-care-center>