

Change point analysis to detect the effect of pruning severity on tree growth

Yutaka Iguchi¹

¹Laboratory of Biology

November 15, 2023

Abstract

The effect of pruning severity on tree growth was analyzed by change point detection using segmented regression. The present study applied this analysis to a well-known published data set including diameter growth response, tree age, pruning severity and pretreatment crown size. First, multiple regression analysis was performed to assess the effect of tree age, pruning severity and pretreatment crown size on diameter growth response. Next, segmented regression analysis was performed to assess the effect of pruning severity on diameter growth response. The results of the multiple regression showed that diameter growth response was significantly influenced by pruning severity and pretreatment crown size. The results of the segmented regression showed that in the whole data set, an abrupt change toward a decrease in diameter growth response was detected at 25% of the live crown removed. However, in the group of fully crowned and open-grown, diameter growth response continuously decreased with increasing pruning severity with no significant abrupt change, whereas in the group of 70-90% live crown, diameter growth response did not significantly decrease up to the break point (53% crown removed) and then abruptly decreased. This may be the first study to show the numerical evaluation of the effect of pruning severity on tree growth by change point analysis.

Yutaka Iguchi*

Laboratory of Biology, Yamashita-cho 1-10-6, Okaya City, Nagano Prefecture, 394-0005, Japan

* Correspondence:

Corresponding Author

bio-igu@f8.dion.ne.jp

1. Introduction

Tree pruning aims to craft canopy structure and shape by removing and shortening branches and encouraging growth in selected areas of the crown (Gilman et al., 2006). However, there has been a long-standing controversial issue of how pruning severity affects tree growth (Clark & Matheny, 2010). Pruning can negatively affect growth through excessive intensity (Rais et al., 2020). Therefore, most pruning prescriptions are based on empirical data combining operational needs with tree growth responses (Maurin & DesRochers, 2013; Shimada, 2017) or based on previous pruning studies. For example, O'Hara (1991) has been frequently cited as a well-known review suggesting that one-third of the live crown could be pruned without serious growth impact (Robbins, 2000; Clark & Matheny, 2010; Rais et al. 2020; Suchocka et al., 2021). However, O'Hara (1991) did not show how the removal limit of one-third was estimated.

The present article proposes a change point analysis to detect the effect of pruning severity on tree growth. Regarding O'Hara's (1991) estimation, it seems plausible that there exists an abrupt change point in the relationship between pruning severity and tree growth. Therefore, the present article aims to introduce a segmented regression model as a tool of the detection of an abrupt change point (Muggeo, 2008) and apply it to the data of O'Hara (1991).

Segmented regression models (also called broken-line models) are regression models where the relationships between the response and one or more explanatory variables are piecewise linear, namely represented by two or more straight lines connected at unknown values (Muggeo, 2008). Change point analysis using segmented regression models has been rarely reported in arboriculture and its related fields except for Hilbert et al. (2022). Therefore, the use of segmented regression and the results shown here may help further studies explore the relationship between pruning severity and tree growth.

As mentioned by O'Hara (1991) as well as other studies (Pothier et al., 2013, Shimada, 2023), the growth of diameter or circumference at breast height is considered an index of tree vigor. Therefore, the present article explores the relationship between diameter growth response and other measurement variables shown in O'Hara (1991).

2. Materials and Methods

2.1. Data and Analysis

The data analyzed here were obtained from Table 1 of O'Hara (1991). Some pruning severity values (% crown removed) were shown as the interval such as 15 – 35. In this case, the mid-value of the interval such as 25 was used for computational convenience.

In the following sections, a multiple regression analysis is first performed to examine the influence of the tree age, pruning severity and pretreatment crown size on diameter growth response. Next, a segmented regression model is employed to detect a change point in the relationships between pruning severity and growth response diameter.

Statistical analysis was performed using the stats, car, and segment packages in the R software (R Core Team, 2023) at a significance level of 0.05.

2.2. Multiple Regression Analysis

In the multiple regression analysis, the response variable was diameter growth response, and the explanatory variables were tree age, pruning severity (percent of crown removed), and pretreatment crown size. There were missing data in the Table 1 of O'Hara (1991). Therefore, multiple imputation was used to fill in the missing data. The calculation was performed using the function mice in the mice package and the function lm in the stats package.

The variable pretreatment crown size was defined as a categorical variable. The category names were derived from the Table 1 of O'Hara (1991): 70% live crown, 80% live crown, 90% live crown, fully crowned, and open-grown. The 70% live crown category served as the reference category.

2.3 Segmented Regression Analysis

In the segmented regression analysis, the response variable was diameter growth response, and the explanatory variable was pruning severity. The analysis was performed using the function segmented in the segmented package. In order to avoid multiple values of the response variable for one value of the explanatory variable, small random numbers with the range $[-0.01, 0.01]$ were added to the explanatory variable.

The random numbers were created by the function `runif` in the `stats` package. Score test by the function `pscore.test` in the `segmented` package was also used to test for the existence of a breakpoint.

3. Results

3.1. Results of Multiple Regression Analysis

The results of the multiple regression analysis are shown in Table 1. Diameter growth response was significantly influenced by pruning severity and pretreatment crown size. In the categories of pretreatment crown size, the fully crowned category significantly influenced diameter growth response and the open-grown category was almost significant.

Variable	Regression Coefficient	SE	t value	df	p value
Pruning Severity	-0.72	0.08	-8.07	23.1	0.0
Age	-0.09	0.27	-0.32	16.5	0.76
Pretreatment Crown Size					
80% Live Crown	-8.65	10.93	-0.79	22.2	0.44
90% Live Crown	-5.7	10.07	-0.57	23.1	0.58
Fully Crowned	-13.79	5.74	-2.4	22.3	0.03
Open-grown	-14.91	7.76	-1.92	22.8	0.07

Table 1: Multiple regression analysis showing the predictors of diameter growth response.

3.2. Results of Segmented Regression Analysis

First, a segmented regression was applied to all the data sets of pruning severity and diameter growth response. The results are depicted in Figure 1. A breakpoint was found at 25% crown removed and almost significantly supported by score test ($p = 0.07$). The first segmented line ($\beta = -0.25$, $0 < \text{severity} < 25$) was not significant ($p = 0.57$).




Figure 1: Segmented regression of diameter growth response on pruning severity. The data were obtained from the Table 1 of O'Hara (1991).

Second, based on the results of the multiple regression analysis, the data were divided into two groups regarding pretreatment crown size: one consists of fully crowned and open-grown, and the other consists of 70% live crown, 80% live crown, and 90% live crown. The results of the segmented regression are depicted in Figure 2.

In the group of fully crowned and open-grown, a breakpoint was found at 67% crown removed, but not significantly supported by score test ($p = 0.78$). The first segmented line ($\beta = -0.81$, $0 < \text{severity} < 67$) was significantly negative ($p < 0.001$) (the left panel of Figure 2).

On the other hand, in the group of 70-90% live crown, a breakpoint was found at 53% crown removed and significantly supported by score test ($p < 0.001$). The first segmented line ($\beta = -0.16$, $0 < \text{severity} < 53$) was not significant ($p = 0.14$) (the right panel of Figure 2).




Figure 2: Segmented regression of diameter growth response on pruning severity in the two groups based on pretreatment crown size. The data were obtained from the Table 1 of O'Hara (1991).

4. Discussion

Several previous studies have referred to O'Hara (1991) and explained that the removal of up to one-third, or 33% of the live crown did not noticeably reduce diameter growth (Robbins, 2000; Clark & Matheny, 2010; Kirby, 2016; Rais et al. 2020; Suchocka et al., 2021). Nevertheless, there remained the question of how the limit value of 33% was obtained by calculation.

The present study applied segmented regression to the data of O'Hara (1991) and revealed that in the whole data set, an abrupt change toward a decrease in diameter growth response was detected at 25% of the live crown removed. This value was a little smaller than the limit value of 33% estimated by O'Hara (1991) and cited by the above-mentioned studies.

O'Hara (1991) mentioned the effect of different pretreatments on tree growth, but did not take it into consideration when evaluating the crown removal limit. However, the present results revealed that the different pretreatments produced different relationships between pruning severity and growth response diameter.

In the group of fully crowned and open-grown, diameter growth response continuously decreased with increasing pruning severity with no significant abrupt change. Therefore, it seemed difficult to determine the crown removal limit. However, in the group of 70-90% live crown, diameter growth response did not significantly decrease up to the break point (53% crown removed) and then abruptly decreased. Consequently, the difference in pretreatments was found to have a great impact on the relationships between pruning severity and growth response diameter.

5. Conclusion

This is probably the first study that evaluated the effect of pruning severity on tree growth by change point analysis using segmented regression. The advantage of segmented regression is that it enables us to analyze continuous data as they are. In other words, segmented regression does not treat continuous predictors such as pruning severity as categorical predictors. Therefore, the present study with segmented regression was able to analyze the relationship between pruning severity and growth response diameter to estimate the limit

value up to which pruning can be done without serious growth losses. The results will recommend the use of change point analysis in future studies on pruning severity.

Acknowledgements

The author declares that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest. The author confirms sole responsibility for the following: study conception and design, data collection, analysis and interpretation of results, and manuscript preparation.

References

- Clark, J. R., & Matheny, N. (2010). The Research Foundation to Tree Pruning: A Review of the Literature. *Arboriculture & Urban Forestry*, 36, 110-120. <https://doi.org/10.48044/jauf.2010.015>
- Gilman, E. F., Anderson, P. J., & Harchick, C. (2006). Pruning Lower Branches of Live Oak (*Quercus virginiana* Mill.) Cultivars and Seedlings during Nursery Production: Balancing Growth and Efficiency. *Journal of Environmental Horticulture*, 24, 201-206. <https://doi.org/10.24266/0738-2898-24.4.201>
- Hilbert, D. R., Koeser, A. K., Roman, L. A., Andreu, M. G., Hansen, G., Thetford, M., & Northrop, R. J. (2022). Selecting and Assessing Underutilized Trees for Diverse Urban Forests: A Participatory Research Approach. *Frontiers in Ecology and Evolution*, 10, 759693. <https://doi.org/10.3389/fevo.2022.759693>
- Kirby, J. W. (2016). Stem and Branch Diameter Response in Pruned Douglas-Fir Plantations (*Pseudotsuga menziesii* var. *menziesii*): Implications for Volume and Clear Wood Production in the US Pacific Northwest. Master's Thesis, Seattle: University of Washington.
- Maurin, V., & DesRochers, A. (2013). Physiological and Growth Responses to Pruning Season and Intensity of Hybrid Poplar. *Forest Ecology and Management*, 304, 399-406. <https://doi.org/10.1016/j.foreco.2013.05.039>
- Muggeo, V. M. R. (2008). Segmented: An R Package to Fit Regression Models with Broken-Line Relationships. *R News*, 8, 20-25
- O'Hara, K. L. (1991). Technical Commentary: A Biological Justification for Pruning in Coastal Douglas-Fir Stands. *Western Journal of Applied Forestry*, 6, 59-63. <https://doi.org/10.1093/wjaf/6.3.59>
- Pothier, D., Fortin, M., Auty, D., Delisle-Boulianne, S., Gagné, L. V., & Achim, A. (2013). Improving Tree Selection for Partial Cutting Through Joint Probability Modelling of Tree Vigor and Quality. *Canadian Journal of Forest Research*, 43, 288-298. <https://doi.org/10.1139/cjfr-2012-0402>
- R Core Team (2023). R: A Language and Environment for Statistical Computing. R Foundation for Statistical Computing, Vienna.
- Rais, A., Poschenrieder, W., van de Kuilen, J. W. G., & Pretzsch, H. (2020). Impact of Spacing and Pruning on Quantity, Quality and Economics of Douglas-Fir Sawn Timber: Scenario and Sensitivity Analysis. *European Journal of Forest Research*, 139, 747-758. <https://doi.org/10.1007/s10342-020-01282-8>
- Robbins, J. M. (2000). Influence of Spacing and Crown Recession on Wood Quality of Intensively-Managed Young Growth Douglas-Fir. Master's Thesis, Corvallis: Oregon State University.
- Shimada, H. (2017). Maintenance of Trees in a Residential Section Utilizing Pruning Method under Retaining the Apical of the Main Branch of Crown. *Tree and Forest Health*, 21, 127-131. https://doi.org/10.18938/treeforesthealth.21.3_127

Shimada, H., & Hosono, T. (2023). Effects of Pruning on the Growth of Bamboo-leaf Oak (*Quercus myrsinifolia*): A Comparison between Crown Reduction and Leader Retention Methods. *Tree and Forest Health*, 27, 1-12.

Suchocka, M., Swoczyna, T., Kosno-Jończy, J., & Kalaji, H. M. (2021). Impact of Heavy Pruning on Development and Photosynthesis of *Tilia cordata* Mill. *Trees*. *PLoS ONE*, 16, e0256465. <https://doi.org/10.1371/journal.pone.0256465>