

Manuscript submitted to the Journal of Geophysical Research – Oceans

Cascading weather events amplify the coastal thermal conditions prior to the shelf transit of Hurricane Sally (2020)

B. Dzwonkowski^{1,2*}, S. Fournier³, G. Lockridge², J. Coogan⁴, Z. Liu¹, and K. Park⁵

¹University of South Alabama, Mobile, AL, 36688, U.S.A.

²Dauphin Island Sea Lab, Dauphin Island, AL 36528, U.S.A.

³Jet Propulsion Laboratory, California Institute of Technology, Pasadena, CA 91109, U.S.A.

⁴Woods Hole Oceanographic Institution, Woods Hole, MA 02543, U.S.A.

⁵Texas A&M University at Galveston, Galveston, TX 77554, U.S.A.

*briandz@disl.org

Abstract

Changes in tropical cyclone intensity prior to landfall represent a significant risk to human life and coastal infrastructure. Such changes can be influenced by shelf water temperatures through their role in mediating heat exchange between the ocean and atmosphere. However, the evolution of shelf sea surface temperature during a storm is dependent on the initial thermal conditions of the water column, information that is often unavailable. Here, observational data from multiple monitoring stations and satellite sensors were used to identify the sequence of events that led to the development of storm-favorable thermal conditions in the Mississippi Bight prior to the transit of Hurricane Sally (2020), a storm that rapidly intensified over the shelf. The annual peak in depth-average temperature of $>29^{\circ}\text{C}$ that occurred prior to the arrival of Hurricane Sally was the result of two distinct warming periods caused by a cascade of weather events. The event sequence transitioned the system from below average to above average thermal conditions over a 25-d period. The transition was initiated with the passage of Hurricane Marco (2020), which mixed the upper water column, transferring heat downward and minimizing the cold bottom water reserved over the shelf. The subsequent reheating of the upper ocean by a positive surface heat flux, followed by a period of downwelling winds, effectively elevated shelf-wide thermal conditions for the subsequent storm. The climatological coupling of warm sea surface temperature and downwelling winds suggest regions with such characteristics are at an elevated risk for storm intensification over the shelf.

Plain Language Abstract:

Changes in tropical cyclone intensity prior to landfall represent a significant risk to human life and coastal infrastructure. Such changes can be influenced by sea surface temperatures as they provide the energy to tropical cyclones. However, the changes in sea surface temperatures during a storm depend on heat content in the coastal ocean before the storm arrives and the relevant information is

often unavailable. Measurements from multiple monitoring stations and satellite sensors were used to identify the events that led to the development of favorable temperatures in the Mississippi Bight prior to the arrival of Hurricane Sally (2020), a storm that rapidly intensified in the coastal ocean. A sequence of two warming periods transitioned the coastal ocean from below average to above average temperatures over a 25-d period. The transition was initiated by Hurricane Marco (2020) which pumped heat from the surface of the ocean to the bottom. This was followed by a reheating of the upper ocean and the offshore transport of warm coastal waters which packed the shelf with thermal energy that could feed the subsequent storm. The results suggest that coastlines with certain ocean and wind conditions may have an elevated risk of storm strengthening before landfall.

Key Points:

- Annual peak in depth-average temperature occurred prior to shelf transit of Hurricane Sally providing favorable intensification conditions
- Depth-average temperature was set up by a distinct cascade of weather events including persistent downwelling conditions
- Climatological coupling of ocean temperature and wind conditions suggest an elevated risk of storm intensification risk over the shelf

Key Words: tropical cyclones, coastal ocean, cascading events, temperature, downwelling, Hurricane Sally

1. Introduction

Landfall tropical cyclones are extreme events that represent a major threat to coastal regions across large swaths of the globe (Emanuel 2005; Keim et al. 2007). An important factor limiting society’s ability to mitigate loss of property and life centers around storm intensity, where forecasting progress has been slow relative to storm track (Kaplan and DeMaria 2003; Kaplan 2010; DeMaria et al. 2014). Limitations in intensity forecasting are particularly important as storms approach shorelines where planning and response decisions have direct effects on the impacts of these events. Overcoming difficulties in intensity forecasting will have added importance in a warming future climate where storms are expected to be stronger and have increased precipitation rates (e.g., Bacmeister et al. 2018; Liu et al. 2019; Vecchi et al. 2019; Knutson et al. 2020; Chu et al. 2020). This appears to be particularly concerning in coastal regions as Chu et al. (2020) found there was a 150% increase in the expected number of category 3-5 storms making landfall under a carbon dioxide doubling scenario (but no additional increase for a quadrupling scenario). Furthermore, Emanuel (2017) highlighted that storms may intensify more rapidly before landfall under future warming climate scenarios, potentially making intensity prediction more challenging.

Landfall storm intensity is connected to coastal ocean conditions through sea surface temperature. The importance of this connection is highlighted by incorporation of sea surface temperature into the intensity potential parameter

developed by Emanuel (1999) which indicates that the larger the difference between the sea surface and upper atmosphere temperatures, the stronger the potential wind speed. As a result, sea surface temperature modifications during a storm event have the potential to influence the evolution of a tropical cyclone. Such changes in sea surface temperature occur through a combination of vertical mixing, horizontal advection, upwelling and surface heat flux (Potter et al. 2017). While there can be significant storm to storm variability in the relative impacts of these four mechanisms (e.g., Potter et al. 2017), vertical mixing is generally found to be the dominant driver of upper ocean cooling (Price 2009). Thus, this mixing potential implies that pre-storm thermal stratification (or lack thereof) has a significant impact on storm intensity at landfall. Shelves with high thermal stratification prior to storm passage have been linked to significant ahead-of-eye cooling and thus storm attenuation (Glenn et al. 2016; Seroka et al. 2016; Park et al. 2019). In contrast, homogeneous shelves have been shown to support sustained intensities or even intensification (Miles et al. 2017; Potter et al. 2019; Pun et al. 2019; Zhang et al. 2019; Dzwonkowski et al. 2020). Consequently, the initial shelf conditions over which air-sea interactions operate can significantly impact storm evolution prior to landfall.

The sequence of events that lead to particularly favorable or unfavorable shelf conditions can be challenging to assess because the importance of any particular storm event and its associated shelf crossing trajectory are extremely difficult to predict. As highlighted by Dzwonkowski et al. (2020), the sequence of events that setup the extreme shelf heat content prior to Hurricane Michael (2018) began nearly a month prior to the shelf intensification of this storm. Thus, planned data collections of such preconditioning are difficult, requiring most observational studies be based on platforms of opportunity. This issue is further exacerbated by the sparse availability of real-time subsurface information on shelf conditions, limiting the ability of forecasters to understand changes in sea surface temperature prior to landfall. Thus, developing a better understanding of the event sequence that drives conditions favorable to storm intensification is critical to forecasting landfall intensity of tropical cyclones. Furthermore, an enhanced understanding of event sequencing could also be used to better characterize the susceptibility of individual shelf regions to risks posed by tropical cyclones; for example, contributing event distributions could be used to develop co-occurrence probabilities.

The intense Atlantic storm activity of 2020, with 8 landfall tropical cyclones in the Gulf of Mexico, provided multiple potential opportunities to improve the understanding of shelf setup processes. Of the storms entering the Gulf of Mexico, Hurricane Sally transited the Mississippi Bight and made landfall in Orange Beach, Florida on the morning of September 16 (Berg and Reinhart 2021). While Hurricane Sally only achieved the status of a category 2 storm, the storm was only 1 knot below the wind speed required for classification as a major hurricane (Category 3-5). More importantly, it was a slow-moving storm (mean shelf transit speed of $\sim 8 \text{ km h}^{-1}$) that rapidly intensified over the shelf (20 knots over 12 h) leading to an extended period of high wind and waves in the

region. The specific coastal ocean response to the passage of Hurricane Sally is the focus of another complimentary manuscript under preparation.

Here, we focus on the causal chain of events that led to an amplification of the heat content over the shelf in the Mississippi Bight prior to the arrival of Hurricane Sally. Similar to the work of Dzwonkowski et al. (2020) that focused on hurricane intensification from the summer 2018, the findings of this study highlight that events nearly a month prior to the arrival of Hurricane Sally were critical to setting up the shelf conditions. In the cases of Hurricane Michael (Dzwonkowski et al. 2020) and Hurricane Sally (this study), the passage of a tropical cyclone initiated the sequence of events that generated favorable shelf conditions for intensification of a subsequent storm. However, the initial storm response as well as the subsequent details in this case study deviate in notable ways and the findings suggest a coupling of climatological patterns that could be expected to produce hurricane ‘friendly’ shelf regions.

2. Data and Analysis

The data sources, processing, and analysis follow similar methods as those in Dzwonkowski et al. (2018, 2020) with appropriately modified descriptions below.

2.1. Tropical cyclone data

Position data for Hurricanes Marco and Sally were acquired from the best track files for each storm as provided by National Oceanic and Atmospheric Administration (NOAA) National Hurricane Center (<ftp://ftp.nhc.noaa.gov/atcf/btk/>).

2.2. In-situ data and associated analysis

To understand the atmospheric and oceanic conditions on the shelf prior to the arrival of Hurricane Sally, field and reanalysis data from various sources were obtained. Standard meteorological data from two NOAA National Data Buoy Center (NDBC) stations were used (Fig. 1): buoy station 42012 offshore of Orange Beach, AL (ORB, https://www.ndbc.noaa.gov/station_page.php?station=42012) and DPIA1 on Dauphin Island, AL (https://www.ndbc.noaa.gov/station_page.php?station=dpia1). The meteorological data included air temperature, relative humidity/dew point temperature, and wind speed and direction. These stations were the closest measurements available to the main mooring site (CP) and have relatively long records (9 and 30 years at ORB and DPIA1, respectively). The sites are generally similar in magnitude and direction, however, there were significant differences during the period when Hurricane Sally was transiting the shelf. In addition, incoming solar radiation data from the Dauphin Island station of the Alabama Real-time Coastal Observing system were obtained for the study period (https://arcos.disl.org/stations/disl_stations?stationnew=628). This site is less than 500 m away from DPIA1, and both sites are collectively referred to as site DI for this study (Fig. 1). Additional key atmospheric variables, including solar radiation and long-wave radiation, were obtained from the National Centers for Environmental Prediction (NCEP)

North American Region Reanalysis (NARR) for the grid cell closest to site ORB (<https://www.esrl.noaa.gov/psd/data/gridded/data.narr.html>). The NARR outputs are on $1/3^\circ$ grid (~ 32 -km resolution) and were interpolated from the 3-h outputs to hourly to match the DI and ORB data.

The hydrographic data were primarily derived from a long-term mooring station (site CP) on the 20 m isobath to the west-southwest of Mobile Bay (Fig. 1). The site provides a relatively long-term (15 years over the 2005-2020 period) perspective of shelf conditions from a suite of instruments. The instrument suite has changed over time, but typically consists of two CTD instruments to measure temperature and salinity at the bottom (~ 0.4 m above bottom, mab or depth of ~ 19.8 m) and near-surface (~ 15.5 mab or depth of ~ 4.5 m), and 4-9 thermistors throughout the water column to measure thermal structure. Details of the mooring configurations and aspects of the data processing can be found in various studies (e.g., Tzeng et al. 2016; Coogan et al. 2019; Dzwonkowski et al. 2020). Similar to the summer of 2018, the late summer of 2020 featured an additional data stream from a CTD on a surface buoy (~ 100 m apart from site CP), providing data at ~ 19.6 mab (i.e., ~ 0.6 m below the surface). However, the extent of the data stream in 2020 was short-lived, reporting intermittently between July 31 and September 3. In order to fill the significant gaps in this surface measurement as well as provide a similar in-situ sea surface temperature measured in previous years, sea surface temperature data from ORB (~ 1 -2 m below the surface) were combined with those from site CP for available data (10 years in 2009-2020, no available data in 2018). These data are available at the NOAA National Center for Environmental Information (NCEI: see the Acknowledgements for citation links).

The heat content/flux calculations were carried out following typical procedures. For heat content, the depth-average temperature was used given the shelf focus of the study (Price 2009; Potter et al. 2019; Pun et al. 2019) and was calculated with water column temperature observations interpolated to a 1-m grid. While sea surface temperature is the means through which tropical cyclones interact with the ocean, this study focuses on depth-average temperature as averaging over the expected mixing depth of a storm has been suggested or shown to be the most relevant property for storm intensification (Balaguru et al. 2015, 2018), particularly over a shallow shelf (Price 2009; Potter et al. 2019). The sensible and latent heat fluxes were calculated with the TOGA-COARE algorithms (Fairall et al. 1996) similar to other studies in the region (Coogan et al. 2019; Dzwonkowski et al. 2020). Incoming solar radiation data from site DI compared well to the NARR reanalysis data at CP and were used for the subsequent analysis. Upward solar radiation as well as long-wave radiation were taken from the NARR data. At a few time points, the sum of the downward solar radiation from DI and the upward solar radiation term from NARR resulted in small negative values, which were fixed to zero. Conversion between relative humidity and dew point temperature was conducted following standard algorithms (<https://bmcnoldy.rsmas.miami.edu/Humidity.html>) (August 1828; Magnus 1844; Alduchov and Eskridge 1996). Note that the dew point temper-

ature data stream failed at ORB on September 15 prior to Hurricane Sally. As such, relative humidity at site DI was used in the heat flux calculations at site ORB thereafter. While no marine heatwaves were identified in this study period following the methods of Hobday et al. (2016), the 90th percentile was used as a reference point for extreme thermal conditions in the depth-average temperature time series.

Current velocity data throughout the water column were obtained from a Nortek Acoustic Doppler Waves and Current profiler (AWAC) at site CP. During the passage of Hurricane Sally, movement of the AWAC changed the pitch, roll, and heading of the instrument. Proprietary firmware from Nortek (Surge v1.15.03) was used to re-map the bins to account for the change in orientation during this event period which resulted in the loss of some near-surface bins during Hurricane Sally (bin ranges changed 1.5-17.0 mab to 1.0-14.5 mab on September 16). The resulting data were processed following the procedure in Dzwonkowski et al. (2014) where a 40-h low-pass Lanczos filter was used to highlight low-frequency circulation patterns. These data are available at the Dauphin Island Sea Lab Data Management Center (<https://www.disl.org/research/data-management-center>) and NOAA NCEI (see the Acknowledgements for citation links).

The hydrographic and current velocity data at site CP were used to determine several water column stability parameters similar to those analyzed in Dzwonkowski et al. (2018). Direct measurements of the bulk shear squared (bulk S^2), bulk buoyancy frequency squared (bulk N^2), bulk Richardson number (bulk Ri) were determined:

- (1)
- (2)
- (3)

where Δu and Δv are along and across-shelf velocity differences between the upper and lower layers, respectively; Δz is the distance between the layer velocities; g is gravitational acceleration; $\Delta \rho$ is the density difference between the layers; and ρ_0 is the average density. Richardson number values less than or equal to the critical value ($Ri_{cr} = 0.25$) are generally accepted to be indicative of mixing, however higher values ($0.25 < Ri < 1$) have been associated with mixing in some studies (e.g., Canuto et al. 2001). The bulk S^2 was calculated with an upper layer velocity at 17.0 mab and a lower layer velocity at 3.0 mab as these values provide a first order estimate of the shear. Similarly, density measurement at the near-surface and bottom CTD were used to determine bulk N^2 .

2.3. Satellite data and associated analysis

For determining the regional sea surface conditions, several satellite-based data sets were obtained. Sea surface temperature data were obtained from the Jet

Propulsion Laboratory (JPL) Multi-scale Ultra-high Resolution (MUR) sea surface temperature product (https://podaac.jpl.nasa.gov/Multi-scale_Ultra-high_Resolution_MUR-SST) which is available from 2003-present. This is a 1-km resolution product that blends 1-km infrared sensor data and 25-km microwave sensor data. A regional climatology for the northeastern Gulf of Mexico was developed using the available 17 years of data. The MUR sea surface temperature data were favored over the longer duration Optimal Interpolated sea surface temperature (OISST) product because of the shelf emphasis of the study region where higher spatial resolution was prioritized over the coarser resolution of the OISST product. In addition, Terra/MODIS L2 chlorophyll-a data (1 km resolution) from August 18 2020 were used to provide a qualitative view of riverine/estuarine waters over the shelf prior to the arrival of Hurricane Marco. Such qualitative use of chlorophyll-a data from standard ocean color algorithms has been common in many previous studies (e.g., Dzwonkowski and Yan 2005; Walker et al. 2005).

3. Results

3.1. Water column conditions

The importance of event sequencing can be seen in the evolution of the depth-average temperature, a proxy for shelf heat content, which experienced a dramatic increase over the inner shelf during the month prior to Hurricane Sally increasing from $<27^{\circ}\text{C}$ in mid-August to $>29.5^{\circ}\text{C}$, the 2020 maximum, by September 10-11 (Fig. 2a). The peak was rapidly attenuated in response to the shelf crossing of Hurricane Sally (September 14-16), which initiated the fall transition to cooler, less stratified shelf conditions. Comparing the 2020 depth-average temperature evolution to the climatological conditions showed a transition from below average to above average conditions. While this increase in depth-average temperature did not reach the threshold for a ‘marine heatwave’, the depth-average temperatures were well above values that have been statistically associated with rapidly intensifying tropical cyclones, e.g., $<28.5^{\circ}\text{C}$ per Kaplan and DeMaria (2003). Notably, this month-long period was marked by two episodes of warming (August 17-27 and September 5-11), both of which were associated with downwelling favorable wind conditions (Fig. 2b). The first of these events was generated by the passage of Hurricane Marco to the south of the Mississippi Bight (Fig. 1), while the second was associated with a more typical summer wind pattern that occurred prior to the arrival of Hurricane Sally on the shelf.

3.2. Response to Hurricane Marco

The first warming period (August 17-27), with an increase in depth-average temperature of $\sim 2.5^{\circ}\text{C}$, was associated with the passage of tropical cyclone Marco and occurred as the shelf was transitioning from upwelling to downwelling conditions (Fig. 2a,b). Despite the storm being well to the southwest of the Alabama shelf, the cyclonic winds associated with Marco produced strong downwelling conditions ($\sim 10\text{ m s}^{-1}$ westward wind on August 23-26 in Fig. 2b), which amplified the transition from upwelling to downwelling that began around August

18. During the initial transition, the water column velocity data showed generally weak currents in both the along and across-shelf directions (Fig. 2c,d). Similarly, the thermal structure of the water column at site CP showed relaxing isotherms sloping downward with time (Fig. 2e).

The impact of Hurricane Marco was clearly visible in both the velocity and temperature structure on August 22-26. Unsurprisingly, the strong downwelling winds produced a very strong downwelling circulation with depth-average along-shelf currents around 60 cm s^{-1} and opposing across-shelf flows in the near-surface (onshore) and bottom (offshore) on the order of 30 cm s^{-1} (Fig. 2c,d). This downwelling circulation generated a rapid thermal response on August 22-23 that began homogenizing the water column temperature to $\sim 29^\circ\text{C}$, a value slightly lower than the pre-storm surface temperature (Fig. 2e). Shortly thereafter (August 24), the surface cooled by $0.5\text{-}1^\circ\text{C}$ while the bottom continued to warm and then became completely mixed (August 26). Interestingly, these changes in the thermal structure resulted in no significant change in the depth-average temperature during this period (Fig. 2a).

The salinity data at site CP brought additional context to thermal response during the passage of Hurricane Marco (Fig. 2f). The initial salinity response to the downwelling wind from Hurricane Marco showed a freshening of both the surface and bottom at site CP, dropping about 4 and 6 psu, respectively, on August 24-25. On August 26, two days after Marco passed just south of the Mississippi River Delta, the surface and bottom salinity began converging, an indication of mixing, and remained uniform for approximately a day. At the same time, the across-shelf current was dramatically reduced to $\sim 5 \text{ cm s}^{-1}$ at both near-surface and near-bottom (Fig. 2d).

Following the shelf response to the passage of Hurricane Marco, the system rapidly re-stratified with surface to bottom salinity differences of 4 psu and increasing to 8 psu by September 1. This resulted in stronger salinity difference relative to the pre-Marco conditions (Fig. 2f). In contrast, the thermal stratification was much slower to recover with surface to bottom temperature differences nearing the pre-Marco conditions around September 3-4, which lasted only for a short period of time till September 5 (Fig. 2e). Importantly, the post-Marco thermal structure was significantly different with a much thicker upper warm layer. This was particularly evident in the mid-water column structure where the 28.5°C [26°C] isotherm was at around 14 m [17 m] depth on September 3 compared to around 7 m [13 m] depth during the mid-August upwelling conditions (Fig. 2e). Despite having nearly the same surface to bottom temperature difference there was a $\sim 1.3^\circ\text{C}$ increase (from $\sim 27^\circ\text{C}$ in mid-August to $\sim 28.3^\circ\text{C}$ on September 5) in depth-average temperature, resulting in a much more favorable thermal environment for the subsequent tropical cyclone (i.e., Hurricane Sally).

3.3. Pre-Hurricane Sally Response

This favorable thermal environment was provided with an additional boost from the second warming period on September 5-11, with an increase in depth-average

temperature of $\sim 1.3^\circ\text{C}$, which was also associated with downwelling favorable wind conditions (Fig. 2a,b). The system response to the downwelling wind forcing was consistent with expectations, i.e., westward along-shelf flow and opposing across-shelf flow in the surface and bottom (Fig. 2c,d). Water temperatures below 26°C at the bottom of site CP on September 3-5 were replaced by much warmer water (Fig. 2e) and the prolonged period of downwelling wind conditions eventually homogenized the water column by September 10. This was well before Hurricane Sally began to exert a significant influence over the shelf region. At this point (September 11), the warming trend in the depth-average temperature was arrested and remained around 29.4°C until the inner core of Hurricane Sally began approaching the shelf break on September 14 (Fig. 2a).

4. Discussion

As noted, there were two critical periods of increasing shelf heat content associated with downwelling events. In addition to the individual behaviors of these events, the time interval between them (August 28 - September 4) was also critical to the final pre-Sally shelf conditions. The key aspects of the processes driving the evolution of the thermal environment in the event sequence are discussed below.

4.1. Downward transfer of heat from mixing during Hurricane Marco

Given that site CP was over a 100 km north of the center of Hurricane Marco, the relative importance of this event to the shelf heat budget is difficult to assess with the available observations. As can be seen from the time series of the current velocity and temperature structure (Fig. 2c,d,e), the cold water that was advected offshore in the bottom boundary layer during Marco eventually returned by early September during the subsequent upwelling event. This suggests the advective impact of Marco on the shelf heat content was short lived. However, there was a much thicker upper warm layer relative to the prior upwelling period (September 4 vs. August 15), that suggests heat was transferred deeper into the water column, effectively storing thermal energy at depth for Hurricane Sally, due to mixing during Hurricane Marco.

To determine whether significant mixing was likely during the passage of Hurricane Marco, water column stability characteristics at site CP were examined as Hurricane Marco moved through the region (Fig. 3). Beginning on August 23, the bulk S^2 and bulk N^2 started to exhibit opposing trends with shear increasing and stratification decreasing with S^2 having notable diurnal peaks. Diurnal cycles in S^2 have been previously identified at this site and have been linked with near-inertial oscillations that modulate the low frequency structure of the horizontal velocities (Dzwonkowski et al. 2018). This interaction of the high and low frequencies generated periods of high and low shear (e.g., thin red and blue lines in Fig. 2d). At the same time, the low frequency downwelling flow was advecting the downwelling front offshore reducing the bulk N^2 (Fig. 3a). The resulting bulk Ri maintained values below 1 from August 24-26 with

several periods nearing the critical value of 0.25 (Fig. 3b). During this period of low bulk Ri , stratification was present but declined till August 26 when N^2 becomes zero, indicating the strong shear observed was generating mixing in the water column. This becomes more apparent on August 25 when the surface and bottom salinities converge to 29 psu.

Interestingly, the initial site response was quite different from what might be expected from a water column solely experiencing mixing (i.e., converging of surface and bottom mass properties) as both the surface and bottom salinities dropped synchronously (August 23, Fig. 2f). This highlights the role of downwelling advection in the system response to Hurricane Marco. In this case, satellite-derived ocean color imagery showed an extensive river plume surging across much of the Mississippi Bight on August 18 (Fig. 1). Thus, the onshore surface advection of this offshore buoyancy source coupled with the offshore transport of coastal water in the bottom boundary layer likely generated the synchronous drop in salinity. All of which was occurring in a highly sheared (relative to stratification) water column that would have been expected to erode the stratification through mixing.

Consequently, these data indicate that Hurricane Marco did induce mixing that transferred heat downward into the water column over the shelf. This is consistent with findings of Dzwonkowski et al. (2020) that showed a similar downward transfer of heat when Tropical Storm Gordon passed through the Mississippi Bight region approximately a month prior to Hurricane Michael, another storm that intensified while crossing the shelf. In the case of Hurricane Marco, the return of cool bottom temperatures to site CP indicated that the shelf was not ‘cleared’ of its cold water reserve ($<26^\circ\text{C}$ on September 3-5 in Fig. 2e). This may, in part, have been due to the presence of the extensive surface river plume emanating from the Mississippi River Delta, which would limit the mixing depth generated from the winds of Hurricane Marco. While buoyancy supplied by river plumes support hurricane strengthening in the open ocean (Reul et al. 2015; Balaguru et al. 2020; Domingues et al. 2021), the river discharge in this case may have indirectly inhibited the strengthening of a subsequent storm (Hurricane Sally) by limiting the amount of heat that was mixed to depth over the shelf by Hurricane Marco.

4.2. Post-Hurricane Marco thermal recovery

The next component of this event sequence was a re-warming of the upper ocean after Hurricane Marco as a result of favorable atmospheric conditions. Between August 25 and September 4, the region experienced positive net daily surface heat flux with the exception of August 30 (Fig. 2g) which allowed the depth-average temperature to increase slightly immediately after the storm and maintained a temperature around 29°C for approximately a week between August 27 and September 2 (Fig. 2a). While the full water column initially warms in response to the surface heat flux, over time the thermal structure showed a clear divergence with the upper water column re-warming and the lower water column cooling. Thus, the post-Marco positive net surface heat flux

sustained the depth-average temperature over the inner shelf despite upwelling conditions advecting cooler bottom water that was displaced offshore during the previous downwelling condition (i.e., winds associated with Hurricane Marco). Importantly, the warm surface layer extended to a much greater depth relative to the pre-Marco conditions. This response is also largely consistent with the findings of Dzwonkowski et al. (2020) that noted the breakdown of stratification was critical in the rapid rewarming of the upper water column and was a key component of the compounding processes that drove a coastal marine heatwave in 2018 before landfall of Hurricane Michael. It should be noted that eventually the cooler bottom water associated with the upwelling did reduce the shelf heat content, dropping the depth-average temperature by approximately 0.5°C on September 4-5 (Fig. 2a). Nonetheless, the depth-average temperature at site CP was still notably higher than the pre-Marco conditions and both contributing factors (i.e., vertical mixing and rewarming) would be expected to have occurred over the broader shelf region significantly increasing the heat content across the Mississippi Bight region.

4.3. Impact of the pre-Sally downwelling

This increase in depth-average temperature across the shelf was amplified by a final downwelling event leading up to the arrival of Hurricane Sally. This persistent downwelling event effectively cleared the inner shelf of any cool water (Fig. 2e) and would be expected to continue advecting the pool of cooler bottom shelf water well beyond the mooring location. Such cold pool behavior was captured in mid-Atlantic Bight with coupled glider observations and numerical modeling outputs during Hurricane Sandy in 2012 (Miles et al. 2017). The cold pool advection was shown to be consistent with the scaling arguments of Austin and Lentz (2002), where frontal distance offshore (x_{ek}) could be represented as:

$$x_{\text{ek}}(t) = X_o + \sqrt{\int_0^t \frac{2*U_s}{\alpha} dt} \quad (4)$$

for events beyond an inertial period where t is time, X_o is the initial position of the front, α is shelf slope, and U_s is Ekman surface transport $= \tau_s / (\rho_o \cdot f)$ where τ_s is along-shelf wind stress, ρ_o is a reference density, and f is the Coriolis parameter. Applying this scaling to the period of time prior to the arrival of Hurricane Sally when the 28.5°C contour, a water temperature associated with rapid intensification, was observed to be passing by site CP (~September 9) suggests that the edge of the cold pool would be ~48 km south of the CP/ORB mooring locations (64 km offshore), putting the foot of the 28.5°C contour around the 40 m isobath, more than half the distance to the shelf break. This is consistent with the sea surface temperature data on September 14 (not shown), which showed the broader shelf area being above 28.5°C . Thus, this sequence of events leading up to the arrival of Hurricane Sally, i.e., the erosion of cooler bottom water from Hurricane Marco coupled with the subsequent offshore advection of the remaining bottom water, likely created a very large region of the shelf packed with very warm water. Similar such pre-storm thermal conditioning on the Texas shelf was associated with the intensification of Hurricane Harvey in

2017 (Potter et al. 2019).

4.4. Climatology coupling of temperature and wind

Given that both warm upper ocean temperatures and downwelling favorable winds were primary drivers in amplifying the coastal ocean heat content, the likelihood of their co-occurrence during this period of peak hurricane activity represents a potentially important climatological coupling. Long-term observations of wind stress from both ORB (9-year) and DI (30 years) indicated downwelling winds dominated from mid-August to mid-September in this region (Fig. 4). While there was significant interannual variability in the strength of the wind stress, the mean of every year during this active storm period (August 15- September 14) was downwelling favorable. The relatively shorter time series at site ORB suggested that the 2020 study period was associated with anomalously strong downwelling winds during this time period, however the longer perspective of site DI showed that there were other years similar in strength or stronger.

Similarly, MUR sea surface temperature data over the shelf indicated the 2020 study period prior to Sally was generally above average but not extreme (not shown). In general, the climatological MUR sea surface temperature conditions from mid-August to mid-September showed the fall transition to cooler temperatures, with the transition beginning at shallow coastal waters first (Fig. 5). Thus, the typical wind conditions during this time period coupled with the climatological sea surface temperature gradient would be expected to regularly feature downwelling events that deliver warmer offshore surface water to the coast and advect any reserve of colder bottom water offshore. This coupling of seasonal characteristics provides a coastal setting that is climatologically set-up to produce favorable conditions for tropical storm intensification as they transit the shelf. Thus, understanding the potential coupling of the drivers associated with the event sequence identified in this study is an important step for improving probabilistic representations of potential risk associated with hazards from tropical storms.

To expand the analysis beyond the Alabama coast, wind data from the NARR reanalysis was used to solve the $U_s/$ term from Equation 4 throughout the Gulf of Mexico and south Atlantic coast (Fig. 6). This term can be used to highlight regions that experience periods of significant downwelling during the peak of the Atlantic hurricane season and can potentially lead to increased heat content over the shelf for storm intensification. The average winds from August 1 to October 31 across 41 years (1979-2020) of data were used, and slopes were calculated from the shoreline to the 100m isobath. The distribution of this parameter showed the northern Gulf of Mexico was largely downwelling favorable during this time period with the Louisiana coast (i.e., coastline of the central northern Gulf of Mexico) being a hot spot for potential intensification due to the wide shelf and average westward winds. Such potential for favorable shelf thermal conditions in the northern Gulf of Mexico during the peak of hurricane season may contribute to the observed and simulated tendency of

tropical storms to rapidly intensify before landfall in this region (Landseas and Franklin 2013 and Emanuel 2017).

While the prospective of this study focuses on the potential importance of the resulting elevated heat content in the coastal ocean for the intensification of Hurricane Sally, the depth-average temperature in and of itself can be viewed as a coastal hazard. There is a rapidly growing body of work that highlights the impacts of marine heatwaves on ecosystems (e.g., Garrabou et al. 2009; Wernberg et al. 2013; Frölicher and Laufkötter 2018). As such, the event sequence that elevates the depth-average temperature and the potential climatological coupling of its drivers should be considered in the management of thermally sensitive ecosystems (e.g., coral reefs).

5. Conclusion

While continental shelves are relatively small regions of the global ocean, these regions are a critical transition zone that tropical cyclones must cross prior to landfall. Despite short periods of time spent over these regions (relative to the life span of a tropical cyclone), important modifications of storm characteristics can occur from the air-sea interactions. Importantly, understanding the initial coastal ocean conditions may become increasingly critical as some recent work indicates a slowing of mid-latitude tropical cyclones under warming climate scenarios (Yamaguchi et al. 2021). While much uncertainty surrounds such future behavior, a slowing of storms would provide more time over continental shelf environments allowing for more air-sea interactions prior to landfall. Thus, the potential slower speed of storms as well as the observed slower decay rate over land (Li and Chakraborty 2020), both emphasize the importance of understanding the processes that set up the initial thermal conditions in the coastal ocean, a primary source of storm energy before landfall.

In the case of Hurricane Sally, observational data from regional monitoring systems provided context for the state of the initial thermal conditions as well as the event sequence that drove the evolution of the observed hydrographic conditions. This study found that the depth-average temperature prior to Hurricane Sally reached 29.4°C, the 2020 annual peak and was approximately 0.7°C above the 15-year mean for the study time period. This peak in heat content was driven by a sequence of events that included the passage of Hurricane Marco and a persistent weather pattern that generated strong downwelling conditions. Interestingly, this work suggests that an indirect impact of freshwater over the shelf (i.e., limiting the downward transfer of heat during Marco and subsurface storage) may have limited the thermal energy available to Sally. However, the subsequent advection of a downwelling front approximately 64 km offshore likely exposed more than 50% of shelf region to nearly homogeneous water column conditions with temperatures exceeding 28.5°C. Thus, the event sequence shaping the thermal conditions over the shelf effectively produced a runaway for intensification as Hurricane Sally crossed the shelf.

Given the identified event sequence and the climatological patterns of wind

stress and warm upper ocean temperature during the study time period, hurricanes passing through this region of the Gulf of Mexico are likely to encounter favorable shelf conditions for intensification. The identification of other regions with such characteristics during hurricane season may provide additional context for anticipating the intensity response prior to landfall.

Acknowledgements

This work would not have been possible without the many folks that have passed through the Tech Support Group at the Dauphin Island Sea Lab to help maintain site CP over the years. We thank Rich Pawlowicz at the University of British Columbia for the freely available MATLAB `m_map` toolbox and Chris Landsea of NOAA NHC for providing links to early versions of the NOAA NHC best track data for 2020 storms. This research was made possible by the NOAA RESTORE Science Program (NA17NOS4510101 and NA19NOS4510194) and the NASA Physical Oceanography program (80NSSC21K0553). All data that were not downloaded from public sources have been made publicly available through or have been submitted to the NOAA National Center for Environmental Information (NCEI). In particular, site CP data are available at the following links: <https://accession.nodc.noaa.gov/0211052> ; and <https://data.nodc.noaa.gov/cgi-bin/iso?id=gov.noaa.nodc:0203749> . To be determined as the data has been submitted. A portion of this work was conducted at the Jet Propulsion Laboratory, California Institute of Technology, under contract with NASA. We thank the NASA Goddard Space Flight Center, Ocean Ecology Laboratory, Ocean Biology Processing Group for the Moderate-resolution Imaging Spectroradiometer (MODIS) Terra ocean color data; 2014 Reprocessing. NASA OB.DAAC, Greenbelt, MD, USA. doi:10.5067/AQUA/MODIS/MODIS_OC.2014.0.

References

- Alduchov, O.A. and Eskridge, R.E., 1996. Improved Magnus form approximation of saturation vapor pressure. *Journal of Applied Meteorology and Climatology*, 35(4), 601-609.
- August, E.F., 1828. Ueber die Berechnung der Expansivkraft des Wasserdunstes. *Ann. Phys. (Berlin)*, 89(5), 122-137.
- Austin, J. and Lentz, S., 2002. The inner shelf response to wind-driven upwelling and downwelling. *Journal of Physical Oceanography*, 32, 2171-2193.
- Bacmeister, J.T., Reed, K.A., Hannay, C., Lawrence, P., Bates, S., Truesdale, J.E., et al., 2018. Projected changes in tropical cyclone activity under future warming scenarios using a high-resolution climate model. *Climatic Change*, 146, 547-560.
- Balaguru, K., Foltz, G.R., Leung, L.R., D'Asaro, E., Emanuel, K.A., Liu, H., and Zedler, S.E., 2015. Dynamic Potential Intensity: An improved representation of the ocean's impact on tropical cyclones. *Geophysical Research Letters*, 42(16), 6739-6746.

Balaguru, K., Foltz, G.R., Leung, L.R., Hagos, S.M., and Judi, D.R., 2018. On the use of ocean dynamic temperature for hurricane intensity forecasting. *Weather and Forecasting*, 33(2), 411-418.

Balaguru, K., Foltz, G.R., Leung, L.R., Kaplan, J., Xu, W., Reul, N. and Chapron, B., 2020. Pronounced impact of salinity on rapidly intensifying tropical cyclones. *Bulletin of the American Meteorological Society*, 101(9), pp.E1497-E1511.

Berg, R. and Reinhart, B.J., 2021. *Hurricane Sally (AL192020): 11-17 September 2021*. National Hurricane Center Tropical Cyclone Report, NOAA/National Weather Service/National Hurricane Center.

Canuto, V.M., Howard, A., Cheng, Y., and Dubovikov, M.S., 2001. Ocean turbulence. Part I: One-point closure model-momentum and heat vertical diffusivities. *Journal of Physical Oceanography*, 31, 1413-1426.

Chu, J.E., Lee, S.S., Timmermann, A., Wengel, C., Stuecker, M.F. and Yamaguchi, R., 2020. Reduced tropical cyclone densities and ocean effects due to anthropogenic greenhouse warming. *Science advances*, 6(51), p.eabd5109.

Coogan, J., Dzwonkowski, B., and Lehrter, J., 2019. Effects of coastal upwelling and downwelling on hydrographic variability and dissolved oxygen in Mobile Bay. *Journal of Geophysical Research: Oceans*, 124(2), 791-806.

DeMaria, M., Sampson, C.R., Knaff, J.A., and Musgrave, K.D., 2014. Is tropical cyclone intensity guidance improving? *Bulletin of the American Meteorological Society*, 95(3), 387-398.

Domingues, R., Le Hénaff, M., Halliwell, G., Zhang, J.A., Bringas, F., Chardon, P. et al., 2021. Ocean conditions and the intensification of three major Atlantic hurricanes in 2017. *Monthly Weather Review*, 149(5), 1265-1286.

Dzwonkowski, B. and Yan, X.-H., 2005. Development and application of a neural network based ocean colour algorithm in coastal waters. *International Journal of Remote Sensing*, 26(6), 1175-1200.

Dzwonkowski, B., Park, K., Lee, J., Webb, B.M., and Valle-Levinson, A., 2014. Spatial variability of flow over a river-influenced inner shelf in coastal Alabama during spring. *Continental Shelf Research*, 74, 25-34.

Dzwonkowski, B., Fournier, S., Park, K., Dykstra, S.L., and Reager, J.T., 2018. Water column stability and the role of velocity shear on a seasonally stratified shelf, Mississippi Bight, northern Gulf of Mexico. *Journal of Geophysical Research: Oceans*, 123(8), 5777-5796.

- Dzwonkowski, B., Coogan, J., Fournier, S., Lockridge, G., Park, K., and Lee, T., 2020. Compounding impact of severe weather events fuels marine heatwave in the coastal ocean. *Nature Communications*, *11*, 4623, doi:10.1038/s41467-020-18339-2.
- Emanuel, K., 1999. Thermodynamic control of hurricane intensity. *Nature*, *401*(6754), 665-669.
- Emanuel, K., 2005. Increasing destructiveness of tropical cyclones over the past 30 years. *Nature*, *436*, 686-688.
- Emanuel, K., 2017. Will global warming make hurricane forecasting more difficult? *Bulletin of the American Meteorological Society*, *98*(3), 495-501.
- Fairall, C., Bradley, E., Rogers, D., Edson, J., and Young, G., 1996. Bulk parameterization of air-sea fluxes for Tropical Global Atmosphere Coupled-Ocean Atmosphere Response Experiment. *Journal of Geophysical Research*, *101*, 3747-3764.
- Frölicher, T.L. and Laufkötter, C., 2018. Emerging risks from marine heat waves. *Nature Communications*, *9*, 650, doi:10.1038/s41467-018-03163-6.
- Garrabou, J., Coma, R., Bensoussan, N., Bally, M., Chevaldonne, P., Cigliano, M. et al., 2009. Mass mortality in Northwestern Mediterranean rocky benthic communities: effects of the 2003 heat wave. *Global Change Biology*, *15*, 1090-1103.
- Glenn, S.M., Miles, T.N., Seroka, G.N., Xu, Y., Forney, R.K., Yu, F., et al., 2016. Stratified coastal ocean interactions with tropical cyclones. *Nature Communications*, *7*, 10887, doi:10.1038/ncomms10887.
- Hobday, A.J., Alexander, L.V., Perkins, S.E., Smale, D., Straub, S., Olive, E.C.J. et al., 2016. A hierarchical approach to defining marine heatwaves. *Progress in Oceanography*, *141*, 227-238.
- Kaplan, J. and DeMaria, M., 2003. Large-scale characteristics of rapidly intensifying tropical cyclones in the North Atlantic basin. *Weather and Forecasting*, *18*(6), 1093-1108.
- Kaplan, J., DeMaria, M., and Knaff, J.A., 2010. A revised tropical cyclone rapid intensification index for the Atlantic and eastern North Pacific basins. *Weather and Forecasting*, *25*(1), 220-241.
- Keim, B.D., Muller, R.A., and Stone, G.W., 2007. Spatiotemporal patterns and return periods of tropical storm and hurricane strikes from Texas to Maine. *Journal of Climate*, *20*(14), 3498-3509.
- Knutson, T., Camargo, S.J., Chan, J.C.L., Emanuel, K., Ho, C.-H., Kossin, J. et al., 2020. Tropical cyclones and climate change assessment: Part II: Projected response to anthropogenic warming. *Bulletin of the American Meteorological Society*, *101*(3), E303-E322.

- Landsea, C. W., and J. L. Franklin, 2013: Atlantic hurricane database uncertainty and presentation of a new database format. *Monthly Weather Review*, 141, 3576–3592, doi:10.1175/MWR-D-12-00254.1.
- Li, L. and Chakraborty, P., 2020. Slower decay of landfalling hurricanes in a warming world. *Nature*, 587, 230-234.
- Liu, M., Vecchi, G.A., Smith, J.A., and Knutson, T.R., 2019. Causes of large projected increases in hurricane precipitation rates with global warming. *npj Climate and Atmospheric Science*, 2, 38, doi:10.1038/s41612-019-0095-3.
- Magnus, G., 1844. Versuche über die Spannkkräfte des Wasserdampfs. *Ann. Phys. (Berlin)*, 137(2), 225-247.
- Miles, T., Seroka, G., and Glenn, S., 2017. Coastal ocean circulation during Hurricane Sandy. *Journal of Geophysical Research: Oceans*, 122(9), 7095-7114.
- Park, J.-H., Yeo, D.-E., Lee, K., Lee, H., Lee, S.W., Noh, S., et al. Rapid decay of slowly moving Typhoon Soulik (2018) due to interactions with the strongly stratified northern East China Sea. *Geophysical Research Letters*, 46(24), 14595-14603.
- Potter, H., Drennan, W.M., and Graber, H.C., 2017. Upper ocean cooling and air-sea fluxes under typhoons: A case study. *Journal of Geophysical Research: Oceans*, 122(9), 7237-7252.
- Potter, H., DiMarco, S.F., and Knapp, A.H., 2019. Tropical cyclone heat potential and the rapid intensification of Hurricane Harvey in the Texas Bight. *Journal of Geophysical Research: Oceans*, 124(4), 2440-2451.
- Price, J.F., 2009. Metrics of hurricane-ocean interaction: vertically-integrated or vertically-averaged ocean temperature? *Ocean Science*, 5(3), 351-368.
- Pun, I.-F., Chan, J.C.L., Lin, I.-I., Chan, K.T.F., Price, J.F., Ko, D.S. et al., 2019. Rapid intensification of Typhoon Hato (2017) over shallow water. *Sustainability*, 11(13), 3709, doi:10.3390/su11133709.
- Reul, N., Chapron, B., Grodsky, S.A., Guimbard, S., Kudryavtsev, V., Foltz, G.R. and Balaguru, K., 2021. Satellite observations of the sea surface salinity response to tropical cyclones. *Geophysical Research Letters*, 48(1), p.e2020GL091478.
- Seroka, G., Miles, T., Xu, Y., Kohut, J., Schofield, O. and Glenn, S., 2016. Hurricane Irene sensitivity to stratified coastal ocean cooling. *Monthly Weather Review*, 144(9), 3507-3530.
- Tzeng, M.W., Dzwonkowski, B., and Park, K., 2016. Data processing for a small-scale long-term coastal ocean observing system near Mobile Bay, Alabama. *Earth and Space Science*, 3(12), 510-522.
- Vecchi, G.A., Delworth, T.L., Murakami, H., Underwood, S.D., Wittenberg, A.T., Zeng, F. et al., 2019. Tropical cyclone sensitivities to CO₂ doubling:

Roles of atmospheric resolution, synoptic variability and background climate changes. *Climate Dynamics*, 53(9-10), 5999-6033.

Walker, N.D., Leben, R.R., and Balasubramanian, S., 2005. Hurricane-forced upwelling and chlorophyll a enhancement within cold-core cyclones in the Gulf of Mexico. *Geophysical Research Letters*, 32, L18610, doi:10.1029/2005GL023716.

Wernberg, T., Smale, D.A., Tuya, F., Thomsen, M.S., Langlois, T.J., de Bettignies, T. et al., 2013. An extreme climatic event alters marine ecosystem structure in a global biodiversity hotspot. *Nature Climate Change*, 3, 78-82.

Yamaguchi, M. and Maeda, S., 2020. Slowdown of Typhoon translation speeds in mid-latitudes in September influenced by the Pacific Decadal Oscillation and global warming. *Journal of the Meteorological Society of Japan*, 98(6), 1321-1334.

Zhang, Z., Wang, Y., Zhang, W., and Xu, J., 2019. Coastal ocean response and its feedback to Typhoon Hato (2017) over the South China Sea: A numerical study. *Journal of Geophysical Research: Atmospheres*, 124(24), 13731-13749.




Figure 1. Map of the Mississippi Bight with high resolution (1 km) Terra MODIS

ocean color data from August 18 2020, highlighting the large plume of the Mississippi River water impacting the study region prior to the passage of Hurricane Marco on August 24. Locations of monitoring stations used in the Alabama coastal region are shown: site CP (CP,), DPIA1 (DI,) and the NOAA Orange Beach Buoy (ORB,) as well as the 20, 100, and 200 m isobaths (three solid lines). In addition, the estimated foot of a downwelling front using 28.5°C isotherm on September 14 is shown (dotted line). The inset shows the study region (black box) in the broader context of the Gulf of Mexico region with storm tracks for Hurricane Marco (blue) and Hurricane Sally (black).




Figure 2. Evolution of the environmental data encompassing the period prior to, during and post landfall of Hurricane Sally including (a) depth-average temperature at site CP (red), a proxy for shelf heat content, (b) along and across-shelf wind speed components at ORB, (c) along-shelf depth-average currents at site CP, (d) near-surface and near-bottom across-shelf currents at site CP, (e) vertical temperature structure at site CP, (f) salinity at site CP, and (g) estimated surface heat flux at ORB. The inverse triangles at the top of each panel indicates the closest times to landfall for Hurricane Marco (8/24/2020) and landfall for Hurricane Sally (09/16/2020). Also shown in (a) are the long-term mean (black line) and the 90th percentile threshold (dotted line) based on the 15-year (2005-2020) data, respectively. In (e), the black contours indicate isotherms (26°C, 28.5°C and 30°C) and the horizontal black line at the bottom indicates the downwelling period (September 9-14) over which a scaled advection of an observed front (using the 28.5°C isotherm) was applied to estimated offshore distance.




Figure 3. Time series of (a) the bulk buoyancy frequency (bulk N^2) and bulk horizontal vertical shear squared (bulk S^2) and (b) the bulk Richardson number (bulk Ri) at site CP prior to, during and after the passage of Hurricane Marco. Also shown are the critical Ri ($Ri_{cr} = 0.25$), the horizontal red dashed line in (b), and the date of Marco passing just to the south of the Mississippi River Delta ().




Figure 4. Mean along-shelf wind stress during the period between August 15 and September 15 from long-term records at ORB (red) and DI (black), 9 and 30 years, respectively. The negative wind stress indicates westward wind that favor downwelling. Gaps in the time series occur in years where data was not available for the time period.




Figure 5. Hovmoller diagram of the climatological MUR sea surface temperature pattern during August-September, derived from the 2004-2020 data, across the Mississippi Bight shelf along the longitude of site CP. The dashed and dotted lines indicate the locations of site CP and the shelf break.




Figure 6 - Relative comparison of the $\frac{U_s}{\alpha}$ term in the scaling argument (Eq. 4) based on average winds from August 1 to October 31 (1979-2020) using wind data from the NARR model where red denotes downwelling winds and blue denotes upwelling winds. The darker the shading the more predominant the advective response is expected to be in that region.