

**Table1: Baseline Clinical and Procedural Characteristics**

	Total (n=209)	Anticoagulation			Bleeding History		
		Warfarin (n=152)	DOAC (n=57)	P- value*	Positive Bleeding History (n=147)	No Bleeding History (n=62)	p- value *
Age	74.6 ± 7.8	74.3±8.0	75.3±7.0	0.392	74.6± 7.4	74.6± 8.6	0.97
Sex (% male)	57%	58%	56%	0.88	59%	55%	0.65
Body mass index (BMI)	29.7 ± 7.0	29.7±7.0	29.5±7.0	0.846	29.7 ± 6.6	29.7 ± 7.8	0.98
Obstructive sleep apnea (OSA)	33(15.8%)	24(15.8%)	9(15.8%)	1.0	25 (17.0%)	8 (12.9%)	0.54
Hypertension	195(93.3%)	141(92.7%)	54(94.7%)	0.763	137 (93.2%)	58 (93.5%)	1.0
Diabetes mellitus	63(30.1%)	45(29.6%)	18(31.6%)	0.866	43(29.3%)	20 (32.3%)	0.74
History of Stroke	42(20.1%)	34(22.4%)	8(14.0%)	0.245	29 (19.7%)	13 (21.0%)	0.85
Myocardial infarction	3(1.4%)	3(2.0%)	0	0.564	2 (1.4%)	1 (1.6%)	1.0
Peripheral arterial disease	7(3.3%)	7(4.6%)	0	0.193	6 (4.1%)	1 (1.6%)	0.68
Coronary artery disease	54(25.8%)	44(28.9%)	10(17.5%)	0.111	44 (29.9%)	10 (16.1%)	0.039
Congestive heart failure	48(23.0%)	38(25.0%)	10(17.5%)	0.275	37 (25.2%)	11 (17.7%)	0.28
Ejection fraction	53.0 ± 8.9	52.3±9.2	55 ±7.8	0.052	52.4 ± 9.3	54.4 ± 7.8	0.14
Chronic kidney disease	16(7.7%)	16(10.5%)	0	0.009	14 (9.5%)	2 (3.2%)	0.16
Paroxysmal (versus persistent) atrial fibrillation	97(46.4%)	80(52.6%)	17(29.8%)	0.005	69 (46.9%)	28 (45.2%)	0.88
HAS-BLED score†	4[3-4]	4 [3-4]	4 [3-4]	0.329	4[4-4]	3[3-3]	<0.00 1
CHA <sub>2</sub> DS <sub>2</sub> -VASc score†	4 [3-5]	4 [3-5]	4 [3-4]	0.170	4 [3-5]	4[3-4]	0.243
Device size (mm)	25.2 ± 3.4	25.2±3.3	25.4±3.5	0.651	25.6 ± 3.4	24.5 ± 3.1	0.032
Follow up (months)	23.5 ± 7.1	25.3±7.1	18.6± 4.3	<0.0001	24.2 ± 7.1	21.7 ± 6.8	0.016

\* P value computed from Student's t test for continuous variables, Fisher exact test for categorical values. Equality of medians test used for data presented as median and interquartile range (HAS-BLED and CHA<sub>2</sub>DS<sub>2</sub>-VASc scores).

† data presented as median and interquartile range