

Table2. A multivariate logistic regression model for initial IVIG resistance in patients with KD.

Variates	β	SE	Walds	P value	OR	95%CI
<i>Initial IVIG resistance-PT</i>						
Platelet	0.005	0.002	7.523	0.006*	1.01	1.00-1.01
Creatinine	-0.034	0.020	2.830	0.093	0.97	0.93-1.01
Urea nitrogen	-0.168	0.125	1.804	0.179	0.85	0.66-1.08
ALT	-0.002	0.001	2.365	0.124	0.99	0.99-1.00
PT	-0.091	0.041	5.060	0.024*	0.91	0.84-0.99
<i>Initial IVIG resistance-APTT</i>						
Platelet	0.004	0.002	1.323	0.038*	1.00	1.00-1.01
Creatinine	-0.019	0.022	0.707	0.400	0.98	0.94-1.03
Urea nitrogen	-0.190	0.133	2.017	0.152	0.83	0.64-1.07
ALT	-0.002	0.001	2.110	0.146	0.99	0.99-1.00
Potassium	0.933	0.291	10.29	0.001*	2.54	1.44-4.50

APTT	-0.051	0.019	7.033	0.008*	0.95	0.92-0.99
------	--------	-------	-------	--------	------	-----------

Initial IVIG resistance-D-dimer

Platelet	0.004	0.002	4.257	0.039*	1.00	1.00-1.01
Creatinine	-0.016	0.023	0.532	0.466	0.98	0.94-1.03
Urea nitrogen	-0.175	0.133	1.746	0.186	0.84	0.65-1.09
ALT	0.000	0.001	0.350	0.554	1.00	0.99-1.00
Potassium	0.953	0.300	10.06	0.002*	2.59	1.44-4.67
Total bilirubin	-0.021	0.012	3.419	0.064	0.98	0.96-1.00
D-dimer	-0.129	0.063	4.237	0.040*	0.88	0.78-0.99

Initial IVIG resistance-ATIII

Creatinine	-0.046	0.021	5.024	0.025*	0.96	0.92-0.99
Urea nitrogen	-0.144	0.127	1.293	0.256	0.87	0.68-1.11
ALT	-0.002	0.001	2.285	0.132	1.00	0.99-1.00
ATIII	0.030	0.008	14.17	<0.001*	1.03	1.01-1.05

Abbreviations: IVIG, intravenous immunoglobulin; PT, prothrombin time; APTT, activated partial thromboplastin time; AT, antithrombin; ALT, alanine aminotransferase

*Statistically significant ($P<0.05$)