

**Table 1. Characteristics of pediatric lymphoma patients who completed Pediatric PROMIS-25 surveys at diagnosis in Lilongwe, Malawi, stratified by outcome.**

	All patients N = 75	Alive N = 30	Died/LTFU N = 45	p-value
<b><i>Patient Characteristics, n (%)</i></b>				
Age, median [IQR]	10 [8-12]	9 [7-12]	10 [9-12]	0.197
Male sex	51 (68)	20 (67)	31 (69)	1.00
HIV-negative	66 (90)	25 (89)	41 (91)	0.666
Diagnosis				0.018
Burkitt lymphoma	61 (81)	20 (67)	41 (91)	
Hodgkin lymphoma	14 (19)	10 (33)	4 (9)	
Stage III/IV	48 (66)	16 (53)	32 (74)	0.106
Lanksy Performance Score $\leq 70$	50 (70)	15 (52)	35 (82)	0.009
Moderate-to-Severe Acute Malnutrition	54 (72)	19 (63)	35 (78)	0.111
<b><i>Laboratory Results, median [IQR]</i></b>				
Hemoglobin, g/dL	10.0 [8.5-11.4]	10.0 [9.4-11.0]	10.0 [8.2-11.5]	0.751
Lactate dehydrogenase, IU/L	555 [378-895]	460 [354-662]	739 [460-1333]	0.006
Lactate dehydrogenase > 2x ULN, n (%) *	30 (40)	10 (33)	20 (44)	0.350
White blood cell count, $10^3/\mu\text{L}$	8.3 [5.6-13.1]	8.9 [5.8-14.1]	8.1 [5.4-12.2]	0.441
Platelet count, $10^3/\mu\text{L}$	330 [237-437]	357 [292-421]	280 [168-486]	0.158
Albumin, g/dL	3.5 [3.1-3.8]	3.6 [3.4-4.0]	3.2 [3.0-3.8]	0.101
<b><i>PROMIS-25 Domain Scores, mean [SD]</i></b>				
Mobility	31.5 (9.1)	34.1 (9.5)	29.8 (8.6)	0.049
Anxiety	61.9 (9.8)	61.0 (10.1)	62.4 (9.6)	0.528
Depressive Symptoms	62.2 (6.7)	61.4 (6.9)	62.8 (6.6)	0.402
Fatigue	61.7 (9.2)	60.8 (9.2)	62.3 (9.3)	0.513
Peer Relationships	49.3 (7.3)	48.4 (7.9)	49.9 (7.0)	0.376
Pain Interference	59.6 (9.5)	58.4 (10.0)	60.5 (9.2)	0.353
Pain Intensity, median [IQR]	9 [4-10]	8.5 [3.3-10.0]	10.0 [6.0-10.0]	0.073
Higher PROMIS scores for Mobility and Peer Relationships reflect better functioning and relationships; lower symptom scores reflect better symptom experiences.				
*LDH Upper Limit of Normal (ULN) for age: Age 2-12 = 295 U/L, Age 13-17 = 190 U/L. LTFU = Lost to Follow-up; HIV= Human Immunodeficiency Virus; ULN= Upper limit of normal; IQR = Interquartile range; SD = Standard Deviation				

