

Table 3. Complications

Complication type (30 days)	CPER n/162 (%)	NCEPR n/179 (%)	Clavien- Dindo Grade (post-op)
No complication	150/162 (92.6%)	164 (91.6%)	n/a
Ruptured ectopic following initial expectant/medical management	3/162 (1.9%)	4/179 (2.2%)	n/a
Conversion from laparoscopy to laparotomy	1 (0.6%)	3 (1.7%)	n/a
Significant additional intra-op procedure*	0	2 (1.1%)	n/a
Anaphylaxis to muscle relaxant requiring ITU admission ⁺	1 (0.6%)	0	n/a
Methotrexate toxicity	1 (0.6%)	0	n/a
Transfusion pre-op	0	1 (0.6%)	n/a
Transfusion pre- methotrexate	1 (0.6%)	0	n/a
Concurrent LRTI	0	1 (0.6%)	1
Transfusion post-op	0	2 (1.1%)	2
Post-op pyrexia	1 (0.6%)	0	1
Re-presentation post-op pain/ bleeding	3 (1.9%)	1 (0.6%)	1
Re-presentation post-op infection (PID) ⁺	1 (0.6%)	0	1
Re-presentation post-op pain + rising bhcg ^{**}	1 (0.6%)	0	2
Repeat laparoscopy ^{***}	0	1 (0.6%)	3b

* One patient required- adheysiolysis, salpingectomy + bilateral ovarian cystectomy and one patient required salpingectomy + small bowel resection

** required further management with methotrexate

*** first laparoscopy for evacuation of blood and second laparoscopy for oophorectomy after ongoing rising HCG

⁺ same patient