

Table 2: Summary of ultrasound diagnoses detected in the study population (N = 1341)

Diagnosis	N (%)
No abnormalities	884 (65.9)
Fibroids	196 (14.6)
Para-ovarian cyst	103 (7.7)
Endometriosis	66 (4.9)
Ovarian cyst	51 (3.8)
Simple cyst	21 (1.6)
Dermoid cyst	19 (1.4)
Haemorrhagic cyst	8 (0.6)
Cystadenoma	2 (0.1)
Fibroma	1 (0.1)
Adenomyosis	41 (3.1)
Polycystic ovaries	36 (2.7)
Congenital uterine anomalies	30 (2.2)
Arcuate	12 (0.9)
Subseptate	12 (0.9)
Bicornuate	3 (0.2)
Unicornuate	3 (0.2)
Pelvic vein thrombus	11 (0.8)
Pelvic adhesions	8 (0.6)
Endometrial polyp	6 (0.4)
IUCD in situ	5 (0.4)
Ovarian hyperstimulation	5 (0.4)
Hydrosalpinx	3 (0.2)
Intrauterine adhesions	3 (0.2)
Ovarian torsion	2 (0.1)
Other *	5 (0.4)

*includes urinary retention, vaginal wall cyst, metastatic bowel cancer, cervical mucous retention cyst

