

Table 2. Significant hits in urine and plasma obtained from Volcano plots, including mean fold-changes of normalized intensity in the inhibition session versus control session, and mean fold-change of normalized intensity in the PM subjects versus EM-UM subjects.

<i>m/z</i>	RT (min)	Ionization Mode	<i>Inhibition session vs Control</i>		<i>PM subjects (n = 6) vs session (n = 37)</i>		<i>EM-UM subjects (n = 37)</i>	
			Fold- change	P-value (adjusted)	Fold- change	P-value		
Plasma								
416.3159	4.16	+	0.60	4.48×10^{-4}	5.93×10^{-3}	3.28×10^{-7}		
595.3236	5.25	-	0.16	4.10×10^{-5}	4.41×10^{-2}	2.10×10^{-5}		
597.3382	5.25	+	0.13	9.38×10^{-8}	6.52×10^{-2}	2.95×10^{-5}		
Urine								
220.1543	2.51	+	0.54	2.63×10^{-7}	0.34	1.44×10^{-5}		
416.3159	4.16	+	0.67	4.53×10^{-3}	1.82×10^{-2}	3.94×10^{-6}		
432.3108	4.11	+	0.56	1.39×10^{-2}	8.56×10^{-3}	3.28×10^{-7}		
444.3108	4.35	+	0.67	3.50×10^{-5}	2.06×10^{-2}	3.94×10^{-6}		