

1 **Table 3. Adjusted associations Between Participant Characteristics and Timing of**  
2 **Introduction of specific allergen:**  
3

	<b>Peanut</b>	<b>Milk</b>	<b>Egg</b>
<b>Participant Characteristics Associated with: Early Introduction of specific allergen (6 months or before)</b>	<b>Odds Ratio (95% CI)</b>	<b>Odds Ratio (95% CI)</b>	<b>Odds Ratio (95% CI)</b>
<b>White (vs Black)</b>	3.6 (1.5-9.8)	2.9 (1.2-7.2)	3.1 (1.3-9.0)
<b>Birth year 2008-2016 (vs pre-2008)</b>	1.4 (0.4-9.6)	0.8 (0.2-2.8)	2.1 (0.5-14.0)
<b>Birth year 2017-2019 (vs pre-2008)</b>	6.3 (1.5-44.4)	2.2 (0.5-9.4)	3.5 (0.8-24.7)
<b>Multiple (vs single) Food Allergies</b>	1.6 (0.7-4.3)	0.8 (0.2-2.5)	0.5 (0.2-1.2)
<b>Female (vs Male)</b>	1.9 (0.9-4.1)	1.1 (0.5-2.6)	1.0 (0.5-2.0)
<b>Eczema (vs no eczema)</b>	1.3 (0.5-3.9)	4.6 (1.4-17.3)	0.7 (0.3-1.7)
<b>Rush University Site (vs Lurie)</b>	0.8 (0.2-2.4)	1.9 (0.6-5.7)	0.9 (0.3-2.8)
<b>Children's National (DC) Site (vs Lurie)</b>	2.4 (1.0-6.0)	0.8 (0.2-2.5)	1.1 (0.4-2.9)
<b>Cincinnati Children's Site (vs Lurie)</b>	0.3 (0.1-0.1)	0.6 (0.2-1.6)	0.4 (0.1-1.0)
<b>Patient Characteristics Associated with: Later introduction of specific allergen (12 or more months of age)</b>			
<b>White (vs Black)</b>	0.2 (0.1-0.4)	0.2 (0.1-0.4)	0.4 (0.2-0.8)
<b>Birth year 2008-2016 (vs pre-2008)</b>	0.7 (0.2-1.8)	2.6 (0.7-11.3)	0.5 (0.2-1.4)

<b>Birth year 2017-2019 (vs pre-2008)</b>	0.1 (0.02-0.2)	0.2 (0.02-1.4)	0.1 (0.03-0.3)
<b>Multiple (vs single) Food Allergies</b>	0.8 (0.4-1.6)	2.5 (0.6-11.7)	1.3 (0.6-2.6)
<b>Female (vs Male)</b>	0.6 (0.3-1.1)	1.2 (0.4-3.0)	0.9 (0.5-1.6)
<b>Eczema (vs no eczema)</b>	0.5 (0.2-1.2)	0.3 (0.1-1.0)	1.5 (0.7-3.3)
<b>Rush University Site (vs Lurie)</b>	0.6 (0.2-1.5)	0.9 (0.3-2.8)	1.3 (0.5-3.5)
<b>Children's National (DC) Site (vs Lurie)</b>	0.5 (0.2-1.0)	1.3 (0.3-4.9)	1.3 (0.5-3.3)
<b>Cincinnati Children's Site (vs Lurie)</b>	0.7 (0.3-1.6)	0.9 (0.2-3.0)	1.3 (0.6-2.9)

\*CI: Confidence Interval

4

5