

Table 2: Characteristics of patients who received anticoagulation

Patient no.	Age (in years)	Gender	Diagnosis	Location of Tumor Thrombus	Anticoagulation	Reason for anticoagulation
1	2.5	Female	Wilms tumor	IVC, Renal vein and hepatic vein	Lovenox	To avoid progression of IVC thrombus
2	3	Male	Neuroblastoma	IVC and right atrium	Unfractionated heparin and Lovenox	IVC and intracardiac thrombus
3	4	Female	Wilms tumor	Renal vein, hepatic vein, IVC and right atrium	Lovenox	IVC and intracardiac thrombus
4	10	Male	Embryonal sarcoma	IVC, portal and hepatic veins	Lovenox	To avoid further progression of IVC Thrombus. Also developed line-associated thrombus.
5	12	Male	Rhabdomyosarcoma	IVC, Right atrium	Unfractionated heparin and Lovenox	Intracardiac tumor thrombus
6	14	Female	Osteosarcoma	Right atrium and pulmonary arteries	Unfractionated heparin and Lovenox	Pulmonary embolism, Intracardiac thrombus
7	15	Female	Ewing sarcoma	IVC and Pulmonary arteries	Unfractionated heparin and Lovenox	Pulmonary embolism
8	15	Male	Clear cell sarcoma	Renal vein and right atrium	Lovenox	Symptomatic thrombus with lower extremity swelling and pain
9	16	Female	Osteosarcoma	Pulmonary arteries	Lovenox	To prevent further progression
10	18	Male	Embryonal carcinoma	IVC and renal vein	Unfractionated heparin and Lovenox	Renal vein thrombosis (RVT)