

Table 2. Diagnostic imaging use by age group in low-value injury admissions

Age group	All	1-15	16-64	65-74	75+
N (%)	19163 (100)	1937 (10.1)	4714 (24.6)	2414 (12.6)	10098 (52.7)
No imaging	898 (4.7)	335 (17.3)	235 (5)	78 (3.2)	250 (2.5)
X-ray	15680 (81.8)	1087 (56.1)	3656 (77.6)	2033 (84.2)	8904 (88.2)
CT (any)	11952 (62.4)	740 (38.2)	3097 (65.7)	1562 (64.7)	6553 (64.9)
CT (head)	7438 (38.8)	555 (28.7)	1300 (27.6)	935 (38.7)	4648 (46)
CT (thoraco-abdominal)	2612 (13.6)	89 (4.6)	1004 (21.3)	365 (15.1)	1154 (11.4)
CT (spine)	3907 (20.4)	169 (8.7)	1356 (28.8)	491 (20.3)	1891 (18.7)
CT (pan scan)	323 (1.7)	7 (0.4)	224 (4.8)	28 (1.2)	64 (0.6)
CT (other)	3670 (19.2)	101 (5.2)	938 (19.9)	548 (22.7)	2083 (20.6)
MRI	555 (2.9)	39 (2)	252 (5.3)	84 (3.5)	180 (1.8)
AA	567 (3)	17 (0.9)	266 (5.6)	86 (3.6)	198 (2)

CT: Computed tomography, MRI: Magnetic resonance imaging, AA: Angiography / arteriography