

Conservation Laws and Explicit Solution of system of Fractional-Order Coupled Nonlinear Hirota Equations by Lie Symmetry Analysis

H. Gandhi^{1,a)}, A. Tomar^{2,b)} and D. Singh^{3,c)}

^{1,2}*Amity School of Applied Science, Amity University, Haryana, India*

³*Amity Institute of Applied Science, Amity University, Noida, U.P, India.*

^{a)} hemantgandhi1977@gmail.com,

^{b)} Corresponding Author: amitmath.14@gmail.com, ^{c)} dsingh@ggn.amity.edu

Abstract: The main objective of this research article is to summarize the study of the application of Lie symmetry reduction to the fractional order coupled nonlinear complex Hirota system of partial differential equations. By the efficient use of symmetries and explicit solutions, this system reducing to nonlinear fractional ordinary differential equations (FODEs) with the application of Erdyli-Kober (E-K) operators for fractional derivatives and integrals depending on real order θ . Investigating the convergent series solution along with the adjoint system and providing the conservation laws by Noether's theorem.

Keywords: Lie symmetry reduction, Nonlinear system of Hirota equations, Erdyli-Kober operators, Noether's theorem.

2010 MSC: 34A08, 26A33, 76M60

1. Introduction

Fractional calculus is the theory of fractional integrals and derivatives of arbitrary order which is evolved towards the end of 17th century. Evidently, fractional calculus is old as classical calculus; many physicists and mathematicians [1, 3-5] like Lagranges, Euler, Abel, Gronwald, Letnikov, Riemann and Caputo have done great contribution on fractional-order derivatives. In the current research, there have been fast development in the field of fractional order systems involving in science, engineering and technology i.e. bioscience, fluid flow, electromagnetic, viscoelasticity, pandemic analysis, medicine, infectious modeling, drug therapy, image processing and diffusion wave equations [9, 10, 15, 16, 32].

Every researcher or mathematician has open challenge to overcome the difficulties and to find the exact solution of fractional-order systems. The person, who knows the utility of

fractional derivatives and integrals, is only authorized to work on such kind of fractional partial differential equations (FPDEs), because each definition has its own limitations and benefits. Frenandez et al. [42] represented the series analysis of fractional operators involving Mittag-Leffler functions. Guariglia et al. [43] explained fractional derivatives of Riemann-Zeta function. Srivastava et al. [44] described the utility of fractional-order system in the biological population model with carrying capacity. The similarity analysis for strong shocks in non-ideal gas and numerical simulation of Ito coupled system have been analyzed by Arora et al. [11, 35]. Authors [6-8, 17, 18, 28, 30] suggested semi analytic and computational methodologies for obtaining the series and exact solutions of FODEs and FPDEs involving Laplace transform method, Sine-Cosine technique, homotopy analysis, variation iteration method, homotopy perturbation method, reduced differential transform and power series technique but these methods are useful to provide numerical or approximate series solutions. In our recent work, we are interested in application of Lie symmetry analysis [2], which has capability to provide us explicit and exact solution of fractional and integer-order problems. Authors [19-23, 25, 26] have imposed the invariance analysis and E-K operators on system of FPDEs with R-Lfractional derivative approach. Nowadays, researchers are working on conservation laws analysis of the systems. Sneddon [24] explained the application of fractional E-K differ-integral operators in beautiful manner, which is capable to convert the system of time fractional FPDEs into FODEs. Noether's theorem [27, 31, 33] established a relation between conservation laws and symmetry of differential equations and applied on FPDEs without Lagrangian operators. Recently, authors [34, 36-39, 47] provided invariance structure, explicit exact solutions with power series solution and conservation analysis of Boussinesq-Burger's system, Drinfeld-Sokolov-Satsuma-Hirota coupled KdV and m-KdV equations via Lie symmetry analysis. Biswas et al. [12-14] have worked on dual dispersion, power laws, conservation laws and optimal quasi-solitons by Lie symmetry analysis. Chauhan and Arora [29] has obtained the complete analysis of time fractional Kupershmidt equation. Recently, Gandhi et al. [40, 41, 50] have applied symmetry reduction on multi-ordered time-fractional KdV equations and Hirota-Satsoma-coupled Korteweg-de-Vries equations to obtain the explicit solutions with convergence and conservation laws; he concluded that the fractional-order parameter θ can control the output of solution of fractional mathematical models and in physical and mathematical aspects, the conservation laws play very crucial role to discuss the consistency of system. Zhang et al. [45] promoted the

conservation laws of Fokkar-Plank equation with power diffusion. Bruzon et al. [46] focused on the study of similarity solutions and application of new conservation theorem on Cooper-Shepard-Sodano equation. New soliton solution of time-fractional Drinfeld-Sokolov-Satsuma-Hirota system in dispersive water waves has been illustrated by Ray et al. [48], he claimed the analytical solution with Adomian polynomials and Tanh-method. A generalized two-component Hunter-Saxton system has been studied by Yang et al. [49]. Dubey et al. [51] has suggested the copulation of fractional homotopy perturbation and fractional homotopy analysis method with Sumudu transforms on local-fractional Laplace equation. The Hydon method has been used by Chatibi et al. [52] to construct discrete symmetries for a family of ordinary, partial, and fractional differential equations. Using the Lie point symmetry, the generalized invariant structure of the (2+1)-dimensional Date-Jimbo-Kashiwara-Miwa equation has been proposed by Chauhan et al. [53].

In present article, a nonlinear time fractional complex Hirota system of partial differential equations in Riemann-Liouville (RL) definition is considered by Lie symmetry approach. The main goal of this work is to elaborate the utilization of symmetry analysis on nonlinear fractional system of four equations with their conservation analysis. Hirota equations have many applications in propagation of sound and optical pulses in water crystals waveguide along with single mode fibers study. The complex Hirota system [21] is explained as follows:

$$\begin{cases} \frac{\partial^\theta z_1}{\partial t^\theta} + \frac{\partial^3 z_1}{\partial x^3} + 6(|z_1|^2 + |z_2|^2) \frac{\partial z_1}{\partial x} = 0, \\ \frac{\partial^\theta z_2}{\partial t^\theta} + \frac{\partial^3 z_2}{\partial x^3} + 6(|z_1|^2 + |z_2|^2) \frac{\partial z_2}{\partial x} = 0, \end{cases} \quad (1)$$

where $z_1(x, t) = u(x, t) + i v(x, t)$ and $z_2(x, t) = w(x, t) + i z(x, t)$,

and $(|z_1|^2 + |z_2|^2) = u^2 + v^2 + w^2 + z^2 = \sum u^2$.

The system reduced to:

$$\begin{cases} \partial_t^\theta u + u_{xxx} + 6u_x \left(\sum u^2 \right) = 0, \\ \partial_t^\theta v + v_{xxx} + 6v_x \left(\sum u^2 \right) = 0, \\ \partial_t^\theta w + w_{xxx} + 6w_x \left(\sum u^2 \right) = 0, \\ \partial_t^\theta z + z_{xxx} + 6z_x \left(\sum u^2 \right) = 0, \end{cases} \quad (2)$$

here $\partial_t^\theta u, \partial_t^\theta v, \partial_t^\theta w$ and $\partial_t^\theta z$ are partial derivatives of u, v, w and z with fractional parameter θ ($0 < \theta < 1$) and independent variables 'x' and 't' respectively. In continuation, section 2 is dedicated to preliminaries and invariance study of the system with the systematic use of Lie symmetry approach to find explicit solution in section 3. Section 4 is devoted to application of Erdlyi-Kober operators for conversion of system of FPDEs into FODEs. In section 5, power series solution processed with convergence analysis under implicit function theorem. Finally, adjoint system and conservation laws are studied via Noether's theorem is proposed in section 6 and section 7. The conclusion and remarks are presented with references at the end of work.

2. Preliminaries

In this part, author would like to design some fundamentals and definitions for the sake of understanding the methodologies and key points, concerned with local fractional real order derivatives and integrals and their applications in fractional calculus.

Definition 2.1. The Caputo explained the fractional-order derivative of function $F(t)$ as

$$D_t^\theta (F(t)) = \frac{1}{\Gamma(\lambda - \theta)} \int_0^t (t - \rho)^{\lambda - \theta - 1} F^\lambda(\rho) d\rho \text{ for } \lambda - 1 < \theta \leq \lambda; \lambda \in N; t > 0. \quad (3)$$

Definition 2.2. The RL derived the definition of fractional-order derivative of $F(t)$ as

$$D_t^\theta (F(t)) = \frac{1}{\Gamma(\lambda - \theta)} \frac{d^\lambda}{dt^\lambda} \int_0^t (t - \rho)^{\lambda - \theta - 1} F(\rho) d\rho \text{ for } \lambda - 1 < \theta \leq \lambda; \lambda \in N; t > 0. \quad (4)$$

Definition 2.3. Let the function $u(x, t)$ with variables ‘ x ’ and $t > 0$ then RL fractional partial order derivative is proposed as

$$\partial_t^\theta(u(x, t)) = \begin{cases} \frac{1}{\Gamma(\lambda - \mu)} \frac{\partial^\lambda}{\partial t^\lambda} \int_0^t (t - \rho)^{\lambda - \theta - 1} u(\rho, x) d\rho & \text{for } \lambda - 1 < \theta < \lambda, \lambda \in N, \\ \frac{\partial^\lambda u}{\partial t^\lambda} & \text{for } \theta = \lambda. \end{cases} \quad (5)$$

Definition 2.4. The Leibnitz described the product rule under application of RL fractional-order derivatives in the form

$$D_t^\theta(U.V) = \sum_{\lambda=0}^{\infty} \binom{\theta}{\lambda} D_t^{\theta-\lambda}(U).D_t^\lambda(V) \quad ; \theta > 0 \text{ with } \binom{\theta}{\lambda} = \frac{(-1)^\lambda \theta \Gamma(n - \theta)}{\Gamma(1 - \theta) \Gamma(\theta + 1)}. \quad (6)$$

Definition 2.5. The E-Kober generalized fractional differential operator $(E_\partial^{\tau, \mu} \omega)(\zeta)$ is given by

$$(E_\partial^{\tau, \mu} \omega)(\zeta) = \prod_{\lambda=0}^{m-1} \left(\tau + \lambda - \frac{1}{\partial} z \frac{d}{dz} \right) (K_\partial^{\tau+\mu, m-\mu} \omega)(z) \text{ with } \zeta > 0, \partial > 0 \text{ and } \mu > 0; \quad (7)$$

$$m = \begin{cases} [\mu] + 1, & \mu \notin N \\ \mu, & \mu \in N \end{cases}$$

Definition 2.6. E-Kober generalized fractional-order integral operator $(K_\partial^{\tau, \mu} \omega)(\zeta)$ is

$$(K_\partial^{\tau, \mu} \omega)(\zeta) = \begin{cases} \frac{1}{\Gamma(\mu)} \int_1^\infty (v-1)^{\mu-1} v^{-(\tau+\mu)} \omega(\zeta v^{1/\partial}) dv, & \mu > 0, \\ \omega(\zeta) & , \quad \mu = 0. \end{cases} \quad (8)$$

3. Methodology

In this section, authors would like to pursue the steps and process of fractional Lie symmetry reduction to coupled time fractional order system of FPDEs.

Let us assume the system of FPDEs with fractional order θ

$$\begin{cases} \partial_t^\theta u = F_1(t, x, u, v, w, u_x, v_x, w_x, z_x, u_{xx}, v_{xx}, w_{xx}, z_{xx} \dots); \\ \partial_t^\theta v = F_2(t, x, u, v, w, u_x, v_x, w_x, z_x, u_{xx}, v_{xx}, w_{xx}, z_{xx} \dots); \\ \partial_t^\theta w = F_3(t, x, u, v, w, u_x, v_x, w_x, z_x, u_{xx}, v_{xx}, w_{xx}, z_{xx} \dots); \\ \partial_t^\theta z = F_4(t, x, u, v, w, u_x, v_x, w_x, z_x, u_{xx}, v_{xx}, w_{xx}, z_{xx} \dots); \end{cases} \quad 0 < \theta < 1. \quad (9)$$

The infinitesimal transformations with single parametric notation in fractional Lie symmetry analysis is expressed as:

$$\begin{cases} \bar{t} = \bar{t}(x, t, u, v, w, z; \varepsilon) = t + \varepsilon \tau(x, t, u, v, w, z) + o(\varepsilon^2); \\ \bar{x} = \bar{x}(x, t, u, v, w, z; \varepsilon) = x + \varepsilon \xi(x, t, u, v, w, z) + o(\varepsilon^2); \\ \bar{u} = \bar{u}(x, t, u, v, w, z; \varepsilon) = u + \varepsilon \eta(x, t, u, v, w, z) + o(\varepsilon^2); \\ \bar{v} = \bar{v}(x, t, u, v, w, z; \varepsilon) = v + \varepsilon \phi(x, t, u, v, w, z) + o(\varepsilon^2); \\ \bar{w} = \bar{w}(x, t, u, v, w, z; \varepsilon) = w + \varepsilon \mu(x, t, u, v, w, z) + o(\varepsilon^2); \\ \bar{z} = \bar{z}(x, t, u, v, w, z; \varepsilon) = z + \varepsilon \mu(x, t, u, v, w, z) + o(\varepsilon^2). \end{cases} \quad (10)$$

The vector field generated by infinitesimals is taken as:

$$X = \tau \partial_t + \xi \partial_x + \eta_1 \partial_u + \eta_2 \partial_v + \eta_3 \partial_w + \eta_4 \partial_z \quad (11)$$

$$\text{with } \tau = \left. \frac{d\bar{t}}{d\varepsilon} \right|_{\varepsilon=0}, \quad \xi = \left. \frac{d\bar{x}}{d\varepsilon} \right|_{\varepsilon=0}, \quad \eta_1 = \left. \frac{d\bar{u}}{d\varepsilon} \right|_{\varepsilon=0}, \quad \eta_2 = \left. \frac{d\bar{v}}{d\varepsilon} \right|_{\varepsilon=0}, \quad \eta_3 = \left. \frac{d\bar{w}}{d\varepsilon} \right|_{\varepsilon=0}, \quad \eta_4 = \left. \frac{d\bar{z}}{d\varepsilon} \right|_{\varepsilon=0}.$$

Here, $\xi, \tau, \eta_1, \eta_2, \eta_3$ and η_4 are obtained infinitesimals operators from (11), $\eta_1^{\theta,t}, \eta_2^{\theta,t}, \eta_3^{\theta,t}$ and $\eta_4^{\theta,t}$ are the fractional extended infinitesimals of order θ and $\eta_1^x, \eta_1^{xx}, \eta_1^{xxx}, \eta_2^x, \eta_2^{xx}, \eta_2^{xxx}, \eta_3^x, \eta_3^{xx}, \eta_3^{xxx}$ and $\eta_4^x, \eta_4^{xx}, \eta_4^{xxx}$ are extended infinitesimals of integer-order described

$$\begin{aligned}
\eta_1^x &= D_x(\eta_1) - u_x D_x(\xi) - u_t D_x(\tau) ; \\
\eta_1^{xx} &= D_x(\eta_1^x) - u_{xx} D_x(\xi) - u_{xt} D_x(\tau) ; \\
\eta_1^{xxx} &= D_x(\eta_1^{xx}) - u_{xxx} D_x(\xi) - u_{xxt} D_x(\tau) ; \\
\eta_2^x &= D_x(\eta_2) - v_x D_x(\xi) - v_t D_x(\tau) ; \\
\eta_2^{xx} &= D_x(\eta_2^x) - v_{xx} D_x(\xi) - v_{xt} D_x(\tau) ; \\
\eta_2^{xxx} &= D_x(\eta_2^{xx}) - v_{xxx} D_x(\xi) - v_{xxt} D_x(\tau) ; \\
\eta_3^x &= D_x(\eta_3) - w_x D_x(\xi) - w_t D_x(\tau) ; \\
\eta_3^{xx} &= D_x(\eta_3^x) - w_{xx} D_x(\xi) - w_{xt} D_x(\tau) ; \\
\eta_3^{xxx} &= D_x(\eta_3^{xx}) - w_{xxx} D_x(\xi) - w_{xxt} D_x(\tau) ; \\
\eta_4^x &= D_x(\eta_4) - z_x D_x(\xi) - z_t D_x(\tau) ; \\
\eta_4^{xx} &= D_x(\eta_4^x) - z_{xx} D_x(\xi) - z_{xt} D_x(\tau) ; \\
\eta_4^{xxx} &= D_x(\eta_4^{xx}) - z_{xxx} D_x(\xi) - z_{xxt} D_x(\tau),
\end{aligned} \tag{12}$$

where ' D_x ' is total derivative operator defined as:

$$D_x = \frac{\partial}{\partial x} + u_x \frac{\partial}{\partial u} + u_{xx} \frac{\partial}{\partial u_{xx}} + \dots + v_x \frac{\partial}{\partial v_x} + v_{xx} \frac{\partial}{\partial v_{xx}} + \dots + w_x \frac{\partial}{\partial w_x} + w_{xx} \frac{\partial}{\partial w_{xx}} + \dots \tag{13}$$

The extended infinitesimal function of θ -th order $\eta_1^{\theta,t}$ concerned to RL fractional derivative is described by:

$$\eta_1^{\theta,t} = D_t^\theta(\eta_1) + \xi.D_t^\theta(u_x) - D_t^\theta(\xi u_x) + D_t^\theta(D_t(\tau)u) - D_t^{\theta+1}(\tau u) + \tau.D_t^{\theta+1}(u). \tag{14}$$

Also, $D_t^{\theta+1}(f(t)) = D_t^\theta(D_t(f(t)))$, then above expression simplified to

$$\eta_1^{\theta,t} = D_t^\theta(\eta_1) + \xi.D_t^\theta(u_x) - D_t^\theta(\xi u_x) + \tau.D_t^\theta(u) - D_t^\theta(\tau u_t). \tag{15}$$

Applying the generalized Leibnitz rule on (15), we obtain

$$\eta_1^{\theta,t} = D_t^\theta(\eta_1) - \theta.D_t^\theta(\tau) \frac{\partial^\theta u}{\partial t^\theta} - \sum_{\lambda=1}^{\infty} \binom{\theta}{\lambda} D_t^\lambda(\xi) D_t^{\theta-\lambda}(u_x) - \sum_{\lambda=1}^{\infty} \binom{\theta}{\lambda+1} D_t^{\lambda+1}(\tau) D_t^{\theta-\lambda}(u). \tag{16}$$

Using the generalized Leibnitz rule and chain rule (5) the term $D_t^\theta(\eta)$ in (16) can be defined as

$$\begin{aligned}
D_t^\theta(\eta_1) = & \frac{\partial^\theta \eta_1}{\partial t^\theta} + \left(\eta_{1u} \frac{\partial^\theta u}{\partial t^\theta} - u \frac{\partial^\theta(\eta_{1u})}{\partial t^\theta} \right) + \left(\eta_{1v} \frac{\partial^\theta v}{\partial t^\theta} - v \frac{\partial^\theta(\eta_{1v})}{\partial t^\theta} \right) + \left(\eta_{1w} \frac{\partial^\theta w}{\partial t^\theta} - w \frac{\partial^\theta(\eta_{1w})}{\partial t^\theta} \right) \\
& \left(\eta_{1z} \frac{\partial^\theta z}{\partial t^\theta} - z \frac{\partial^\theta(\eta_{1z})}{\partial t^\theta} \right) + \sum_{\lambda=1}^{\infty} \binom{\theta}{\lambda} \frac{\partial^\lambda(\eta_{1u})}{\partial t^\lambda} D_t^{\theta-\lambda}(u) + \sum_{\lambda=1}^{\infty} \binom{\theta}{\lambda} \frac{\partial^\lambda(\eta_{1v})}{\partial t^\lambda} D_t^{\theta-\lambda}(v) + \\
& \sum_{\lambda=1}^{\infty} \binom{\theta}{\lambda} \frac{\partial^\lambda(\eta_{1w})}{\partial t^\lambda} D_t^{\theta-\lambda}(w) + \sum_{\lambda=1}^{\infty} \binom{\theta}{\lambda} \frac{\partial^\lambda(\eta_{1z})}{\partial t^\lambda} D_t^{\theta-\lambda}(z) + \sigma_1 + \sigma_2 + \sigma_3 + \sigma_4,
\end{aligned} \tag{17}$$

where,

$$\begin{cases}
\sigma_1 = \sum_{\lambda=2}^{\infty} \sum_{m=2}^{\lambda} \sum_{k=2}^m \sum_{r=0}^{k-1} \binom{\theta}{\lambda} \binom{\lambda}{m} \binom{k}{r} \frac{t^{\lambda-\theta}}{k! \Gamma(\lambda+1-\theta)} (-u)^r \frac{\partial^m}{\partial t^m} (u^{k-r}) \frac{\partial^{\lambda-m+k} \eta_1}{\partial t^{\lambda-m} \partial u^k}, \\
\sigma_2 = \sum_{\lambda=2}^{\infty} \sum_{m=2}^{\lambda} \sum_{k=2}^m \sum_{r=0}^{k-1} \binom{\theta}{\lambda} \binom{\lambda}{m} \binom{k}{r} \frac{t^{\lambda-\theta}}{k! \Gamma(\lambda+1-\theta)} (-v)^r \frac{\partial^m}{\partial t^m} (v^{k-r}) \frac{\partial^{\lambda-m+k} \eta_1}{\partial t^{\lambda-m} \partial v^k}, \\
\sigma_3 = \sum_{\lambda=2}^{\infty} \sum_{m=2}^{\lambda} \sum_{k=2}^m \sum_{r=0}^{k-1} \binom{\theta}{\lambda} \binom{\lambda}{m} \binom{k}{r} \frac{t^{\lambda-\theta}}{k! \Gamma(\lambda+1-\theta)} (-w)^r \frac{\partial^m}{\partial t^m} (w^{k-r}) \frac{\partial^{\lambda-m+k} \eta_1}{\partial t^{\lambda-m} \partial w^k}, \\
\sigma_4 = \sum_{\lambda=2}^{\infty} \sum_{m=2}^{\lambda} \sum_{k=2}^m \sum_{r=0}^{k-1} \binom{\theta}{\lambda} \binom{\lambda}{m} \binom{k}{r} \frac{t^{\lambda-\theta}}{k! \Gamma(\lambda+1-\theta)} (-z)^r \frac{\partial^m}{\partial t^m} (z^{k-r}) \frac{\partial^{\lambda-m+k} \eta_1}{\partial t^{\lambda-m} \partial z^k}.
\end{cases} \tag{18}$$

Finally, the expression for θ -th order extended infinitesimal $\eta_1^{\theta,t}$ of the form

$$\begin{aligned}
\eta_1^{\theta,t} = & \frac{\partial^\theta \eta_1}{\partial t^\theta} + (\eta_{1u} - \theta D_t(\tau)) \frac{\partial^\theta u}{\partial t^\theta} - u \frac{\partial^\theta(\eta_{1u})}{\partial t^\theta} + \left(\eta_{1v} \frac{\partial^\theta v}{\partial t^\theta} - v \frac{\partial^\theta(\eta_{1v})}{\partial t^\theta} \right) + \\
& \left(\eta_{1w} \frac{\partial^\theta w}{\partial t^\theta} - w \frac{\partial^\theta(\eta_{1w})}{\partial t^\theta} \right) + \left(\eta_{1z} \frac{\partial^\theta z}{\partial t^\theta} - z \frac{\partial^\theta(\eta_{1z})}{\partial t^\theta} \right) + \\
& \sum_{\lambda=1}^{\infty} \left[\binom{\theta}{\lambda} \frac{\partial^\lambda \eta_{1u}}{\partial t^\lambda} - \binom{\theta}{\lambda+1} D_t^{\lambda+1}(\tau) \right] D_t^{\theta-\lambda}(u) + \sum_{\lambda=1}^{\infty} \binom{\theta}{\lambda} \frac{\partial^\lambda(\eta_{1v})}{\partial t^\lambda} D_t^{\theta-\lambda}(v) \\
& + \sum_{\lambda=1}^{\infty} \binom{\theta}{\lambda} \frac{\partial^\lambda(\eta_{1w})}{\partial t^\lambda} D_t^{\theta-\lambda}(w) + \sum_{\lambda=1}^{\infty} \binom{\theta}{\lambda} \frac{\partial^\lambda(\eta_{1z})}{\partial t^\lambda} D_t^{\theta-\lambda}(z) \\
& - \sum_{\lambda=1}^{\infty} \binom{\theta}{\lambda} D_t^\lambda(\xi) D_t^{\theta-\lambda}(u_x) + \sigma_1 + \sigma_2 + \sigma_3 + \sigma_4.
\end{aligned} \tag{19}$$

Similarly, expressions for $\eta_2^{\theta,t}$, $\eta_3^{\theta,t}$ and $\eta_4^{\theta,t}$ are also obtained

$$\begin{aligned}
\eta_2^{\theta,t} &= \frac{\partial^\theta \eta_2}{\partial t^\theta} + (\eta_{2v} - \theta D_t(\tau)) \frac{\partial^\theta v}{\partial t^\theta} - v \frac{\partial^\theta (\eta_{2v})}{\partial t^\theta} + \left(\eta_{2u} \frac{\partial^\theta u}{\partial t^\theta} - u \frac{\partial^\theta (\eta_{2u})}{\partial t^\theta} \right) + \\
&\quad \left(\eta_{2w} \frac{\partial^\theta w}{\partial t^\theta} - u \frac{\partial^\theta (\eta_{2w})}{\partial t^\theta} \right) + \left(\eta_{2z} \frac{\partial^\theta z}{\partial t^\theta} - u \frac{\partial^\theta (\eta_{2z})}{\partial t^\theta} \right) + \\
&\quad \sum_{\lambda=1}^{\infty} \left[\left(\frac{\theta}{\lambda} \right) \frac{\partial^n \eta_{2v}}{\partial t^n} - \left(\frac{\theta}{\lambda+1} \right) D_t^{\lambda+1}(\tau) \right] D_t^{\theta-\lambda}(v) + \sum_{\lambda=1}^{\infty} \left(\frac{\theta}{\lambda} \right) \frac{\partial^\lambda (\eta_{2u})}{\partial t^\lambda} D_t^{\theta-\lambda}(u) + \\
&\quad \sum_{\lambda=1}^{\infty} \left(\frac{\theta}{\lambda} \right) \frac{\partial^\lambda (\eta_{2w})}{\partial t^\lambda} D_t^{\theta-\lambda}(w) + \sum_{\lambda=1}^{\infty} \left(\frac{\theta}{\lambda} \right) \frac{\partial^\lambda (\eta_{2z})}{\partial t^\lambda} D_t^{\theta-\lambda}(z) - \\
&\quad \sum_{\lambda=1}^{\infty} \left(\frac{\theta}{\lambda} \right) D_t^\lambda(\xi) D_t^{\theta-\lambda}(v_x) + \sigma_5 + \sigma_6 + \sigma_7 + \sigma_8.
\end{aligned} \tag{20}$$

$$\begin{aligned}
\eta_3^{\theta,t} &= \frac{\partial^\theta \eta_3}{\partial t^\theta} + (\eta_{3w} - \theta D_t(\tau)) \frac{\partial^\theta w}{\partial t^\theta} - w \frac{\partial^\theta (\eta_{3w})}{\partial t^\theta} + \left(\eta_{3u} \frac{\partial^\theta u}{\partial t^\theta} - u \frac{\partial^\theta (\eta_{3u})}{\partial t^\theta} \right) + \\
&\quad \left(\eta_{3v} \frac{\partial^\theta v}{\partial t^\theta} - u \frac{\partial^\theta (\eta_{3v})}{\partial t^\theta} \right) + \left(\eta_{3z} \frac{\partial^\theta z}{\partial t^\theta} - u \frac{\partial^\theta (\eta_{3z})}{\partial t^\theta} \right) + \\
&\quad \sum_{\lambda=1}^{\infty} \left[\left(\frac{\theta}{\lambda} \right) \frac{\partial^n \eta_{3w}}{\partial t^n} - \left(\frac{\theta}{\lambda+1} \right) D_t^{\lambda+1}(\tau) \right] D_t^{\theta-\lambda}(w) + \sum_{\lambda=1}^{\infty} \left(\frac{\theta}{\lambda} \right) \frac{\partial^\lambda (\eta_{3u})}{\partial t^\lambda} D_t^{\theta-\lambda}(u) \\
&\quad + \sum_{\lambda=1}^{\infty} \left(\frac{\theta}{\lambda} \right) \frac{\partial^\lambda (\eta_{3v})}{\partial t^\lambda} D_t^{\theta-\lambda}(v) + \sum_{\lambda=1}^{\infty} \left(\frac{\theta}{\lambda} \right) \frac{\partial^\lambda (\eta_{3z})}{\partial t^\lambda} D_t^{\theta-\lambda}(z) - \\
&\quad \sum_{\lambda=1}^{\infty} \left(\frac{\theta}{\lambda} \right) D_t^\lambda(\xi) D_t^{\theta-\lambda}(w_x) + \sigma_9 + \sigma_{10} + \sigma_{11} + \sigma_{12}.
\end{aligned} \tag{21}$$

and

$$\begin{aligned}
\eta_4^{\theta,t} &= \frac{\partial^\theta \eta_4}{\partial t^\theta} + (\eta_{4z} - \theta D_t(\tau)) \frac{\partial^\theta z}{\partial t^\theta} - z \frac{\partial^\theta (\eta_{4z})}{\partial t^\theta} + \left(\eta_{4u} \frac{\partial^\theta u}{\partial t^\theta} - u \frac{\partial^\theta (\eta_{4u})}{\partial t^\theta} \right) \\
&\quad + \left(\eta_{4w} \frac{\partial^\theta w}{\partial t^\theta} - u \frac{\partial^\theta (\eta_{4w})}{\partial t^\theta} \right) + \left(\eta_{4v} \frac{\partial^\theta z}{\partial t^\theta} - v \frac{\partial^\theta (\eta_{4z})}{\partial t^\theta} \right) \\
&\quad + \sum_{\lambda=1}^{\infty} \left[\left(\frac{\theta}{\lambda} \right) \frac{\partial^n \eta_{4z}}{\partial t^n} - \left(\frac{\theta}{\lambda+1} \right) D_t^{\lambda+1}(\tau) \right] D_t^{\theta-\lambda}(z) + \sum_{\lambda=1}^{\infty} \left(\frac{\theta}{\lambda} \right) \frac{\partial^\lambda (\eta_{4u})}{\partial t^\lambda} D_t^{\theta-\lambda}(u) \\
&\quad + \sum_{\lambda=1}^{\infty} \left(\frac{\theta}{\lambda} \right) \frac{\partial^\lambda (\eta_{4w})}{\partial t^\lambda} D_t^{\theta-\lambda}(w) + \sum_{\lambda=1}^{\infty} \left(\frac{\theta}{\lambda} \right) \frac{\partial^\lambda (\eta_{4v})}{\partial t^\lambda} D_t^{\theta-\lambda}(v) \\
&\quad - \sum_{\lambda=1}^{\infty} \left(\frac{\theta}{\lambda} \right) D_t^\lambda(\xi) D_t^{\theta-\lambda}(z_x) + \sigma_{13} + \sigma_{14} + \sigma_{15} + \sigma_{16}.
\end{aligned} \tag{22}$$

$$\sigma_i = \sum_{\lambda=2}^{\infty} \sum_{m=2}^{\lambda} \sum_{k=2r=0}^m \sum_{k-1}^{\lambda} \binom{\theta}{\lambda} \binom{\lambda}{m} \binom{k}{r} \frac{t^{\lambda-\theta}}{k! \Gamma(\lambda+1-\theta)} (-u)^r \frac{\partial^m}{\partial t^m} (u^{k-r}) \frac{\partial^{\lambda-m+k} \eta}{\partial t^{\lambda-m} \partial u^k}. \quad (23)$$

For expressions σ_i ; $i=5, 6, 7 \dots 16$ given by (23) vanishes.

4. Lie symmetry reduction of nonlinear system of Hirota equations

Applying prolongation on set of Hirota system of FPDEs (2), we obtain

$$\begin{cases} \eta_1^{\theta,t} + \eta_1^{xxx} + 6\eta_1^x \left(\sum u^2 \right) + 12u_x (u\eta_1 + v\eta_2 + w\eta_3 + z\eta_4) = 0; \\ \eta_2^{\theta,t} + \eta_2^{xxx} + 6\eta_2^x \left(\sum u^2 \right) + 12v_x (u\eta_1 + v\eta_2 + w\eta_3 + z\eta_4) = 0; \\ \eta_3^{\theta,t} + \eta_3^{xxx} + 6\eta_3^x \left(\sum u^2 \right) + 12w_x (u\eta_1 + v\eta_2 + w\eta_3 + z\eta_4) = 0; \\ \eta_4^{\theta,t} + \eta_4^{xxx} + 6\eta_4^x \left(\sum u^2 \right) + 12z_x (u\eta_1 + v\eta_2 + w\eta_3 + z\eta_4) = 0. \end{cases} \quad (24)$$

Using above explained facts in the methodology, substituting the values of extended infinitesimals and solving system of obtained fractional PDEs, we obtain the following infinitesimals

$$\begin{aligned} \xi &= \frac{p_1 x}{3} + p_2, \quad \tau = \frac{p_1 t}{\theta}, \quad \eta_1 = \frac{-p_1 u}{3} + p_3 v + p_4 w + p_5 z, \\ \eta_2 &= -p_3 u - \frac{p_1 v}{3} + p_6 w + p_7 z, \\ \eta_3 &= -p_4 u - p_6 v - \frac{p_1 w}{3} + p_8 z, \\ \eta_4 &= -p_5 u - p_7 v - p_8 w - \frac{p_1 z}{3}, \end{aligned} \quad (25)$$

where $p_i (i=1, 2, \dots, 8)$ are components of standard basis of vector field

$$X_i = \xi \partial_x + \tau \partial_t + \eta_1 \partial_u + \eta_2 \partial_v + \eta_3 \partial_w + \eta_4 \partial_z. \quad (26)$$

Lie algebra has the following vectors

$$\begin{aligned}
X_1 &= \frac{x}{3}\partial_x + \frac{t}{\theta}\partial_t - \frac{u}{3}\partial_u - \frac{v}{3}\partial_v - \frac{w}{3}\partial_w - \frac{z}{3}\partial_z, \\
X_2 &= \partial_x, \quad X_3 = v\partial_u - u\partial_v, \quad X_4 = w\partial_u - u\partial_w, \\
X_5 &= z\partial_u - u\partial_z, \quad X_6 = w\partial_v - v\partial_w, \\
X_7 &= z\partial_v - v\partial_z, \quad X_8 = z\partial_w - w\partial_z.
\end{aligned} \tag{27}$$

We are interested to evaluate the optimal solution of system of equations by choosing first infinitesimal generator with the help of characteristic equation.

$$\frac{3dx}{x} = \frac{\theta dt}{t} = \frac{-3du}{u} = \frac{-3dv}{v} = \frac{-3dw}{w} = \frac{-3dz}{z}. \tag{28}$$

Explicit solution obtained with ‘ α ’ as similarity variable

$$u = t^{-\theta/3}f(\alpha), \quad v = t^{-\theta/3}g(\alpha), \quad w = t^{-\theta/3}h(\alpha), \quad z = t^{-\theta/3}k(\alpha) \text{ and } \alpha = xt^{-\theta/3}. \tag{29}$$

5. Application of EK differ-integral operators

Applying the R-L fractional derivative as:

$$\begin{aligned}
\partial_t^\theta u &= \partial_t^\lambda \left[\frac{1}{\Gamma(\lambda - \theta)} \int_0^t (t-s)^{\lambda-\theta-1} s^{-\theta/3} f(xs^{-\theta/3}) ds \right]; \quad q = t/s \\
&= \partial_t^\lambda \left[\frac{t^{\lambda-\frac{2\theta}{3}}}{\Gamma(\lambda - \theta)} \int_1^\infty (q-1)^{\lambda-\theta-1} q^{-\left(\lambda-\frac{2\theta}{3}+1\right)} f(\alpha q^{\theta/3}) dq \right].
\end{aligned} \tag{30}$$

Applying E-K integral operator then above expression found to be

$$\partial_t^\theta u = \partial_t^\lambda \left[t^{\lambda-\frac{2\theta}{3}} \left(\kappa_{\frac{3}{\theta}}^{1-\frac{\theta}{3}, \lambda-\theta} f \right) \alpha \right]. \tag{31}$$

Also, using $t \partial_t f(\alpha) = t x \left(-\frac{\theta}{3} \right) t^{-\frac{\theta}{3}-1} f'(\alpha) = -\frac{\theta}{3} \alpha \frac{d}{d\alpha} (f(\alpha))$ in above equation (31), we obtain

$$\begin{aligned}
\partial_t^\theta u &= \partial_t^\lambda \left(\partial_t \left[t^{\lambda - \frac{2\theta}{3}} \left(\kappa_{\frac{3}{\theta}}^{1 - \frac{\theta}{3}, \lambda - \theta} f \right) \alpha \right] \right) \\
&= \partial_t^{\lambda-1} \left[t^{\lambda - \frac{2\theta}{3} - 1} \left(\lambda - \frac{4\theta}{3} - \frac{\theta}{3} \alpha \frac{d}{d\alpha} \right) \left(\kappa_{\frac{3}{\theta}}^{1 - \frac{\theta}{3}, \lambda - \theta} f \right) \alpha \right].
\end{aligned} \tag{32}$$

Repeating above arguments $(\lambda-1)$ times and applying E-K differential operator, we obtain

$$\partial_t^\theta u = t^{-\frac{2\theta}{3}} \left(P_{\frac{3}{\theta}}^{1 - \frac{4\theta}{3}, \theta} f \right) \alpha. \tag{33}$$

System of FPDEs is reduced to system of FODEs which is given by

$$\begin{cases}
\left(P_{\frac{3}{\theta}}^{1 - \frac{4\theta}{3}, \theta} f \right) \alpha + f'''(\alpha) + 6f'(\alpha) \left(\sum f^2 \right) = 0; \\
\left(P_{\frac{3}{\theta}}^{1 - \frac{4\theta}{3}, \theta} g \right) \alpha + g'''(\alpha) + 6g'(\alpha) \left(\sum f^2 \right) = 0; \\
\left(P_{\frac{3}{\theta}}^{1 - \frac{4\theta}{3}, \theta} h \right) \alpha + h'''(\alpha) + 6h'(\alpha) \left(\sum f^2 \right) = 0; \\
\left(P_{\frac{3}{\theta}}^{1 - \frac{4\theta}{3}, \theta} k \right) \alpha + k'''(\alpha) + 6k'(\alpha) \left(\sum f^2 \right) = 0,
\end{cases} \tag{34}$$

$$\text{where } \sum f^2 = f^2 + g^2 + h^2 + k^2.$$

6. Power series solution of system and its convergence:

$$\text{Set } f(\alpha) = \sum_{n=1}^{\infty} a_n \alpha^n, \quad g(\alpha) = \sum_{n=1}^{\infty} b_n \alpha^n, \quad h(\alpha) = \sum_{n=1}^{\infty} c_n \alpha^n, \quad k(\alpha) = \sum_{n=1}^{\infty} d_n \alpha^n. \tag{35}$$

From above equation, we obtained

$$\left\{ \begin{array}{l} \sum_{n=0}^{\infty} \frac{\Gamma\left(2 - \frac{4\theta}{3} + \frac{n\theta}{3}\right)}{\Gamma\left(2 - \frac{\theta}{3} + \frac{n\theta}{3}\right)} a_n \alpha^n + \sum_{n=0}^{\infty} (n+3)(n+2)(n+1) a_{n+3} \alpha^n + 6 \left(\sum_{n=0}^{\infty} (n+1) a_{n+1} \alpha^n \right) [S] = 0; \\ \sum_{n=0}^{\infty} \frac{\Gamma\left(2 - \frac{4\theta}{3} + \frac{n\theta}{3}\right)}{\Gamma\left(2 - \frac{\theta}{3} + \frac{n\theta}{3}\right)} b_n \alpha^n + \sum_{n=0}^{\infty} (n+3)(n+2)(n+1) b_{n+3} \alpha^n + 6 \left(\sum_{n=0}^{\infty} (n+1) b_{n+1} \alpha^n \right) [S] = 0; \\ \sum_{n=0}^{\infty} \frac{\Gamma\left(2 - \frac{4\theta}{3} + \frac{n\theta}{3}\right)}{\Gamma\left(2 - \frac{\theta}{3} + \frac{n\theta}{3}\right)} c_n \alpha^n + \sum_{n=0}^{\infty} (n+3)(n+2)(n+1) c_{n+3} \alpha^n + 6 \left(\sum_{n=0}^{\infty} (n+1) c_{n+1} \alpha^n \right) [S] = 0; \\ \sum_{n=0}^{\infty} \frac{\Gamma\left(2 - \frac{4\theta}{3} + \frac{n\theta}{3}\right)}{\Gamma\left(2 - \frac{\theta}{3} + \frac{n\theta}{3}\right)} d_n \alpha^n + \sum_{n=0}^{\infty} (n+3)(n+2)(n+1) d_{n+3} \alpha^n + 6 \left(\sum_{n=0}^{\infty} (n+1) d_{n+1} \alpha^n \right) [S] = 0, \end{array} \right. \quad (36)$$

$$\text{where } S = \left(\sum_{n=1}^{\infty} a_n \alpha^n \right)^2 + \left(\sum_{n=1}^{\infty} b_n \alpha^n \right)^2 + \left(\sum_{n=1}^{\infty} c_n \alpha^n \right)^2 + \left(\sum_{n=1}^{\infty} d_n \alpha^n \right)^2. \quad (37)$$

Substituting $n=0$ in (36), we get

$$\left\{ \begin{array}{l} a_3 = -\frac{\Gamma(2-4\theta/3)}{3!\Gamma(2-\theta/3)} a_0 - a_1 S_0; \\ b_3 = -\frac{\Gamma(2-4\theta/3)}{3!\Gamma(2-\theta/3)} b_0 - b_1 S_0; \\ c_3 = -\frac{\Gamma(2-4\theta/3)}{3!\Gamma(2-\theta/3)} c_0 - c_1 S_0; \\ d_3 = -\frac{\Gamma(2-4\theta/3)}{3!\Gamma(2-\theta/3)} d_0 - d_1 S_0, \end{array} \right. \quad \text{where } S_0 = a_0^2 + b_0^2 + c_0^2 + d_0^2, \quad (38)$$

and

$$\left\{ \begin{aligned} a_{n+3} &= \frac{n!}{(n+3)!} \frac{\Gamma\left(2 - \frac{4\theta}{3} + \frac{n\theta}{3}\right)}{\Gamma\left(2 - \frac{\theta}{3} + \frac{n\theta}{3}\right)} a_n - \frac{6}{(n+3)(n+2)} \sum_{k=0}^n a_{n+1-k} (a_{n-k}^2 + b_{n-k}^2 + c_{n-k}^2 + d_{n-k}^2) \\ b_{n+3} &= \frac{n!}{(n+3)!} \frac{\Gamma\left(2 - \frac{4\theta}{3} + \frac{n\theta}{3}\right)}{\Gamma\left(2 - \frac{\theta}{3} + \frac{n\theta}{3}\right)} b_n - \frac{6}{(n+3)(n+2)} \sum_{k=0}^n b_{n+1-k} (a_{n-k}^2 + b_{n-k}^2 + c_{n-k}^2 + d_{n-k}^2) \\ c_{n+3} &= \frac{n!}{(n+3)!} \frac{\Gamma\left(2 - \frac{4\theta}{3} + \frac{n\theta}{3}\right)}{\Gamma\left(2 - \frac{\theta}{3} + \frac{n\theta}{3}\right)} c_n - \frac{6}{(n+3)(n+2)} \sum_{k=0}^n c_{n+1-k} (a_{n-k}^2 + b_{n-k}^2 + c_{n-k}^2 + d_{n-k}^2) \\ d_{n+3} &= \frac{n!}{(n+3)!} \frac{\Gamma\left(2 - \frac{4\theta}{3} + \frac{n\theta}{3}\right)}{\Gamma\left(2 - \frac{\theta}{3} + \frac{n\theta}{3}\right)} d_n - \frac{6}{(n+3)(n+2)} \sum_{k=0}^n d_{n+1-k} (a_{n-k}^2 + b_{n-k}^2 + c_{n-k}^2 + d_{n-k}^2) \end{aligned} \right. \quad (39)$$

So the explicit power series solution of system is given by:

$$\left\{ \begin{aligned} f(\alpha) &= a_0 + a_1(xt^{-\theta/3}) + a_2(xt^{-\theta/3})^2 + \left[-\frac{\Gamma(2-4\theta/3)}{3!\Gamma(2-\theta/3)} a_0 - a_1 S_0 \right] (xt^{-\theta/3})^3 + \sum_{n=1}^{\infty} a_{n+3} (xt^{-\theta/3})^{n+3} \\ g(\alpha) &= b_0 + b_1(xt^{-\theta/3}) + b_2(xt^{-\theta/3})^2 + \left[-\frac{\Gamma(2-4\theta/3)}{3!\Gamma(2-\theta/3)} b_0 - b_1 S_0 \right] (xt^{-\theta/3})^3 + \sum_{n=1}^{\infty} b_{n+3} (xt^{-\theta/3})^{n+3} \\ h(\alpha) &= c_0 + c_1(xt^{-\theta/3}) + c_2(xt^{-\theta/3})^2 + \left[-\frac{\Gamma(2-4\theta/3)}{3!\Gamma(2-\theta/3)} c_0 - c_1 S_0 \right] (xt^{-\theta/3})^3 + \sum_{n=1}^{\infty} c_{n+3} (xt^{-\theta/3})^{n+3} \\ k(\alpha) &= d_0 + d_1(xt^{-\theta/3}) + d_2(xt^{-\theta/3})^2 + \left[-\frac{\Gamma(2-4\theta/3)}{3!\Gamma(2-\theta/3)} d_0 - d_1 S_0 \right] (xt^{-\theta/3})^3 + \sum_{n=1}^{\infty} d_{n+3} (xt^{-\theta/3})^{n+3} \end{aligned} \right. \quad (40)$$

$$\left\{ \begin{aligned} |a_{n+3}| &= \left| \frac{n!}{(n+3)!} \frac{\Gamma\left(2 - \frac{4\theta}{3} + \frac{n\theta}{3}\right)}{\Gamma\left(2 - \frac{\theta}{3} + \frac{n\theta}{3}\right)} \right| |a_n| - \left| \frac{6}{(n+3)(n+2)} \right| \left\| \sum_{k=0}^n a_{n+1-k} (a_{n-k}^2 + b_{n-k}^2 + c_{n-k}^2 + d_{n-k}^2) \right\|; \\ |b_{n+3}| &= \left| \frac{n!}{(n+3)!} \frac{\Gamma\left(2 - \frac{4\theta}{3} + \frac{n\theta}{3}\right)}{\Gamma\left(2 - \frac{\theta}{3} + \frac{n\theta}{3}\right)} \right| |b_n| - \left| \frac{6}{(n+3)(n+2)} \right| \left\| \sum_{k=0}^n b_{n+1-k} (a_{n-k}^2 + b_{n-k}^2 + c_{n-k}^2 + d_{n-k}^2) \right\|; \\ |c_{n+3}| &= \left| \frac{n!}{(n+3)!} \frac{\Gamma\left(2 - \frac{4\theta}{3} + \frac{n\theta}{3}\right)}{\Gamma\left(2 - \frac{\theta}{3} + \frac{n\theta}{3}\right)} \right| |c_n| - \left| \frac{6}{(n+3)(n+2)} \right| \left\| \sum_{k=0}^n c_{n+1-k} (a_{n-k}^2 + b_{n-k}^2 + c_{n-k}^2 + d_{n-k}^2) \right\|; \\ |d_{n+3}| &= \left| \frac{n!}{(n+3)!} \frac{\Gamma\left(2 - \frac{4\theta}{3} + \frac{n\theta}{3}\right)}{\Gamma\left(2 - \frac{\theta}{3} + \frac{n\theta}{3}\right)} \right| |d_n| - \left| \frac{6}{(n+3)(n+2)} \right| \left\| \sum_{k=0}^n d_{n+1-k} (a_{n-k}^2 + b_{n-k}^2 + c_{n-k}^2 + d_{n-k}^2) \right\|. \end{aligned} \right. \quad (41)$$

It is well known that for $n < m$, the expressions like $\left| \frac{\Gamma(n)}{\Gamma(m)} \right| < 1$ and ‘M’ is maximum of the arbitrary coefficients involved in the set of equations with introduction of majorant series. Now, we introduce some another power series

$$U(\chi) = \sum_{n=0}^{\infty} p_n \chi^n; V(\chi) = \sum_{n=0}^{\infty} q_n \chi^n; W(\chi) = \sum_{n=0}^{\infty} r_n \chi^n \text{ and } Y(\chi) = \sum_{n=0}^{\infty} s_n \chi^n, \quad (42)$$

where, $p_n = |a_n|$, $q_n = |b_n|$, $r_n = |c_n|$, $s_n = |d_n|$, $n = 1, 2, 3, \dots$, then, we can have

$$\left\{ \begin{aligned} p_{n+3} &= M \left[p_n + \sum_{k=0}^n p_{n+1-k} (p_{n-k}^2 + q_{n-k}^2 + r_{n-k}^2 + s_{n-k}^2) \right]; \\ q_{n+3} &= M \left[q_n + \sum_{k=0}^n q_{n+1-k} (p_{n-k}^2 + q_{n-k}^2 + r_{n-k}^2 + s_{n-k}^2) \right]; \\ r_{n+3} &= M \left[r_n + \sum_{k=0}^n r_{n+1-k} (p_{n-k}^2 + q_{n-k}^2 + r_{n-k}^2 + s_{n-k}^2) \right]; \\ s_{n+3} &= M \left[s_n + \sum_{k=0}^n s_{n+1-k} (p_{n-k}^2 + q_{n-k}^2 + r_{n-k}^2 + s_{n-k}^2) \right]. \end{aligned} \right. \quad (43)$$

Assuming the implicit functions of system with independent variable χ .

$$\begin{aligned}
U_1(\chi, U) &= U(\chi) - p_0 - p_1\chi - p_2\chi^2 - p_3\chi^3 - M \left[p_n + \sum_{k=0}^n p_{n+1-k} (p_{n-k}^2 + q_{n-k}^2 + r_{n-k}^2 + s_{n-k}^2) \right]; \\
V_1(\chi, V) &= V(\chi) - q_0 - q_1\chi - q_2\chi^2 - q_3\chi^3 - M \left[q_n + \sum_{k=0}^n q_{n+1-k} (p_{n-k}^2 + q_{n-k}^2 + r_{n-k}^2 + s_{n-k}^2) \right]; \\
W_1(\chi, W) &= W(\chi) - r_0 - r_1\chi - r_2\chi^2 - r_3\chi^3 - M \left[r_n + \sum_{k=0}^n r_{n+1-k} (p_{n-k}^2 + q_{n-k}^2 + r_{n-k}^2 + s_{n-k}^2) \right]; \\
Y_1(\chi, Y) &= Y(\chi) - s_0 - s_1\chi - s_2\chi^2 - s_3\chi^3 - M \left[s_n + \sum_{k=0}^n s_{n+1-k} (p_{n-k}^2 + q_{n-k}^2 + r_{n-k}^2 + s_{n-k}^2) \right].
\end{aligned} \tag{44}$$

Here, U_1 , V_1 , W_1 and Y_1 are analytic in a neighborhood of $(0, p_0)$, $(0, q_0)$, $(0, r_0)$ and $(0, s_0)$ respectively, where $U_1(0, p_0) = 0$; $V_1(0, q_0) = 0$, $W_1(0, r_0) = 0$ and $Y_1(0, s_0) = 0$ with $\frac{\partial}{\partial U}(U_1(0, p_0)) \neq 0$, $\frac{\partial}{\partial V}(V_1(0, q_0)) \neq 0$, $\frac{\partial}{\partial W}(W_1(0, r_0)) \neq 0$ and $\frac{\partial}{\partial Y}(Y_1(0, s_0)) \neq 0$ then, with Implicit function theorem [40, 50], we reached at convergence of power series solution.

7. Conservation laws

The study of classical and fractional PDEs is incomplete without the discussion of conservation laws for consistency and stability of system. We obtained the distinct conserved vectors for distinct infinitesimal generators associated with system of time fractional PDEs due to existence of one-to-one correspondence between them. These laws can be evaluated with the aid of Noether's theorem [31, 33]. For the sake of conservation laws, we should discuss the adjoint system along with Lagrangian and components of conserved vectors ω^t and ω^x , which has the continuity equation as follows:

$$D_t(\omega^t) + D_x(\omega^x) = 0. \tag{45}$$

The Lagrangian of system of FPDEs with four new dependent variables P , Q , R and S of independent variables t and x is in the form

$$\begin{aligned}
\ell = & P \left(\partial_t^\theta u + \partial_x^3 u + 6\partial_x u \left(\sum u^2 \right) \right) + Q \left(\partial_t^\theta v + \partial_x^3 v + 6\partial_x v \left(\sum u^2 \right) \right) + \\
& R \left(\partial_t^\theta w + \partial_x^3 w + 6\partial_x w \left(\sum u^2 \right) \right) + S \left(\partial_t^\theta z + \partial_x^3 z + 6\partial_x z \left(\sum u^2 \right) \right)
\end{aligned} \tag{46}$$

With Euler-Lagranges (E-L) equations also known as adjoint equations described as:

$$\frac{\delta \ell}{\delta u} = \frac{\delta \ell}{\delta v} = \frac{\delta \ell}{\delta w} = \frac{\delta \ell}{\delta z} = 0, \quad (47)$$

with E-L operators $\frac{\delta}{\delta u^i} = \frac{\partial}{\partial u^i} + (D_t^\theta)^* \frac{\partial}{\partial D_t^\theta u^i} - D_x \left(\frac{\partial}{\partial u^i_x} \right) + D_{xx} \left(\frac{\partial}{\partial u^i_{xx}} \right) - D_{xxx} \left(\frac{\partial}{\partial u^i_{xxx}} \right)$, (48)

and $(D_t^\theta)^*$ represents the adjoint operator to D_t^θ , which is defined in right-sided Caputo time-fractional derivative of order ‘ θ ’ as:

$$(D_t^\theta)^* = {}^C D_T^\theta u = \frac{(-1)^n}{\Gamma(n-\theta)} \int_t^T (v-t)^{n-1-\theta} D_v^\theta u(v, x) dv; \quad n = [\theta] + 1. \quad (49)$$

The adjoint system of equations formed by system of FPDEs

$$\begin{cases} (D_t^\theta)^* P - P_{xxx} + 6P_x \left(\sum u^2 \right) - 12P \left(\sum uu_x \right) = 0; \\ (D_t^\theta)^* Q - Q_{xxx} + 6Q_x \left(\sum u^2 \right) - 12Q \left(\sum uu_x \right) = 0; \\ (D_t^\theta)^* R - R_{xxx} + 6R_x \left(\sum u^2 \right) - 12R \left(\sum uu_x \right) = 0; \\ (D_t^\theta)^* S - S_{xxx} + 6S_x \left(\sum u^2 \right) - 12S \left(\sum uu_x \right) = 0, \end{cases}$$

where $\sum u^2 = u^2 + v^2 + w^2 + z^2$, (50)

and $\sum uu_x = uu_x + vv_x + ww_x + zz_x$,

for two independent variables x and t with dependents u , v , w and z respectively, the components of conserved vectors ω^t and ω^x are expressed as

$$\begin{aligned} \omega^x &= \xi \ell + W_j \left[\frac{\partial \ell}{\partial u_x^j} - D_x \left(\frac{\partial \ell}{\partial u_{xx}^j} \right) + D_x^2 \left(\frac{\partial \ell}{\partial u_{xxx}^j} \right) \right] + D_x(W_j) \left[\frac{\partial \ell}{\partial u_{xx}^j} - D_x \left(\frac{\partial \ell}{\partial u_{xxx}^j} \right) \right] + D_x^2(W_j) \left[\frac{\partial \ell}{\partial u_{xxx}^j} \right]; \\ \omega^t &= \tau \ell + D_t^{\theta-1}(W_j) \frac{\partial \ell}{\partial D_t^\theta u^j} + I \left(W_j, D_t^\theta \frac{\partial \ell}{\partial D_t^\theta u^j} \right), \end{aligned} \quad (51)$$

Here $W_j = \eta_j - \xi_j u_x - \tau_j u_t$, ℓ is defined above in (46) and ‘ I ’ is integral defined as

$$I(f, g) = \frac{1}{\Gamma(1-\theta)} \int_0^t \int_t^T \frac{f(s, x) g(\mu, x)}{(\mu-s)^\theta} d\mu ds. \quad (52)$$

Case 1: For X_1 , Lie characteristic functions are $W_1 = -\frac{u}{3} - \frac{x}{3}u_x - \frac{t}{\theta}u_t$, $W_2 = -\frac{v}{3} - \frac{x}{3}v_x - \frac{t}{\theta}v_t$,

$W_3 = -\frac{w}{3} - \frac{x}{3}w_x - \frac{t}{\theta}w_t$, $W_4 = -\frac{z}{3} - \frac{x}{3}z_x - \frac{t}{\theta}z_t$ and components of conserved vectors obtained as

given below:

$$\begin{aligned}\omega^x = & \frac{x}{3}\ell + \left(-\frac{u}{3} - \frac{x}{3}u_x - \frac{t}{\theta}u_t\right) \left[6P(\sum u^2) + P_{xx}\right] + P_x \left(\frac{2u_x}{3} + \frac{xu_{xx}}{3} + \frac{tu_{tx}}{\theta}\right) \\ & - P \left(\frac{u_{xx}}{3} + \frac{xu_{xxx}}{3} - \frac{tu_{txx}}{\theta}\right) + \left(-\frac{v}{3} - \frac{x}{3}v_x - \frac{t}{\theta}v_t\right) \left[6Q(\sum u^2) + Q_{xx}\right] \\ & + Q_x \left(\frac{2v_x}{3} + \frac{xv_{xx}}{3} + \frac{tv_{tx}}{\theta}\right) - Q \left(\frac{v_{xx}}{3} + \frac{xv_{xxx}}{3} - \frac{tv_{txx}}{\theta}\right) + \left(-\frac{w}{3} - \frac{x}{3}w_x - \frac{t}{\theta}w_t\right) \left[6R(\sum u^2) + R_{xx}\right] \\ & + R_x \left(\frac{2w_x}{3} + \frac{xw_{xx}}{3} + \frac{tw_{tx}}{\theta}\right) - R \left(\frac{w_{xx}}{3} + \frac{xw_{xxx}}{3} - \frac{tw_{txx}}{\theta}\right) + \left(-\frac{z}{3} - \frac{x}{3}z_x - \frac{t}{\theta}z_t\right) \left[6S(\sum u^2) + S_{xx}\right] \\ & + S_x \left(\frac{2z_x}{3} + \frac{xz_{xx}}{3} + \frac{tz_{tx}}{\theta}\right) - S \left(\frac{z_{xx}}{3} + \frac{xz_{xxx}}{3} - \frac{tz_{txx}}{\theta}\right),\end{aligned}\quad (53)$$

$$\begin{aligned}\omega^t = & PD_t^{\theta-1} \left(-\frac{u}{3} - \frac{x}{3}u_x - \frac{t}{\theta}u_t\right) + I \left(-\frac{u}{3} - \frac{x}{3}u_x - \frac{t}{\theta}u_t, P_t\right) + QD_t^{\theta-1} \left(-\frac{v}{3} - \frac{x}{3}v_x - \frac{t}{\theta}v_t\right) + \\ & I \left(-\frac{v}{3} - \frac{x}{3}v_x - \frac{t}{\theta}v_t, Q_t\right) + RD_t^{\theta-1} \left(-\frac{w}{3} - \frac{x}{3}w_x - \frac{t}{\theta}w_t\right) + I \left(-\frac{w}{3} - \frac{x}{3}w_x - \frac{t}{\theta}w_t, R_t\right) + \\ & SD_t^{\theta-1} \left(-\frac{z}{3} - \frac{x}{3}z_x - \frac{t}{\theta}z_t\right) + I \left(-\frac{z}{3} - \frac{x}{3}z_x - \frac{t}{\theta}z_t, S_t\right).\end{aligned}\quad (54)$$

Case 2: For X_2 , $W_1 = -u_x$, $W_2 = -v_x$, $W_3 = -w_x$, and $W_4 = -z_x$, we have

$$\begin{aligned}\omega^x = & \ell - u_x \left[6P(\sum u^2) + P_{xx}\right] - u_{xx}(P_x) - u_{xxx}P - v_x \left[6Q(\sum u^2) + Q_{xx}\right] - v_{xx}(Q_x) - u_{xxx}Q \\ & - w_x \left[6R(\sum u^2) + R_{xx}\right] - w_{xx}(R_x) - w_{xxx}R - z_x \left[6S(\sum u^2) + S_{xx}\right] - z_{xx}(S_x) - z_{xxx}S,\end{aligned}\quad (55)$$

$$\begin{aligned}\omega^t = & -PD_t^{\theta-1}(u_x) + I(-u_x, P_t) - QD_t^{\theta-1}(v_x) + I(-v_x, Q_t) \\ & - RD_t^{\theta-1}(w_x) + I(-w_x, R_t) - SD_t^{\theta-1}(z_x) + I(-z_x, S_t).\end{aligned}\quad (56)$$

Case 3: For X_3 , $W_1 = v$, $W_2 = -u$, $W_3 = 0$, and $W_4 = 0$, we have

$$\begin{aligned}\omega^x = & v \left[6P(\sum u^2) + P_{xx}\right] - v_x P_x + P v_{xx} - u \left[6Q(\sum u^2) + Q_{xx}\right] + u_x Q_x + Q u_{xx}, \\ \omega^t = & PD_t^{\theta-1}(v) + I(v, P_t) - QD_t^{\theta-1}(u) + I(-u, Q_t).\end{aligned}\quad (57)$$

Case 4: For X_4 , $W_1 = w$, $W_2 = 0$, $W_3 = -u$, and $W_4 = 0$, we have

$$\begin{aligned}\omega^x &= w[6P(\sum u^2) + P_{xx}] - w_x P_x + P w_{xx} - u[6R(\sum u^2) + R_{xx}] + u_x R_x + R u_{xx}, \\ \omega^t &= P D_t^{\theta-1}(w) + I(w, P_t) - R D_t^{\theta-1}(u) + I(-u, R_t).\end{aligned}\quad (58)$$

Case 5: For X_5 , $W_1 = z$, $W_2 = 0$, $W_3 = 0$, and $W_4 = -u$, we have

$$\begin{aligned}\omega^x &= z[6P(\sum u^2) + P_{xx}] - z_x P_x + P z_{xx} - u[6S(\sum u^2) + S_{xx}] + u_x S_x + S u_{xx}, \\ \omega^t &= P D_t^{\theta-1}(z) + I(z, P_t) - S D_t^{\theta-1}(u) + I(-u, S_t).\end{aligned}\quad (59)$$

Case 6: For X_6 , $W_1 = 0$, $W_2 = w$, $W_3 = -v$, and $W_4 = 0$, we have

$$\begin{aligned}\omega^x &= w[6Q(\sum u^2) + Q_{xx}] - w_x Q_x + Q w_{xx} - v[6R(\sum u^2) + R_{xx}] + v_x R_x + R v_{xx}, \\ \omega^t &= Q D_t^{\theta-1}(w) + I(w, Q_t) - R D_t^{\theta-1}(v) + I(-v, R_t).\end{aligned}\quad (60)$$

Case 7: For X_7 , $W_1 = 0$, $W_2 = z$, $W_3 = 0$, and $W_4 = -v$, we have

$$\begin{aligned}\omega^x &= z[6Q(\sum u^2) + Q_{xx}] - z_x Q_x + Q z_{xx} - v[6S(\sum u^2) + S_{xx}] + v_x S_x + S v_{xx}, \\ \omega^t &= Q D_t^{\theta-1}(z) + I(z, Q_t) - S D_t^{\theta-1}(v) + I(-v, S_t).\end{aligned}\quad (61)$$

Case 8: For X_8 , $W_1 = 0$, $W_2 = 0$, $W_3 = z$, and $W_4 = -w$, we have

$$\begin{aligned}\omega^x &= z[6R(\sum u^2) + R_{xx}] - w_x R_x + R w_{xx} - w[6S(\sum u^2) + S_{xx}] + w_x S_x + S w_{xx}, \\ \omega^t &= R D_t^{\theta-1}(z) + I(z, R_t) - S D_t^{\theta-1}(w) + I(-w, S_t).\end{aligned}\quad (62)$$

8. Conclusions

The invariance analysis for fractional-order nonlinear coupled Hirota system of PDEs has been done successfully by the application of Lie symmetry reduction. The obtained infinitesimals and generators are used for reduction of the system into nonlinear FODEs via validity of E-K operators in R-L fractional derivative sense. The power series solution and its convergence are discussed by imposition of majorant series and Implicit function theorem. We applied Euler-Lagranges operators to create the adjoint system of the system and conservation analysis with the use of Noether's theorem for stability and consistency of system. This work

may be applicable to fractional fluid flow, propagation of sound, single mode fiber study and other nonlinear evolution problems.

References:

- [1] I. Podlubny: Fractional Differential Equations: An introduction to fractional derivatives, fractional differential equations. *Academic press, San Diego, CA* (1999).
- [2] P.J. Olver: Application of lie group symmetries to differential equations. *Graduated Text in Mathematics, Springer-Verlag*; 107 (2002).
- [3] K.B. Oldham, J. Spanial: The Fractional Calculus. *Academic press, New York* (1974).
- [4] H.M. Srivastava, R.K. Raina, X.J. Yang: Special Functions in Fractional Calculus and Related Fractional Differ-integral Equations. *World Scientific Publishing Company, Singapore, New Jersey, London and Hong Kong* (2014).
- [5] L. Debnath: Recent applications of fractional calculus to science and engineering. *International Journal of Mathematics and Mathematical sciences*; 54(2003) 3413-3442.
- [6] T. Bakkyaraj, R. Sahdevan: Group formalism of lie transformations to time fractional partial differential equations. *Parmana-Journal of Physics*; 85 (2015) 849-860.
- [7] A.M. Wazwaz: Exact solutions for the ZKMEW equation by using the tanh and sine-cosine methods. *International Journal of Computer Mathematics*; 82 (6) (2005) 699-707.
- [8] M.S. Mohamed, G.A. Gapreel: Reduced differential transform method for nonlinear integral member of Kadomtsev-Petviashvili hierarchy differential equations. *Journal of Egyptian Mathematical Society*; 25(2017)1-7.
- [9] H. Gandhi, A. Tomar, D. Singh: A predicted mathematical cancer tumor growth model of brain and its analytical solution by reduced differential transform method. *AISC, Springer*; 1169(2020) 203-213.
- [10] G.W. Bluman, S. Kumei: Use of group analysis in solving overdetermined systems of ordinary differential equations. *Journal of Mathematical Analysis and Applications*; 138(1) (1989) 95-105.
- [11] R. Arora, A. Tomar, V.P. Singh: Similarity Solutions for Strong Shocks in a Non-Ideal Gas. *Mathematical Modelling and Analysis*; 17(3) (2012) 351-365.
- [12] A. Biswas, M. Song, H. Triki, A.H. Kara, B.S. Ahmad, A. Strong, A. Hama: Solitons, Shock waves, Conservation laws and bifurcation analysis of Boussinesq equation with power law non linearity and dual dispersion. *Appl. Math. Inf. Sci.*; 3 (2014) 949-957.
- [13] A. Biswas, C.M. Khalique: Optical Quasi-solitons by lie symmetry analysis. *Jour. King Saud University-Sci.*; 24 (2012) 271-276.
- [14] A. Bansal, A.H. Kara, A. Biswas, S.P. Moshokoa, M. Belic: Optical soliton perturbation, group invariants and conservation laws of perturbed Fokas-Lenells equation. *Chaos Soliton and Fractals*; 114, (2018) 275-280.
- [15] O.S. Iyiola and F.D. Zaman: A fractional diffusion equation model for cancer tumor. *AIP Adv.*; 4 (2014) 107-121.

- [16] A. Iomin: Super diffusion of cancer on a comb structure. *Journal of Physics, conference series*; 7 (2005) 57-67.
- [17] A.M. Wazwaz: The variational iteration method for solving linear and nonlinear systems of PDEs. *Journal Computer Mathematics and Applications*; 54(2007) 895-902.
- [18] H. Jafari, M. Nazarib, D. Baleanuc, C.M. Khalique: A new approach for solving a system of fractional partial differential equations. *Journal Computers and Mathematics with Applications*; 66 (2013) 838–843.
- [19] G. Wang, T. Xu: Symmetry properties and explicit solutions of the nonlinear time fractional kdv equation. *Boundary value Problem*; (2013) 232-245.
- [20] G. Wang, M.S. Hashemi: Lie symmetry analysis and soliton solutions of time fractional K(m,n) equation. *Pramana-Journal of Physic*; 88(1) (2017) 88-95.
- [21] K. Singla, R.K. Gupta: On invariant analysis of some time fractional nonlinear systems of partial differential equations. *Journal of Mathematical Physics*; 57 (2016) 101504.
- [22] T.M. Garrido, M.S. Bruzon: Lie point symmetries and travelling wave solutions for the generalized drinfeld sokolov system. *Journal of Computational and Theoretical Transport*; 45 (2016) 290-298.
- [23] Y. Zhang: Lie symmetry analysis to general time fractional Korteweg-de-Varies equation. *Fractional Differential Calculus*; 5(2015) 125-135.
- [24] I.N. Sneddon: The use in mathematical physics of Erdelyi-Kober operators and some of their generalizations. Lecture notes, *Springer New York*; 457 (1975) 37-79.
- [25] Q. Huang, R. Zhdanov: The efficiency of lie group approach analysis of Harry- dym equation with Riemann-Liouville derivative. *Physica A*; 209 (2014) 110-118.
- [26] H. Liu, J. Li, L. Liu: Lie symmetry analysis, optimal system and exact solution to fifth order KdV types of equations. *Jour. Math. Anal. Appl.*; 368 (2010) 551-558.
- [27] G. Cicogna, G. Gaeta: Noether theorem for μ -symmetries. *J. Phys. A*; 40 (2007), 11899-11921.
- [28] O.S. Iyiola, O.G. Olyinka: Analytical solutions of time fractional models for homogeneous Gardner equation and nonhomogeneous differential equations. *Ain Shams Engg. Journal*; (2014) 2090-4479.
- [29] A. Chauhan, R. Arora: Time fractional Kupershmidt equation: symmetry analysis and explicit series solution with convergence analysis. *Communications in Mathematics*; 27 (2) (2019), 171-185.
- [30] K. Singla, R.K. Gupta: Conservation laws for certain time fractional nonlinear systems of partial differential equations. *Commun. Nonlinear Sci. Numer. Simul.*; 53 (2017) 10-21.
- [31] E. Noether: Invariant variation problems. *Transp. Theory Stat. Phys.*; 1 (1971) 186-207.
- [32] S.Y Lukashchuk: Conservation laws for time fractional sub diffusion and diffusion-wave equations. *Nonlinear Dynamics*; 80 (2015) 791-802.
- [33] N.H. Ibragimov: A new conservation theorem. *Journal Math. Anal. Appl.*; 333 (2007) 311-328.

- [34] C. Chen, Y.L. Jiang and X.T. Wang: Lie symmetry analysis of time fractional generalized KdV equations with variable coefficients. *Symmetry*; 11 (2019) 1281.
- [35] R. Arora, A. Tomar, V.P. Singh: Numerical simulation of Ito coupled system by homotopy analysis method. *Advanced Science, Engineering and Medicine*; 4 (6) (2012), 522-529.
- [36] S. Sahoo, S. Ray: Invariant analysis with Conservation laws for the time fractional Drinfeld-Sokolov-Satsuma-Hirotaequations. *Choas, Solitons and Fractals*; 104 (2017) 725-733.
- [37] K. Adem, M. Khalique: On the solutions and conservation laws of the coupled Drinfeld-Sokolov-Satsuma-Hirota system. *Boundary value problems*; 1 (2014) 248.
- [38] E. Saberi, S.R. Hejazi: Lie symmetry analysis, conservation laws and exact solutions of time fractional generalized Hirota-Satsoma coupled KdV system. *Physica A*; 492 (2017) 296-307.
- [39] R.M. Jena, S. Chakeraverty, D. Baleanu: Solitary wave solution for a generalized Hirota-Satsuma coupled KdV and mKdV equations. *Alexandria Engg. Journal*; 59 (2020) 2877-2889.
- [40] H. Gandhi, D. Singh, A. Tomar: Explicit solution to general fourth order time fractional KdV equation by Lie symmetry analysis. *AIP*; 2253 (2020) Art id020012.
- [41] H. Gandhi, A. Tomar, D. Singh: Lie symmetry analysis to general fifth order time fractional Korteweg-de-Vriesequation and Its explicit solution. *AISC, Springer*; 1169 (2020) 189-201.
- [42] A. Fernandez, D. Baleanu, H.M. Srivastava: Series representations for fractional-calculus operators Involving Generalised Mittag-Leffler Functions, *Commun. in Nonlin. Sci. and Nume. Simul.*; 67 (2019) 517-527.
- [43] E. Guariglia, C. Cattani, H.M. Srivastava: Fractional derivative of the Riemann Zeta function. *Fractional Dynamics*; (2015) 357-368.
- [44] H.M. Srivastava, V.P. Dubey, R. Kumar, J. Singh, D. Kumar, D.Baleanu:An efficient computational approach for a fractional-order Biological population model with carrying capacity. *Chaos, Solitons & Fractals*; 138,109880 (2020)1-13.
- [45] Z.Y. Zhang, J. Zheng, L.L. Guo, H.F. Wu: Lie symmetries and conservation laws of the Fokker-Planck equation with power diffusion. *Mathematical Methods in the Applied Sciences* (2020);<https://doi.org/10.1002/mma.6582>.
- [46] M.S. Bruzón, E.Recio, T.M. Garrido, A.P. Márquez, R. de la Rosa: On the similarity solutions and conservation laws of the Cooper-Shepard-Sodano equation. *Mathematical Methods in the Applied Sciences*.41 (17), 7325-7332.
- [47] S. Gulsen, S.W. Yao, M. Inc: Lie Symmetry Analysis, Conservation Laws, Power Series Solutions and Conservation Analysis of Time Fractional Generalized Drinfeld-Sokolov Systems. *Symmetry-Basel*;13(5), art. n. 874, (2021).
- [48] S.S. Ray, S. Giri: New soliton solution of time fractional Drinfeld-Sokolov-Satsuma-Hirota system in dispersive water waves. *Mathematical Methods in the Applied Sciences* (2021); <https://doi.org/10.1002/mma.7691>.
- [49] H. Yang, W. Liu, Y. Zaho: Lie symmetry reductions and exact solutions to a generalized two-component Hunter-Saxton system. *AIMS Mathematics* 6(2), (2021) 1087-1100.

- [50] H. Gandhi, A. Tomar, D. Singh: Conservation laws and exact series solution of fractional order Hirota-Satsoma coupled Korteweg-de-Vries system by symmetry analysis. *Mathematical Methods in the Applied Sciences* (2021); <https://doi.org/10.1002/mma.7701>.
- [51] V.P. Dubey, J. Singh, A.M. Alshehri, S. Dubey, D. Kumar: A comparative analysis of schemes for solving local fractional Laplace equations. *Mathematical Methods in the Applied Sciences* (2021); <https://doi.org/10.1002/mma.7642>.
- [52] Y. Chatibi, E.E. Kinani, A. Ouhadan: On the discrete symmetry analysis of some classical and fractional differential equations. *Mathematical Methods in the Applied Sciences*, 44 (4) (2020) 2868-2878.
- [53] A. Chauhan, K. Sharma, R. Arora: Lie symmetry analysis, optimal system, and generalized group invariant solutions of the (2+1)-dimensional Date-Jimbo-Kashiwara-Miwa equation. *Mathematical Methods in the Applied Sciences*; 43 (2020) 8823-8840.