

Clp ATPases differentially affect natural competence development in *Streptococcus mutans*

Satya Deo Pandey and Indranil Biswas #

Department of Microbiology, Molecular Genetics and Immunology, University of Kansas Medical

Center, Kansas City, KS, USA¹

Address correspondence to ibiswas@kumc.edu

ABSTRACT

In naturally competent bacteria, DNA transformation through horizontal gene transfer is an evolutionary mechanism to receive extracellular DNA. Bacteria need to maintain a state of competence to accept foreign DNA, and this is an energy-driven phenomenon that is tightly controlled. In *Streptococcus*, competence development is a complex process that is not fully understood. In this study, we used *Streptococcus mutans*, an oral bacterium, to determine how cell density affects competence development. We found that in *S. mutans* the transformation efficiency is maximum when the transforming DNA was added at low cell density and incubated for 2.5 h before selecting for transformants. We also found that *S. mutans* cells remain competent until the mid-logarithmic phase, after which the competence decreases drastically. Surprisingly, we observed that individual components of Clp proteolytic complexes differentially regulate competence. If the transformation is carried out at the early growth phase, both ClpP protease and ClpX ATPase are needed for competence. In contrast, we found that both ClpC and ClpE negatively affect competence. We also found that if the transformation is carried out at the mid-logarithmic growth phase ClpX is still required for competence, but ClpP negatively affects competence. While the exact reason for this differential effect of ClpP and ClpX on transformation is currently unknown, we found that both ClpC and ClpE have a negative effect on transformation, which was not reported before.

INTRODUCTION

Diverse phyla of both Gram-positive and Gram-negative bacteria develop a special physiological state that allows them to uptake DNA from the environment. This state, which is commonly known as natural competence, is generally transient and develops at different stages of growth depending on the bacterial species. Natural competence development has been thoroughly studied in a few Gram-positive and Gram-negative bacterial species (2-5). In *Bacillus subtilis*, the natural competence is developed when the cells enter the stationary growth phase and the cells retain competence for a short time (6). On the other hand, in *Streptococcus pneumoniae*, competence development occurs in a cell density-dependent manner, which occurs in early to mid-log growth phases (7-9). In some organisms such as *Neisseria gonorrhoeae* and *Helicobacter pylori*, the cells are constitutively competent throughout the growth stages (3, 10-12). It is thought that nutritional status and environmental stresses can modulate competence development (13, 14). Furthermore, induction of competence also depends on certain biopolymers such as chitin, as is the case for *Vibrio cholerae* (5, 15-17).

Streptococci are a diverse group of organisms that are represented by over 100 different species (18, 19). Among the streptococci, a handful of species have been characterized for natural competence development (4, 8). Our knowledge of natural competence development in streptococci mostly comes from studies on *S. pneumoniae* (20, 21). In this bacterium, competence is induced by a short 17-residue unmodified peptide pheromone called a competence-stimulating peptide (CSP). This CSP is secreted by a dedicated ABC transporter that cleaves off a leader peptide during transport from the 45-residue pre-CSP encoded by the *comC* gene. Similarly, in *S. mutans*, 21-residue pre-CSP (CSP-21) peptide undergoes post export processing at the C-terminal end that creates a relatively more potent 18-residue peptide (CSP-18) that works at a much lower concentration than CSP-21 (1). When the extracellular concentration of CSP reaches a critical concentration, a membrane-bound sensor kinase, ComD, senses the signal and is activated by autophosphorylation (22-24). The phosphorylated ComD then transfers the phosphate group to its cognate cytoplasmic response regulator ComE for activation. ComD, ComE and ComC are encoded by the *comCDE* operon (22). The activated ComE then induces expression of a competence-specific master sigma factor, *comX*, and the *comCDE* operon (25). Once ComX is made in the cell, it induces expression of so-called late competence genes that are needed for DNA processing, uptake, and recombination (25-27). Typically, *S. pneumoniae* cultures develop competence in the mid-logarithmic phase and last nearly 30 minutes (28). Exogenous addition of CSP leads to induction of *comX* transcription within 5 min and the transcript disappears by 15 min indicating why competence is

short-lived (28). Furthermore, the ComX protein is also degraded rapidly by the ClpC/P and ClpE/P proteolytic complexes that finally cease the competence (29, 30).

In streptococci, ComX is the master regulator of competence development since it directly controls the expression of the late competence genes (31). However, the expression of *comX* is not always dependent on the ComD/E two-component system (TCS). In certain streptococci, including *Streptococcus thermophilus*, a second signal transduction system called ComR/S is also involved in *comX* activation (32). In this case, *comS* encode a short peptide that is secreted outside the cell as an 11-residue long pheromone called ComX-inducing peptide (XIP) (33, 34). When the XIP reaches a certain density, it is imported back into the cell through OppA, an oligo-permease. XIP then interacts with the ComR transcriptional regulator and stimulates binding the ComR-XIP complex to the promoter of *comX* and *comS* and induces transcription. This ComR/S dependent competence induction is only seen in chemically defined media (CDM) that is devoid of oligopeptides, and in this medium, *S. thermophilus* becomes spontaneously competent (32). The induction of early *com* genes in *S. thermophilus* is similar to *S. pneumoniae* and occurs in the early- to mid-log phase (at low cell density). However, in contrast to *S. pneumoniae* and *S. thermophilus*, in *Streptococcus infantarius*, competence develops at the late log phase (high cell density) and lasts for a short time (35). *S. infantarius* competence development is dependent on the ComR/S pathway and not the ComD/E. Despite the presence of all the genes necessary for natural competence development, some streptococci never develop competence (e.g. *S. gallolyticus*), while some others, such as *S. pyogenes*, develop competence only under specific conditions, such as in biofilm (36, 37).

Streptococcus mutans is an oral pathogen associated with dental caries formation and often causes endocarditis. This organism develops competence at the late log phase that persists several hours after induction (38). Interestingly, in *S. mutans* both ComD/E and ComR/S sensor systems modulate the expression of *comX* depending on the growth media (39-41). In nutrient-rich media, ComD/E appears to play a major role, and competence is induced when exogenous CSP is added. Deletion of the ComDE pathway does not abolish competence but transformability is reduced by several orders of magnitude. On the other hand, the ComR/S system is necessary for competence development in CDM. Deletion of ComR also leads to complete cessation of competence development irrespective of the growth media. In *S. mutans*, the ClpC/P complex was shown to be negatively involved in competence by altering the ComX protein concentration in the cell. However, the effect of ClpC/P was only observed in nutrient-rich medium and not in CDM (42, 43). *S. mutans* regulates genetic competence through multiple layers

of control. Working on *S. mutans*, Son and colleagues (44) show that posttranslational control of ComX by MecA/ClpCP is active under both XIP and CSP stimulation. The MecA/ClpCP blocks adventitious entry into competence by sequestering or intercepting low levels of ComX but permits the competence only when ComX levels exceed a threshold. However, cell-to-cell heterogeneity in MecA levels creates variability in that threshold. Therefore, MecA/ClpCP provides a stochastic switch, located downstream of the already noisy *comX*. In *S. mutans*, a third signal transduction system encoded by HdrRM appears to modulate competence in a cell density-dependent manner (45). The exact mechanism by which HdrRM exerts its effect is currently unknown.

The main goal of this study was to determine the optimum environmental conditions to obtain maximum transformation efficiency under nutritionally rich conditions. Our results indicate that maximum transformation occurs near mid-exponential growth phases and the cells remain competent for a long time. We found that the ClpP protease drastically affects competence in a growth phase-dependent manner. In the early stages of growth, both ClpX ATPase and ClpP positively affect competence, while at a later growth stage, ClpP negatively affects competence while ClpX has a negligible effect.

MATERIALS AND METHODS

Strains, media, and growth conditions

S. mutans UA159 and its isogenic mutants $\Delta clpX$, $\Delta clpE$, $\Delta clpP$, and $\Delta clpX$ (57), as well as GS-5, were routinely grown in Bacto Brain-Heart Infusion (BHI; Becton Dickinson) at 37°C under microaerophilic conditions (candle jars), in the presence or absence of antibiotics, such as erythromycin (Em; 1 or 10 µg/ml). DNase I (Sigma) was added when needed as described in the text.

Optimization of transformation assay for natural competence

Strain UA159 was streaked on a BHI-agar plate from glycerol stock (- 80 °C). The next day, a single colony was inoculated into BHI broth and allowed to grow overnight. The culture was diluted to 1:20 in BHI media containing 5% (v/v) horse serum (heat-inactivated; Sigma) and allowed to grow up to OD₆₀₀ 0.15. Plasmid DNA (pIB184Em; (78)), 1 µg/ml and competence stimulating peptide 18 (CSP18: SGSLSTFFRLFNRSFTQA), 200 nM were added at this point (time, 0) and divided in 4 equal volumes (group A - D). In group A, DNase I (10 U) was added at time 0, and an aliquot was plated at different time points. In group B, culture was allowed to grow for 1 hr and then DNase I (10 U) treatment was done for 15 min before plating. In group C, after 1hr culture was further divided equally into 5 vials.

DNase I was added in each vial separately at different time points for 15 min before plating. In group D, DNase I was not added (Fig. 1A). Plating from each group was done at 1, 1.5, 2, 2.5, and 3 h, and colonies were counted after 96 h. The transformation efficiency was calculated as a percentage by dividing the number of transformants by the total number of colony-forming units (CFU). To verify that the transformants are real and not spontaneous mutants, one colony each with and without antibiotic plates was grown in BHI-media and genomic DNA isolated as described before (78). Using isolated genomic DNA as a template and gene-specific primers in PCR, the purity of the transformants was confirmed. The experiment was done in three replicates and values were presented as mean \pm SD. The statistical significance was calculated using a t-test and the p-values <0.05 were considered significant.

Determination of transformation efficiency at various OD₆₀₀ point

The primary inoculum was prepared as above and cultures were grown in the presence of horse serum to OD₆₀₀ 0.2. At this OD, CSP18 was added and cultures were divided equally into 5 vials. The DNA was added to each vial separately when the OD reached the desired density values (0.2, 0.35, 0.5, 0.65, and 0.8). A parallel set of cultures were also grown to determine the OD of the transforming cultures. Cultures were incubated for one more hour before plating. Transformation efficiency was calculated as described earlier. The experiment was done in triplicate and the values were present as mean \pm SD. The statistical significance was calculated using a t-test and the p-values <0.05 were considered significant.

RESULTS

Early-log transformation and late-log plating yield higher transformation efficiency

To better understand the optimum duration of incubation for DNA uptake by *S. mutans*, we used early-log (OD₆₀₀=0.15) grown culture. Transforming DNA (plasmid) and CSP18 were added at the same time. Our overall experimental strategy is shown in Fig. 1A, where we equally divided culture into four groups, A to D. In group A, we added DNase I immediately (time, 0), which served as control. As expected, we obtained no transformants on the antibiotic selection plate even after prolonged incubation (3 h). The data suggest that DNase I is highly effective in digesting DNA in the BHI medium. The absence of DNA was later confirmed by PCR as well (data not shown). In group B, we added DNase I one-hour post-addition of DNA and CSP18 followed by further incubation for different periods of up to three hours. The idea was to assay the multiplicity of the transformed cells in the first hour of incubation. We observed a 5-fold increase in transformants at 2.5 h when compared to 1 h (TE; 50.5×10^{-5} vs 10×10^{-5}) before it started to decline. We believe this 5-fold increase in the transformants is due to the

multiplication of the transformed cells in the first hour of incubation. In group C, we added DNase I at five different time points (five separate aliquots) at the interval of 30 min. This allowed the cells to adsorb and uptake the DNA for different lengths of time. As shown in Fig 1B, we observed a dramatic increase in transformation efficiency (TE) as the incubation time progressed. After 2.5 h incubation, we obtained a nearly 218- fold increase in transformation compared with one-hour incubation (TE; 2181×10^{-5} vs 10.2×10^{-5}). This 218-fold increase in group C compared to just 5- fold in group B at 2.5 h suggests that length of incubation time plays an important role in overall transformation. In group D, we did not add DNase I and as expected, transformation efficiency was the highest; nearly 488-fold observed at 2.5 h when compared to the first hour (TE; 2.73×10^{-2} vs 5.6×10^{-5}) in group D.

When we compared the transformation efficiency between groups C and D, we observed a higher efficiency in group D than in group C at the longer incubation times such as 2, 2.5, and 3 h. However, surprisingly at the early growth stages (1 and 1.5 h), we noted an increase of 2- and 1.4- fold in the efficiency in group C (Fig. 1B). Since this result was consistent we hypothesize that sheared plasmid in group C could have transformed quicker than the full-length plasmid in D and circularized inside the host. To test the hypothesis, we partially treated the transforming plasmid DNA with 0.1U of DNase I for 1 min or linearized the plasmid by restriction digestion with *XhoI* and used it for transformation. We observed no transformant on the selective plate when partially sheared or linear plasmid was used for transformation (data not shown). We also noticed that there was a more viable number of colonies in group A at early growth stages compared to other groups (nearly 2-fold). The data suggest that the addition of DNase I in growth media (BHI) somehow enhanced the bacterial growth that could have resulted in higher TE in the early growth stage in group C. Overall, we observed that maximum transformation is obtained when the transformation is carried out at OD₆₀₀ 0.15. and incubated for 2.5 h.

Low cell density transformation yields higher transformation efficiency

To understand the role of culture density on transformation, we added DNA and CSP18 at different OD₆₀₀ points (0.15, 0.2, 0.35, 0.4, and 0.5) and incubated them further for 2.5 h before plating. The reason was to induce competence at the time of DNA addition and not before that. As shown in Fig. 2, the maximum transformation was obtained at the earliest time point (OD₆₀₀= 0.15). The transformation was dramatically decreased as the culture density was increased. We obtained no transformant when DNA was added at the mid-log phase (OD₆₀₀ = 0.5). To further confirm these results in another strain of *S. mutans*, we used GS5. We found that overall transformation efficiency was nearly 180-fold less in GS5. However, we also observed that transformation was highest at the earliest time point (OD₆₀₀= 0.15)

and then gradually decreased as the cell density increased, a pattern similar to what was observed with UA159. Taken together, the results suggest that the competence development in UA159 and GS5 starts at very low cell density ($OD_{600} = 0.15$) and continues throughout the growth stages, though with a gradually lower rate, up to the mid-log phase ($OD_{600} = 0.5$) before reducing abruptly. And therefore, the optimal time of incubation, 2.5 h, was necessary to attain overall a maximum TE.

The exact mechanism by which plasmid DNA is taken up by the competent cells is poorly understood. It is thought that at least two copies of plasmid DNA need to be simultaneously taken up by the cell to re-circularize inside the cell to generate a complete functional plasmid molecule (46-48). Thus, the transformation conditions described above may be specifically suitable for plasmid DNA. To rule out the possibility, we transformed a linear DNA that contains an *ermA* gene (same resistance gene carried on the plasmid) that is flanked by ~500-bp homology to the SMU.198 locus (49) and compared it with the plasmid DNA transformation. We added DNA at $OD_{600} 0.15$ and incubated for 1- and 2.5 h. As shown in Fig. 3, we obtained maximum transformation after 2.5 h incubation. To our surprise, we also found that the transformation efficiency of plasmid DNA was ~1.5-fold more as compared to the linear DNA at both incubation time points (Fig. 3).

The *clpP* deletion mutant showed higher TE than WT and the *clpX* deletion mutant when transformed at a higher cell density

Transformation is dependent on the activity of Clp proteolytic complexes in streptococci and other bacteria (32, 42, 44, 50). Others and we have previously shown that ClpP affects transformation in *S. mutans* (51, 52). ClpP forms an active proteolytic complex by association with an ATPase and in *S. mutans*, the major Clp proteolytic complex is ClpXP (53, 54). To study the role of ClpP and ClpX in transformation, we used isogenic $\Delta clpX$ and $\Delta clpP$ mutants of the UA159 strain (55, 56). Both these mutants are clean-deletion without any antibiotic resistance markers (57). We then evaluated the transformation efficiency of the strains by adding CSP18 and DNA at low cell density ($OD_{600} = 0.15$), and without CSP18. We observed a reduction in TE of up to 2-3 orders of magnitude in absence of CSP18 in all the three strains during the incubation up to 3 h. We found that both $\Delta clpX$ and $\Delta clpP$ strains displayed a defect in transformation under both conditions (with and without CSP18) when compared to the wild-type UA159. Interestingly, we observed a higher rate of TE in $\Delta clpP$ than $\Delta clpX$ in culture treated with CSP18; however, an opposite trend was observed in culture without CSP18 treatment (Fig. 4A & B). At 2.5 h incubation in culture with CSP18, both the $\Delta clpP$ and $\Delta clpX$ strains

showed about 2.2- and 15-fold reduction in TE, respectively, as compared to the wild type. The fold difference for the $\Delta clpP$ was slightly reduced (~ 2 -fold) after 3 h incubation.

We then wanted to study whether the effect of Clp proteins is culture density-dependent. To confirm, CSP18 was added at low cell-density ($OD_{600} = 0.15$) to induce competence and the transforming DNA was added at different cell densities ($OD_{600} = 0.2, 0.35, 0.5, 0.65$, and 0.8) followed by 1 h incubation at 37°C . Surprisingly, we observed that $\Delta clpP$ strain generated higher transformation as compared to the wild type UA159 at cell densities between $OD_{600} = 0.35$ to 0.8 . For all the strains, transformation efficiency reached the maximum at $OD_{600} = 0.65$. At this cell density, $\Delta clpP$ strain generated nearly 8-fold more transformants compared to the wild type. However, we observed a 2-fold lower transformation in the $\Delta clpP$ mutant than the wild type UA159 at very low cell density ($OD_{600} = 0.2$). Afterward, it gradually increased and reached the maximum at $OD_{600} = 0.65$. In contrast to the $\Delta clpP$, transformation efficiencies of the $\Delta clpX$ mutant were lower for all the cell densities when compared to the wild type (Fig. 4C). To gain insight into the mechanism, we measured the transcript of *comD* and *comX* by RT-PCR (see appendix Fig A1). To our surprise, we found that the expression of *comD* was higher in the *clpP* mutant as compared to the wild-type and the *clpX* mutant. The significance of this observation is not clear at this moment. Taken together, our observation suggests that ClpP and ClpX modulate transformation differently.

Role of various *clp* proteases on TE of UA159

Since ClpP forms a proteolytic complex with two other ATPases, ClpC and ClpE, we wanted to verify whether these ATPases play any role in transformation. We added DNA and CSP18 at low cell density ($OD_{600} = 0.15$) and incubated the culture for 1 and 2.5 h (Fig. 5A). In a parallel set of experiment, DNA was added at high cell density ($OD_{600} = 0.65$) to a CSP18 pre-treated culture (at $OD_{600} = 0.15$) and incubated further for 1 h (Fig. 5B). As expected, when DNA was added at low cell density ($OD_{600}=0.15$) to the CSP18 pre-treated culture, it produced very few transformants after 1 h incubation when compared to the 2.5 h incubation for all the isogenic strains. However, at 2.5 h of incubation, both the $\Delta clpC$ and $\Delta clpE$ strains produced nearly twice as many transformants as compared to the wild type. As expected, the $\Delta clpP$ and $\Delta clpX$ strains produced nearly 3.5- and 35-fold less transformants when compared to the wild type (Fig. 5A). On the other hand, when DNA was added at high cell density, both $\Delta clpC$ and $\Delta clpE$ strains produced nearly 3-fold more transformants as compared to the wild type. Surprisingly, we found that the $\Delta clpP$ strain produced nearly 4-fold more transformants than the wild type. The transformation

efficiency remained low for the $\Delta clpX$ strain (4-fold lower than the wild type) (Fig. 5B). Taken together, our data indicate that the effect of ClpP protease on transformation is cell density-dependent and different Clp ATPases have different effects on transformation.

DISCUSSION

In this study, we attempted to identify the optimum conditions for natural transformation in *S. mutans* with respect to growth in the rich medium. During this work, we have made several observations that suggest that competence development in *S. mutans* UA159 in the rich medium is rather complex. To avoid the unnecessary step of processing the CSP21 peptide by SepM and subsequent accumulation (49), we used the CSP18 peptide at the time of addition of our transforming DNA. We found that competence development in UA159 is greatly dependent on cell density and the maximum transformation efficiency was obtained when DNA was added at very low cell density. We also noted that to obtain maximum transformants, it is necessary to incubate the culture for 2.5 h after the addition of CSP and DNA. If the cultures are plated after 1 h incubation, very few transformants are obtained (Fig. 1B). This is in contrast to the competence development in *S. pneumoniae* where 1 h post-incubation is sufficient (58). Therefore, it appears that either the DNA uptake process or the post-uptake processing, such as recombination, is rather slow in UA159.

We also found that UA159 can be transformed until the mid-log phase if the competence is induced at low cell density ($OD_{600} = 0.15$) by CSP18. We obtained transformants until OD_{600} of 0.8, after which no transformants were obtained. Our result is consistent with an earlier study that also showed no transformants could be obtained after the cell density reaches OD_{600} of 1.1 (59). However, the reduction of competence is very drastic during the growth phase. We noticed that for the UA159 strain, a change in OD_{600} from 0.2 to 0.35 resulted in nearly a 25-fold reduction in transformation; while for the GS-5 strain, the reduction was about 3-fold. Although the exact reason for this reduction is presently unknown, it is possible that the addition of CSP18 at only low cell density was able to induce the late competence genes necessary for the expression of DNA uptake machinery. Nevertheless, this particular assay does not provide any insight into how long the cells can uptake DNA once the competence is induced.

Generally, transformation of linear DNA with homology to the chromosome is much more efficient than transformation with plasmid DNA (60, 61). This is because during transformation double-stranded

circular plasmid DNA is converted to single-stranded linear DNA during uptake (46-48). This linear DNA is later converted to circular DNA with the help of another transforming molecule. Thus, it is assumed that at least two copies of the plasmid need to be in the recipient cell to generate a complete plasmid molecule after transformation. In contrast, linear DNA with homology to chromosome needs to only integrate into the chromosome locus using homologous recombination. Thus, it was surprising to observe that the transformation efficiency of linear DNA was less than that of plasmid DNA (Fig. 3). While the exact reason for this difference is not clearly understood, we speculate that since the linear DNA needs to integrate into the chromosome by double-crossover recombination, the observed difference could be due to a less efficient recombination process rather than due to DNA uptake. Further systematic experimentation with various chromosomal alleles and different plasmid molecules is needed to confirm the difference.

We noticed that the overall transformation efficiency varies greatly between the UA159 and GS5 strains (Fig 2). Strain-dependent variations in natural competence have been reported in *S. pneumoniae* (62, 63) and other non-streptococci including *Vibrio* and *Neisseria* (64, 65). However, most of the competence-related studies in *S. mutans* have been done with UA159 only. The transformation efficiency observed in GS5 was nearly two orders of magnitude lower than the UA159 strain (Fig. 2). However, both the strains show a similar trend in transformation efficiency with respect to cell density. The results are comparable with the *S. pneumoniae* study where the transformation efficiency among the clinical isolates vary nearly four orders of magnitude (63). It is worth mentioning that others and we have found that many *S. mutans* clinical isolates are either poorly or non-transformable. For example, a recent study by Palmer and colleagues found that about 50% of the clinical isolates (7 out of 15) were recalcitrant to natural transformation (66). When we compared the genomes of UA159 and GS-5, we did not find any known competence-related genes that were missing or altered in GS-5. At present, the exact mechanisms responsible for these inter-strain differences in transformation are unknown. However, the inherent quorum signaling may not be responsible since we induced the competence by exogenous addition of CSP18, thereby bypassing the need for the production and secretion of the CSP. Furthermore, the ComD and ComE proteins were also identical in both strains and were biologically active in GS-5 (67).

It is well known that components of the Clp proteolytic complexes are involved in competence development in streptococci (29, 30, 50, 51, 53, 54, 68). Several groups previously demonstrated that both ClpP and ClpX play a positive role in competence development in *S. mutans* when competence is induced at low cell density ($OD_{600} = 0.15$) (51, 54). Our results also indicated that when the transforming

DNA was added to low cell density, both ClpP and ClpX were required for optimum transformation. On the other hand, both ClpC and ClpE displayed an inhibitory effect on transformation at high cell density (Fig. 5A). It has been previously shown in *S. mutans* and other streptococci that ClpC along with the MecA adaptor protein is involved in the degradation of the ComX sigma factor that is responsible for the expression of late competence genes (42-44, 50, 69). However, the role of ClpE in transformation has not been shown previously. It is not clear whether ClpE also plays a role in the degradation of the ComX protein in *S. mutans*. Surprisingly, we found that the effect of ClpP is opposite to ClpC. Since ClpC associates with ClpP protease to form a proteolytic complex, one would expect both ClpC and ClpP to have similar effects on transformation. It is possible that both ClpC and ClpE ATPases associate with some unknown protease to form a productive proteolytic complex that degrades ComX at low cell density; alternatively, the ATPase activity of ClpC and ClpE interferes with the transformation by some unknown mechanisms. Another explanation could be that during low-cell density, some inhibitory peptides, which are substrates of ClpP, are present in the cell and interfere with the transformation. Several such inhibitory peptides have been identified in *S. mutans* and other streptococci (70-74). The situation is somewhat different when the transforming DNA was added at a higher cell density ($OD_{600} = 0.65$). In this case, the inhibitory peptides are either absent or present in very low amounts. At this cell density, ClpC, ClpE, and ClpP all displayed negative effects on transformation perhaps because the ClpC/P and ClpE/P complexes are involved with the degradation of ComX and the other factors necessary for transformation. The role of ClpX in transformation is somewhat of a mystery since it is needed for transformation both at low- and high cell densities. It is noteworthy that, unlike ClpC and ClpE, ClpX expression is not under CtsR regulon and the level of ClpX remains constant throughout the growth (75). Since ClpX forms complexes with ClpP, we speculate that the ClpX/P complex is responsible for the degradation of inhibitory peptides at both low and high cell densities. However, at high cell density, the effect of the inhibitory peptides is not as prominent as at the low cell density. *S. mutans* although does not encode the Lon protease, it does encode several other proteases such as SMU.2153 and PepO (76, 77). ClpE negatively regulates SMU.2153 level in the cell; however, when we deleted SMU.2153, we did not find any noticeable change in the competence development [(76), data not shown]. On the other hand, PepO appears to negatively influence competence development only in defined or semi-defined media (77). Thus, we believe these two proteases do not play a role or are at least not directly involved in competence development under rich growth media. Further in-depth studies are needed to understand the exact role of these Clp ATPases and ClpP in competence development in *S. mutans* UA159.

Author contributions

Satya Pandey: Conceptualization-Equal, Data curation-Equal, Formal analysis-Equal, Investigation-Equal, Methodology-Equal, Validation-Equal, Visualization-Equal, Writing – original draft-Equal, Writing – review & editing-Supporting; **Indranil Biswas:** Conceptualization-Lead, Data Curation-Equal, Formal analysis-Equal, Funding acquisition-Lead, Investigation-Equal, Methodology-Equal, Project administration-Lead, Resources- Lead, Supervision-Lead, Validation-Equal, Visualization-Equal, Writing – original draft-Supporting, Writing – review & editing-Lead

Acknowledgments

This study was supported by an award (GM128241) from NIGMS (NIH) to IB.

Ethics statement

None required

Conflicts of interest

None declared

Data availability statement

All data are provided in this article.

REFERENCES:

1. Petersen FC, Fimland G, Scheie AA. Purification and functional studies of a potent modified quorum-sensing peptide and a two-peptide bacteriocin in *Streptococcus mutans*. *Molecular microbiology*. 2006;61(5):1322-34.
2. Dubnau D. The regulation of genetic competence in *Bacillus subtilis*. *Molecular microbiology*. 1991;5(1):11-8.
3. Dubnau D, Blokesch M. Mechanisms of DNA Uptake by Naturally Competent Bacteria. *Annu Rev Genet*. 2019;53:217-37.
4. Johnston C, Martin B, Fichant G, Polard P, Claverys JP. Bacterial transformation: distribution, shared mechanisms and divergent control. *Nature reviews Microbiology*. 2014;12(3):181-96.
5. Blokesch M. Natural competence for transformation. *Curr Biol*. 2016;26(23):3255.
6. Dubnau D. Genetic competence in *Bacillus subtilis*. *Microbiol Rev*. 1991;55(3):395-424.
7. Tortosa P, Dubnau D. Competence for transformation: a matter of taste. *Curr Opin Microbiol*. 1999;2(6):588-92.
8. Salvadori G, Junges R, Morrison DA, Petersen FC. Competence in *Streptococcus pneumoniae* and Close Commensal Relatives: Mechanisms and Implications. *Front Cell Infect Microbiol*. 2019;9:94.
9. Waters CM, Bassler BL. Quorum sensing: cell-to-cell communication in bacteria. *Annu Rev Cell Dev Biol*. 2005;21:319-46.
10. Hamilton HL, Dillard JP. Natural transformation of *Neisseria gonorrhoeae*: from DNA donation to homologous recombination. *Molecular microbiology*. 2006;59(2):376-85.
11. Aas FE, Wolfgang M, Frye S, Dunham S, Lovold C, Koomey M. Competence for natural transformation in *Neisseria gonorrhoeae*: components of DNA binding and uptake linked to type IV pilus expression. *Molecular microbiology*. 2002;46(3):749-60.
12. Hofreuter D, Odenbreit S, Puls J, Schwan D, Haas R. Genetic competence in *Helicobacter pylori*: mechanisms and biological implications. *Res Microbiol*. 2000;151(6):487-91.

13. Finkel SE, Kolter R. DNA as a nutrient: novel role for bacterial competence gene homologs. *Journal of bacteriology*. 2001;183(21):6288-93.
14. Aminov RI. Horizontal gene exchange in environmental microbiota. *Front Microbiol*. 2011;2:158.
15. Matthey N, Blokesch M. The DNA-Uptake Process of Naturally Competent *Vibrio cholerae*. *Trends Microbiol*. 2016;24(2):98-110.
16. Seitz P, Blokesch M. DNA-uptake machinery of naturally competent *Vibrio cholerae*. *Proceedings of the National Academy of Sciences of the United States of America*. 2013;110(44):17987-92.
17. Stutzmann S, Blokesch M. Comparison of chitin-induced natural transformation in pandemic *Vibrio cholerae* O1 El Tor strains. *Environ Microbiol*. 2020.
18. Facklam R. What happened to the streptococci: overview of taxonomic and nomenclature changes. *Clin Microbiol Rev*. 2002;15(4):613-30.
19. Andam CP, Hanage WP. Mechanisms of genome evolution of *Streptococcus*. *Infect Genet Evol*. 2015;33:334-42.
20. Claverys JP, Dintilhac A, Mortier-Barriere I, Martin B, Alloing G. Regulation of competence for genetic transformation in *Streptococcus pneumoniae*. *J Appl Microbiol*. 1997;83(S1):32S-41S.
21. Straume D, Stamsas GA, Havarstein LS. Natural transformation and genome evolution in *Streptococcus pneumoniae*. *Infect Genet Evol*. 2015;33:371-80.
22. Martin B, Granadel C, Campo N, Henard V, Prudhomme M, Claverys JP. Expression and maintenance of ComD-ComE, the two-component signal-transduction system that controls competence of *Streptococcus pneumoniae*. *Molecular microbiology*. 2010;75(6):1513-28.
23. Johnsborg O, Havarstein LS. Regulation of natural genetic transformation and acquisition of transforming DNA in *Streptococcus pneumoniae*. *FEMS Microbiol Rev*. 2009;33(3):627-42.
24. Martin B, Granadel C, Campo N, Henard V, Prudhomme M, Claverys JP. Expression and maintenance of ComD-ComE, the two-component signal-transduction system that controls competence of *Streptococcus pneumoniae*. *Molecular microbiology*. 2010;75(6):1513-28.
25. Martin B, Soulet AL, Mirouze N, Prudhomme M, Mortier-Barriere I, Granadel C, et al. ComE/ComE~P interplay dictates activation or extinction status of pneumococcal X-state (competence). *Mol Microbiol*. 2013;87(2):394-411.
26. Morrison DA, Lee MS. Regulation of competence for genetic transformation in *Streptococcus pneumoniae*: a link between quorum sensing and DNA processing genes. *Res Microbiol*. 2000;151(6):445-51.
27. Luo P, Li H, Morrison DA. ComX is a unique link between multiple quorum sensing outputs and competence in *Streptococcus pneumoniae*. *Mol Microbiol*. 2003;50(2):623-33.
28. Alloing G, Martin B, Granadel C, Claverys JP. Development of competence in *Streptococcus pneumoniae*: pheromone autoinduction and control of quorum sensing by the oligopeptide permease. *Molecular microbiology*. 1998;29(1):75-83.
29. Chastanet A, Prudhomme M, Claverys JP, Msadek T. Regulation of *Streptococcus pneumoniae* *clp* genes and their role in competence development and stress survival. *J Bacteriol*. 2001;183(24):7295-307.
30. Weng L, Piotrowski A, Morrison DA. Exit from competence for genetic transformation in *Streptococcus pneumoniae* is regulated at multiple levels. *PLoS One*. 2013;8(5):e64197.
31. Claverys JP, Martin B. Competence regulons, genomics and streptococci. *Molecular microbiology*. 1998;29(4):1126-7.
32. Fontaine L, Wahl A, Flechard M, Mignolet J, Hols P. Regulation of competence for natural transformation in streptococci. *Infect Genet Evol*. 2015;33:343-60.
33. Federle MJ, Morrison DA. One if by land, two if by sea: signalling to the ranks with CSP and XIP. *Mol Microbiol*. 2012;86(2):241-5.

34. Mashburn-Warren L, Morrison DA, Federle MJ. A novel double-tryptophan peptide pheromone controls competence in *Streptococcus* spp. via an Rgg regulator. *Molecular microbiology*. 2010;78(3):589-606.
35. Morrison DA, Guedon E, Renault P. Competence for natural genetic transformation in the *Streptococcus bovis* group streptococci *S. infantarius* and *S. macedonicus*. *J Bacteriol*. 2013;195(11):2612-20.
36. Marks LR, Mashburn-Warren L, Federle MJ, Hakansson AP. *Streptococcus pyogenes* biofilm growth in vitro and in vivo and its role in colonization, virulence, and genetic exchange. *J Infect Dis*. 2014;210(1):25-34.
37. Danne C, Guerillot R, Glaser P, Trieu-Cuot P, Dramsi S. Construction of isogenic mutants in *Streptococcus gallolyticus* based on the development of new mobilizable vectors. *Res Microbiol*. 2013;164(10):973-8.
38. Desai K, Mashburn-Warren L, Federle MJ, Morrison DA. Development of competence for genetic transformation of *Streptococcus* mutants in a chemically defined medium. *J Bacteriol*. 2012;194(15):3774-80.
39. Khan R, Rukke HV, Hovik H, Amdal HA, Chen T, Morrison DA, et al. Comprehensive Transcriptome Profiles of *Streptococcus* mutants UA159 Map Core Streptococcal Competence Genes. *mSystems*. 2016;1(2).
40. Underhill SAM, Shields RC, Kaspar JR, Haider M, Burne RA, Hagen SJ. Intracellular Signaling by the comRS System in *Streptococcus* mutants Genetic Competence. *mSphere*. 2018;3(5).
41. Reck M, Tomasch J, Wagner-Dobler I. The Alternative Sigma Factor SigX Controls Bacteriocin Synthesis and Competence, the Two Quorum Sensing Regulated Traits in *Streptococcus* mutants. *PLoS Genet*. 2015;11(7):e1005353.
42. Dong GF, Tian XL, Gomez ZA, Li YH. Regulated proteolysis of the alternative sigma factor SigX in *Streptococcus* mutants: implication in the escape from competence. *BMC microbiology*. 2014;14.
43. Tian XL, Dong G, Liu T, Gomez ZA, Wahl A, Hols P, et al. MecA protein acts as a negative regulator of genetic competence in *Streptococcus* mutants. *J Bacteriol*. 2013;195(22):5196-206.
44. Son M, Kaspar J, Ahn SJ, Burne RA, Hagen SJ. Threshold regulation and stochasticity from the MecA/ClpCP proteolytic system in *Streptococcus* mutants competence. *Molecular microbiology*. 2018;110(6):914-30.
45. Okinaga T, Niu G, Xie Z, Qi F, Merritt J. The *hdrRM* operon of *Streptococcus* mutants encodes a novel regulatory system for coordinated competence development and bacteriocin production. *Journal of bacteriology*. 2010;192(7):1844-52.
46. Saunders CW, Guild WR. Pathway of plasmid transformation in *Pneumococcus*: open circular and linear molecules are active. *Journal of bacteriology*. 1981;146(2):517-26.
47. Lacks S. Uptake of circular deoxyribonucleic acid and mechanism of deoxyribonucleic acid transport in genetic transformation of *Streptococcus pneumoniae*. *Journal of bacteriology*. 1979;138(2):404-9.
48. Barany F, Tomasz A. Genetic transformation of *Streptococcus pneumoniae* by heterologous plasmid deoxyribonucleic acid. *Journal of bacteriology*. 1980;144(2):698-709.
49. Hossain MS, Biswas I. An extracellular protease, SepM, generates functional competence-stimulating peptide in *Streptococcus* mutants UA159. *Journal of bacteriology*. 2012;194(21):5886-96.
50. Wahl A, Servais F, Drucbert AS, Foulon C, Fontaine L, Hols P. Control of natural transformation in salivarius *Streptococci* through specific degradation of sigmaX by the MecA-ClpCP protease complex. *Journal of bacteriology*. 2014;196(15):2807-16.
51. Lemos JA, Burne RA. Regulation and Physiological Significance of ClpC and ClpP in *Streptococcus* mutants. *Journal of bacteriology*. 2002;184(22):6357-66.

52. Chatteraj P, Banerjee A, Biswas S, Biswas I. ClpP of *Streptococcus mutans* differentially regulates expression of genomic islands, mutacin production, and antibiotic tolerance. *J Bacteriol*. 2010;192(5):1312-23.
53. Kajfasz JK, Martinez AR, Rivera-Ramos I, Abranches J, Koo H, Quivey RG, et al. Role of Clp Proteins in Expression of Virulence Properties of *Streptococcus mutans*. *Journal of bacteriology*. 2009;191(7):2060-8.
54. Kajfasz JK, Abranches J, Lemos JA. Transcriptome analysis reveals that ClpXP proteolysis controls key virulence properties of *Streptococcus mutans*. *Microbiology*. 2011;157(Pt 10):2880-90.
55. Jana B, Tao L, Biswas I. Strain-Dependent Recognition of a Unique Degradation Motif by ClpXP in *Streptococcus mutans*. *mSphere*. 2016;1(6).
56. Tao L, Biswas I. Degradation of SsrA-tagged proteins in streptococci. *Microbiology*. 2015;161(Pt 4):884-94.
57. Tao L, Biswas I. Degradation of SsrA-tagged proteins in streptococci. *Microbiol-Sgm*. 2015;161:884-94.
58. Mortier-Barriere I, Campo N, Berge MA, Prudhomme M, Polard P. Natural Genetic Transformation: A Direct Route to Easy Insertion of Chimeric Genes into the Pneumococcal Chromosome. *Methods in molecular biology*. 2019;1968:63-78.
59. Dufour D, Villemin C, Perry JA, Levesque CM. Escape from the competence state in *Streptococcus mutans* is governed by the bacterial population density. *Molecular oral microbiology*. 2016;31(6):501-14.
60. Mannarelli BM, Lacks SA. Ectopic Integration of Chromosomal Genes in *Streptococcus pneumoniae*. *Journal of bacteriology*. 1984;160(3):867-73.
61. Lopez P, Espinosa M, Stassi DL, Lacks SA. Facilitation of Plasmid Transfer in *Streptococcus pneumoniae* by Chromosomal Homology. *Journal of bacteriology*. 1982;150(2):692-701.
62. Li GL, Liang ZW, Wang XT, Yang YH, Shao ZJ, Li MC, et al. Addition of Hypertransformable Pneumococcal Isolates to Natural Transformation for In Vivo Fitness and Virulence. *Infection and immunity*. 2016;84(6):1887-901.
63. Evans BA, Rozen DE. Significant variation in transformation frequency in *Streptococcus pneumoniae*. *Isme Journal*. 2013;7(4):791-9.
64. Duffin PM, Seifert HS. DNA Uptake Sequence-Mediated Enhancement of Transformation in *Neisseria gonorrhoeae* Is Strain Dependent. *Journal of bacteriology*. 2010;192(17):4436-44.
65. Simpson CA, Podicheti R, Rusch DB, Dalia AB, van Kessel JC. Diversity in Natural Transformation Frequencies and Regulation across *Vibrio* Species. *Mbio*. 2019;10(6).
66. Palmer SR, Miller JH, Abranches J, Zeng L, Lefebure T, Richards VP, et al. Phenotypic Heterogeneity of Genomically-Diverse Isolates of *Streptococcus mutans*. *Plos One*. 2013;8(4).
67. Biswas S, Biswas I. SmbFT, a putative ABC transporter complex, confers protection against the lantibiotic Smb in *Streptococci*. *Journal of bacteriology*. 2013;195(24):5592-601.
68. Liu X, Gallay C, Kjos M, Domenech A, Slager J, van Kessel SP, et al. High-throughput CRISPRi phenotyping identifies new essential genes in *Streptococcus pneumoniae*. *Mol Syst Biol*. 2017;13(5):931.
69. Biornstad TJ, Havarstein LS. ClpC acts as a negative regulator of competence in *Streptococcus thermophilus*. *Microbiology (Reading)*. 2011;157(Pt 6):1676-84.
70. Kaspar J, Shields RC, Burne RA. Competence inhibition by the XrpA peptide encoded within the comX gene of *Streptococcus mutans*. *Mol Microbiol*. 2018;109(3):345-64.
71. Shields RC, O'Brien G, Maricic N, Kesterson A, Grace M, Hagen SJ, et al. Genome-Wide Screens Reveal New Gene Products That Influence Genetic Competence in *Streptococcus mutans*. *J Bacteriol*. 2018;200(2).
72. Ahn SJ, Kaspar J, Kim JN, Seaton K, Burne RA. Discovery of novel peptides regulating competence development in *Streptococcus mutans*. *Journal of bacteriology*. 2014;196(21):3735-45.

73. Kaspar J, Ahn SJ, Palmer SR, Choi SC, Stanhope MJ, Burne RA. A unique open reading frame within the *comX* gene of *Streptococcus mutans* regulates genetic competence and oxidative stress tolerance. *Mol Microbiol*. 2015;96(3):463-82.
74. Kim JN, Stanhope MJ, Burne RA. Core-gene-encoded peptide regulating virulence-associated traits in *Streptococcus mutans*. *Journal of bacteriology*. 2013;195(12):2912-20.
75. Gerth U, Kirstein J, Mostertz J, Waldminghaus T, Miethke M, Kock H, et al. Fine-tuning in regulation of Clp protein content in *Bacillus subtilis*. *Journal of bacteriology*. 2004;186(1):179-91.
76. Biswas S, Dhaked HPS, Keightley A, Biswas I. Involvement of ClpE ATPase in Physiology of *Streptococcus mutans*. *Microbiol Spectr*. 2021;9(3):e0163021.
77. Underhill SAM, Shields RC, Burne RA, Hagen SJ. Carbohydrate and PepO control bimodality in competence development by *Streptococcus mutans*. *Molecular microbiology*. 2019;112(5):1388-402.
78. Hossain MS, Biswas I. An Extracellular Protease, SepM, Generates Functional Competence-Stimulating Peptide in *Streptococcus mutans* UA159. *Journal of bacteriology*. 2012;194(21):5886-96.

FIGURE LEGENDS

Fig. 1. Experimental strategy to evaluate transformation efficiency in UA159. (A). Schematic representation of the strategy to evaluate DNA uptake using DNase I (+). At culture $OD_{600} = 0.15$ (considered time zero; $t, 0$), DNA (pIB184Em; 1 μ g/ml) and CSP18 (200nM) were added, and the culture was equally distributed (5 ml each) in four groups (A-D). In group A, DNase I (10U) was added immediately ($t, 0$). In group B, DNase I was added 15 min before plating (P) at 1 h followed by plating at different time points. In group C, culture was further distributed equally in 5 vials (1 ml each), and DNase I was added into each vial at different time points just 15 min before plating. In group D, no DNase I was added. Plating of an aliquot from the four groups was done at 1, 1.5, 2, 2.5, and 3 h of growth at 37 °C. (B). Transformation efficiency plot of four different groups (A to D). The y-axis is represented on the log₁₀ scale and group A was not shown in the graph due to the 'zero' value. The values represent the 'mean \pm SD' of three independent replicates. Statistical significance (*) between groups C and D was calculated using a t-test.

Fig. 2. Comparison of transformation efficiency of two *S. mutans* strains. Transforming DNA (plasmid) and CSP18 were added at indicated culture densities and incubated for 2.5 h before plating on selective plates. The transformation efficiency values of the two strains are shown with the 'mean \pm SD' of three independent replicates. Statistical significance (*) was calculated using one-way ANOVA and values were compared against OD 0.15 in both the panels.

Fig. 3. Transformation of circular and linear DNA molecules. Transforming DNA (plasmid or linear) and CSP18 were added at $OD_{600} = 0.15$ and the culture was further incubated for 1 h or 2.5 h, as indicated. OD_{600} of the culture was measured at the time of plating. The values represent the 'mean \pm SD' of three replicates. Asterisk (*) shows significant p-values (<0.05) calculated using a t-test.

Fig. 4. Effect of Clp proteins on transformation. A). Transformation efficiency of UA159 (WT) and its isogenic *clpX* and *clpP* mutant strains. Transforming DNA (plasmid) and CSP18 were added at culture density $OD_{600} = 0.15$, and the culture was further incubated for indicated time before plating on selective plates. B). A parallel experiment similar to panel A except CSP18 was not added into the culture. C). Transforming DNA (plasmid) was added at various culture densities as indicated (OD_{600} points between 0.2-0.8) followed by 1 h incubation before plating. Asterisk (*) shows a significant p-value (<0.05) in a particular time point calculated using the t-test.

Fig. 5. Effect of Clp ATPases on cell density A). Transformation efficiency at low cell density. Transforming DNA (plasmid) and CSP18 were added at culture density $OD_{600} = 0.15$, and the culture was further incubated for indicated time before plating on selective plates. B). Transforming DNA (plasmid) was added at two culture densities as indicated (OD_{600} points at 0.15 and 0.65) followed by 1 h incubation before plating. CSP18 was added at $OD_{600} = 0.15$ for both the cultures. Asterisk (*) shows significant p-values (<0.05) calculated using T-test. 'ns' indicates 'not significant'.

APPENDIX





Figure A1: Transcriptional expression of *comD* and *comX* using semiquantitative PCR. RNA was isolated at two experimental conditions, without and with CSP18 (-/+), and OD 0.35 and 0.75. After removing gDNA contamination from the RNA samples, cDNA was prepared using company recommendations. Then PCR was done using gene-specific primers. The band intensities of the PCR products were analyzed using ImageJ. The 16S primer was used as an internal positive control standard. Although expression of *comX* in CSP18 treated cells was higher than the untreated ones for all three strains, their levels were close. However, the transcription of *comD* in the *clpP* mutant was dramatically higher than in the wild type (WT) and the *clpX* mutant. It could be a possibility why *clpP* mutant had higher transformation rates when DNA was added at between 0.35 and 0.65 (Fig. 4C).

(A)



(B)




Fig. 2




Fig. 3




FIG. 4




FIG. 5.

