


EuropeAgriDB v1.0: A New Detailed Database to Quantify Nitrogen Budgets in European Agriculture 1961–2019



Rasmus Einarsson, Alberto Sanz-Cobena, Eduardo Aguilera, Gilles Billen,
Josette Garnier, Hans J.M. van Grinsven, & Luis Lassaletta

rasmus.einarsson@chalmers.se




EuropeAgriDB v1.0

- Cropland N budgets 1961–2019 in 26 European countries
- Builds on established methods
 - Lassaletta et al. (2014)
- Higher detail possible in Europe thanks to combination of
 - FAOSTAT
 - Eurostat
 - national statistics
 - many other sources

50 year trends in nitrogen use efficiency of world cropping systems: the relationship between yield and nitrogen input to cropland

Luis Lassaletta¹, Gilles Billen^{1,2}, Bruna Grizzetti³, Juliette Anglade¹ and Josette Garnier^{1,2}




Cropland production

Agricultural land use

Manure N flows

Synthetic N fertilizers



Synthetic N to cropland and permanent grassland


- Some European countries have a rich history of synthetic N inputs to permanent grassland
- But management varies a lot
- This complicates top-down estimates of **cropland** N budgets
- We systematically combined data sources on synthetic N inputs to cropland and grassland

Typical permanent grassland in the Netherlands



Typical permanent grassland in Spain

Result: Estimated synthetic N inputs (kg N/ha/y)




Data availability on crop areas and N harvests

	All crops except fodder*	Fodder* crops
FAOSTAT	Almost 100% since 1961	Very limited
Eurostat	Generally good from around EU accession (varies)	
National statistics	Not investigated	Varies widely

* fodder: temporary grassland, forage legumes, green maize, other plants harvested green, fodder roots

Result: a comprehensive dataset of cropland production


Crop areas



Fodder crops average 1961–2019:

20% of the harvested area, 37% of the harvested N

Examples from the 26 present-day countries in the database



scientific **data**



OPEN

DATA DESCRIPTOR

Crop production and nitrogen use in European cropland and grassland 1961–2019

Rasmus Einarsson^{1,2} [✉], Alberto Sanz-Cobena³ , Eduardo Aguilera³ , Gilles Billen⁴,
Josette Garnier⁴, Hans J. M. van Grinsven⁵  & Luis Lassaletta² [✉]



<https://www.nature.com/articles/s41597-021-01061-z>