

The Eighteen Degree Water (EDW) is an important element of the Northwest Atlantic ocean heat balance and a reflection of the ocean-atmosphere interaction in this region. The analysis of the North Atlantic (NA) climate change based on the World Ocean Atlas [1] and high-resolution Northwest Atlantic Regional Climatology (NWARC) [2] allows us to assess the decadal variability of the EDW (Fig. 1) and its depth that may help us understand localized heat accumulation in the NWA and NA.




Figure 1. Mean annual and seasonal EDW positions at 10m depth for 6 decades (from 1955 to 2017) in the North Atlantic

NCEI's high-resolution NWARC has also shown that the highest rates of heat gain occur in the Sargasso Sea, southeast of the Gulf Stream path in the region occupied by the EDW [3]. The highest resolution analyses on a 0.1°x0.1°-degree latitude-longitude grid indicates that the EDW heaving is patchy and not as homogeneous as coarser resolution mapping [3,4]. The EDW deepening, or “heaving”, on decadal timescales is most likely caused by increasing intensity of Ekman pumping [5].




Figure 2. Mean annual EDW depth for: (a) 1955-1984, b) 1985-2012 periods, (c) EDW depth anomalies (1985-2012 minus 1955-1984)




Figure 3. (a) Decadal EDW volumes (km³) and (b) EDW anomalies as % of mean EDW volume over 1955-2017 period




Figure 4. EDW shapes for coldest (1965-1974, blue) and warmest (2005-2017, dark-yellow) decades showing EDW heaving: a) Northeast view; b) Southwest view. (For better view - see animation)

The depths of EDW for two ~30-year intervals and their differences are shown in Fig. 2. The volume of EDW depends on many factors of which the most important are Ekman pumping, heat fluxes across the surface, and heat advection within the Gulf Stream and the subtropical recirculation gyre. However, heat accumulation in several “pockets” southeast of the Gulf Stream and its extension seems to be most closely connected to EDW

heaving. Fig. 3 illustrates the variations in the EDW volume and its anomalies (in % comparing to the 60-year mean value) for 6 decades from 1955 to 2017. The difference in EDW ‘heaving’ is quite substantial – the volume of EDW has increased by ~790K km³, which is equal to 35 volumes of all Great Lakes combined. Fig. 4 illustrates the EDW depths change from the coldest (1965-1974) to warmest (2005-2017) decade.

Comparing the eddy-permitting EDW heaving inferred from NCEI's NWARC and the ~30-year shift of the curl of wind-stress, and consequently Ekman pumping, confirms strong resemblance of the eddy-permitting and eddy-resolving EDW heaving patterns to two tightly localized pockets of heat accumulation southeast of the Gulf Stream and its extension [5].

References

1. Locarnini, R., *et al.* (2018). World Ocean Atlas 2018, Vol. 1: Temperature. A. Mishonov Technical Ed. NOAA Atlas NESDIS 81. <https://www.nodc.noaa.gov/OC5/woa18/pubwoa18.html>
2. Seidov, D., O. Baranova, T. Boyer, S. Cross, A. Mishonov, and A. Parsons (2016). Northwest Atlantic Regional Ocean Climatology. NOAA Atlas NESDIS 80; doi: 10.7289/V5/ATLAS-NESDIS-80
3. Seidov, D., A. Mishonov, J. Reagan, and R. Parsons (2017). Multidecadal variability and climate shift in the North Atlantic Ocean, *GRL*, 44; doi: [doi:10.1002/2017GL073644](https://doi.org/10.1002/2017GL073644)
4. Seidov, D., A. Mishonov, J. Reagan, O. Baranova, S. Cross, and A. Parsons (2018). Regional climatology of the Northwest Atlantic Ocean—high-resolution mapping of ocean structure and change, *BAMS*; doi: [10.1175/BAMS-D-17-0205.1](https://doi.org/10.1175/BAMS-D-17-0205.1)
5. Seidov, D., A. Mishonov, J. Reagan, and R. Parsons (2018). Eddy-resolving in situ ocean climatologies of temperature and salinity in the Northwest Atlantic Ocean, *JGR*, in review: MS 2018JC014548.