

Valdir F. Novello (vfnovello@gmail.com)^{1*}, Francisco W. Cruz¹, Mathias Vuille², José L. P. S. Campos¹, Nicolás M. Stríkis³, James Apáestegui⁴, Jean S. Moquet⁵, Vitor Azevedo¹, Angela Ampuero¹, Giselle Utida¹, Xianfeng Wang⁶, Gustavo M. Paula-Santos⁷, Plínio Jaqueto⁸, Luiz C. R. Pessenda⁹, Daniel O. Breecker¹⁰, Ivo Karmann¹

1- Institute of Geosciences, University of São Paulo; 2- Department of Atmospheric and Environmental Sciences, University at Albany; 3- Departamento de Geoquímica, Universidade Federal Fluminense; 4- Instituto Geofísico del Perú, Lima; 5- Institut des Sciences de la Terre d'Orléans; 6- Asian School of the Environment and Earth Observatory of Singapore; 7- Institute of Geoscience University of Campinas; 8- Instituto de Astronomia, Geofísica e Ciências Atmosféricas, University of São Paulo; 9- Center for Nuclear Energy in Agriculture (CENA), University of São Paulo; 10- Jackson School of Geosciences, University of Texas.

Introduction and Objectives

Due to the many factors controlling $\delta^{13}\text{C}$ values in stalagmites, complicating their paleoclimatic and paleoenvironmental interpretation, most studies do not present $\delta^{13}\text{C}$ values, but instead focus mainly on $\delta^{18}\text{O}$ values. This is also the case for most cave studies from tropical South America, where many new $\delta^{18}\text{O}$ stalagmite records covering the last millennia were recently published. Here, we review the $\delta^{13}\text{C}$ values in stalagmites, investigating the influence on this proxy of local hydroclimate, altitude, temperature and vegetation types, by employing a new dataset composed of 25 $\delta^{13}\text{C}$ records (13 of them hitherto unpublished) from stalagmites collected at different sites throughout tropical South America (Fig. 1, $\delta^{13}\text{C}_{2k_SA}$ dataset) with aims to revise and characterize the main factors controlling $\delta^{13}\text{C}$ variability in these stalagmites and provide possible paleoclimate and paleoenvironmental reconstructions for the region.




Fig. 2. Relationship between the $\delta^{13}\text{C}$ from the $\delta^{13}\text{C}_{2k_SA}$ dataset with annual precipitation (a) and annual mean temperature (b) of each study site.




Fig. 4. Comparison between the first Principal Component (PC1) derived from the $\delta^{13}\text{C}_{2k_SA}$ dataset (this study) representing the main mode of hydroclimate variability and PC1 based on the $\delta^{18}\text{O}$ from stalagmites from South America (PC1 of $\delta^{18}\text{O}$) representing the main mode of variability of the South American Monsoon System. The uncertainties associated with the PC1s are shown by the colored and gray shading. λ indicates the percentage of explained variance by each PC.




Fig. 1. Map of tropical South America with vegetation types, precipitation (blue isolines - mm/month, derived from the annual mean for the period from 1998 to 2017) and location of the study sites from the $\delta^{13}\text{C}_{2k_SA}$ dataset.




Fig. 3. Maps of South America with the main loadings of the Principal Component Analysis (PCA) and explained total variance. Blue and red dots represent positive and negative loadings, respectively. The magnitude of the loadings is represented by the size of the dots. The larger the dot, the more representative it's loading is of the respective PC.

Conclusion

The predominance of C3 plants above most of the karst systems studied here is responsible for the low $\delta^{13}\text{C}$ values (<6‰) in most of the speleothems, while local hydroclimate associated with PCP process is the main driver behind its variability during the last two millennia (Fig. 2 a). Local temperature and growth rate play only a minor role in shaping the $\delta^{13}\text{C}$ values in the $\delta^{13}\text{C}_{2k_SA}$ dataset (Fig. 2b). The probable reason for this is that most of the speleothems in our database formed under tropical conditions, characterized by a limited temperature range. Using Monte Carlo Principal Component Analysis, we produce an index of the mean hydrologic conditions and its changes over tropical South America for the last two millennia (Fig. 3), which is closely related to monsoon variability for the period prior to 1750 CE (Fig. 4). The recent break-down in the relationship between monsoon and local hydroclimate may have been caused by the increase in temperature, CO_2 , deforestation and fire during the current warm period; however, further studies are required to test this hypothesis.

Related paper:

Novello V. F.; Cruz, F. W.; Vuille, M.; Campos, J. L. P. S.; Stríkis, N. M.; Apáestegui, J.; Moquet, S. J.; Azevedo, V.; Ampuero, A.; Utida, G.; Wang, X.; Paula-Santos, G. M.; Jaqueto, P.; Pessenda, L. C. R.; Breecker, D. O.; Karmann, I. (2021). Investigating $\delta^{13}\text{C}$ values in stalagmites from tropical South America for the last two millennia. *Quaternary Science Reviews*, 255, 106822, <https://doi.org/10.1016/j.quascirev.2021.106822>.