

Assessing and correcting estimated fCO₂ from carbonate chemistry models of the northeastern US

Corbin T Lanker^{1,2} and Jennie E. Rheuban¹

¹Woods Hole Oceanographic Institution, Department of Marine Chemistry & Geochemistry, ²The Ohio State University School of Earth Sciences




Introduction

Characterizing the ocean carbonate system is important in understanding ocean acidification and broader ocean chemistry. Models can help extrapolate carbonate chemistry parameters from existing chemistry data (Fig. 1)

This investigation explores the ability of a Random Forest Regression Model (RFR), a multiple linear regression model (MLR)¹ and bottle samples to predict the fugacity of CO₂ (fCO₂) in the Northeast US nearshore region from modeled or measured total alkalinity (TA) and dissolved inorganic carbon (DIC)

Figure 1. Spatial Distribution of Bottle Samples (Top) and CTD Casts (Bottom). Bottle samples have measures of TA and DIC, CTD cast measurements are used to model TA and DIC where measures aren't available.



Methods

To compare modeled fCO₂, the publicly available Surface Ocean CO₂ Atlas (SOCAT) was used as a source of underway fCO₂ values measured at 5m depth.

fCO₂ was calculated from measured or modeled TA and DIC using CO2SYS (v. 1.1). Surface ocean chemistry was calculated from a mean of values from a depth range. Surface fCO₂ was then compared to SOCAT fCO₂ values by matching SOCAT datapoints within a specified radial distance and collection date range of the modeled and bottle datapoints. Multiple depth, distance and time ranges were used and compared for the best fit. fCO₂ comparisons were analyzed using Root Mean Square Error and models of linear fit.

Results

-Best fit occurred with:

- Surface depth defined as 2-8m
- Spatial Range of 1km
- Temporal Range of 3 days

-RFR and Biogeochemical MLR were biased to overestimate fCO₂ (Fig. 2)

-Calculated fCO₂ from bottle data generated a similar bias as the modeled data (Fig. 2)

-Physical MLR showed no clear relationship with the SOCAT data (Fig. 2)

-Salinity and sea surface temperature were well matched between the SOCAT data and measurements used to model




Figure 2. Direct comparisons of SOCAT fCO₂ (measured in μatm) with modeled/bottle data with linear fits.

Conclusions

Extrapolating carbonate models beyond the parameters they are intended to predict should be done with an abundance of caution.

For the Northeast US, it is suspected that CO2SYS fails to accurately predict fCO₂ due to high contributions of organic alkalinity for which CO2SYS has not accounted. As such, linear models were used to generate corrections for the models and bottle samples to make fCO₂ values extrapolated from DIC and TA closer to reality (Fig. 3)




Figure 3. Spatial distribution of fCO₂ difference (Model-Measured) for model before correction (Top) and after correction (Bottom)

Acknowledgements

The authors would like to express the utmost gratitude for NSF REU OCE-1852460, the WHOI Summer Student Fellowship, Louise P. Cameron, Z. Aleck Wang, Ivan D. Lima, NOAA Climate Program Office and SOCAT for all their support of this project

¹McGarry, K., Siedlecki, S. A., Salisbury, J., & Alin, S. R. (2021). Multiple linear regression models for reconstructing and exploring processes controlling the carbonate system of the northeast US from basic hydrographic data. *Journal of Geophysical Research: Oceans*, 126, e2020JC016480. <https://doi.org/10.1029/2020JC016480>