

# Non-local Eddy-Mean Kinetic Energy Transfers in Submesoscale-Permitting Ensemble Simulations

Quentin Jamet<sup>1\*</sup>, Stephanie Leroux<sup>1,2</sup>, William K. Dewar<sup>1,3</sup>, Thierry Penduff<sup>1</sup>,  
Julien Le Sommer<sup>1</sup>, Jean-Marc Molines<sup>1</sup>, Jonathan Gula<sup>4,5</sup>

<sup>1</sup>Univ. Grenoble Alpes, CNRS, IRD, Grenoble INP, IGE, Grenoble, France

<sup>2</sup>Ocean Next, Grenoble, France

<sup>3</sup>Dept. of EOAS, Florida State University, Tallahassee, FL, USA

<sup>4</sup>Univ Brest, CNRS, Ifremer, IRD, Laboratoire d'Océanographie Physique et Spatiale (LOPS), IUEM,  
Plouzané, France

<sup>5</sup>Institut Universitaire de France (IUF), Paris, France

## Key Points:

- Ensemble-based eddy-mean decomposition of kinetic energy budget supports the view of an ocean turbulence driven by internal dynamics
- Turbulent fluxes of the cross-energy term provide a potentially strong horizontal constraint on eddy-mean flow interactions
- Non-localities are leading order at small scales and should be accounted for in meso-to-submeso scale range parameterizations

---

\*Univ. Grenoble Alpes, CNRS, IRD, Grenoble INP, IGE, Grenoble, France

Corresponding author: Quentin Jamet, [quentin.jamet@univ-grenoble-alpes.fr](mailto:quentin.jamet@univ-grenoble-alpes.fr)

## Abstract

Understanding processes associated with eddy-mean flow interactions helps our interpretation of ocean energetics, and guides the development of parameterizations. Here, we focus on the non-local nature of Kinetic Energy (KE) transfers between mean and turbulent reservoirs. Transfers are interpreted as non-local when the energy extraction from the mean flow does not locally sustain energy production of the turbulent flow, or vice versa. The novelty of our approach is to use ensemble statistics to define the mean and the turbulent flow. Based on KE budget considerations, we first rationalize the eddy-mean separation in the ensemble framework, and discuss the interpretation of a mean flow  $\langle \mathbf{u} \rangle$  driven by the prescribed (surface and boundary) forcing and a turbulent flow  $\mathbf{u}'$  driven by non-linear dynamics sensitive to initial conditions. We then analyze 120-day long, 20-member ensemble simulations of the Western Mediterranean basin run at  $\frac{1}{60}^\circ$  resolution. Our main contribution is to recognize the prominent contribution of the cross energy term  $\langle \mathbf{u}_h \rangle \cdot \mathbf{u}'_h$  to explain non-local energy transfers. This provides a strong constraint on the horizontal organization of eddy-mean flow KE transfers since this term vanishes identically for perturbations  $(\mathbf{u}'_h)$  orthogonal to the mean flow  $(\langle \mathbf{u}_h \rangle)$ . We also highlight the prominent contribution of vertical turbulent fluxes for energy transfers within the surface mixed layer. Analyzing the scale dependence of these non-local energy transfers supports the local approximation usually made in the development of meso-scale, energy-aware parameterizations for non-eddy models, but points out to the necessity of accounting for these non-local effects in the meso-to-submeso scale range.

## Plain Language Summary

The ocean constantly exchanges energy between its mean and its turbulent reservoirs. However, we are still lacking a clear understanding of these eddy-mean flow interactions, which limits our ability to represent them in numerical ocean simulations that require turbulent closures. Here, we focus on the spatial non-locality of these interactions. We analyze for this the kinetic energy exchanges between the ensemble mean and the residual flow during the decorrelation phase of an ensemble of submesoscale-permitting simulations of the Western Mediterranean basin. Our main contribution is to highlight the prominent role played by turbulent fluxes of the cross-energy term  $\langle \mathbf{u}_h \rangle \cdot \mathbf{u}'_h$  in driving non-local kinetic energy transfers. Through these turbulent fluxes, the energy lost by the ensemble mean flow at one location can be transported over tens of kilometers



before to be transferred to the turbulent flow (or vice versa), making the eddy-mean kinetic energy transfers non-local. We then analyze the geographical organization of these turbulent fluxes, and highlight a potentially strong horizontal constraint on eddy-mean flow interactions owing to their particular dynamics. Finally, we quantify the scale dependence of these non-localities, and suggest that their effects should be accounted for in meso-to-submeso scale range parameterizations.

## 1 Introduction

Meso-scale eddies play a crucial role for the energetic balance of the ocean, providing the main pathway toward dissipative scales (Wunsch & Ferrari, 2004). Understanding how these eddies interact with the mean flow thus helps our interpretation of the ocean circulation, and also serves as a basis for the development of robust parameterizations for ocean models. In order to gain insights from the different processes controlling the energetic of these eddies, it is usual and natural to investigate the different terms contributing to the time rate of change of the Eddy Kinetic Energy (EKE) equation (e.g., Webster, 1961, 1965; Dewar & Bane, 1989). From a point of view of parameterization, evaluating the energy levels of meso-scale ‘*eddies*’ is used to constrain numerical eddy dissipation coefficients through mixing length arguments (Cessi, 2008; Eden & Greatbatch, 2008; Mak et al., 2018; Jansen et al., 2019), thus making these coefficients energy-aware. In this context, the ‘*eddies*’ are associated with unresolved, sub-grid scale physics that need to be parameterized. Processes controlling this physics thus need to be represented based on the *mean*, resolved flow. A particularity of eddy-mean kinetic energy transfers lies in the difference in the terms involved in KE budget of the mean and the turbulent flow. That is, changes in the energy of the mean flow are subject to the divergence of an eddy stress tensor correlated with the mean flow, while changes in the energy of the turbulent flow are subject to a turbulent flux up or down the gradient of the mean flow. Equating the eddy-mean interaction term from these two different perspectives is subject to an assumption of locality, where the energy released by the mean flow at one location is assumed to sustain the growth of eddies at that location (or vice versa for energy backscattering processes). However, recent studies based on Lorenz energy cycles at global (Chen et al., 2014, 2016) and regional (Kang & Curchitser, 2015; Capó et al., 2019) scales have shed light on the strong non-locality of these transfers at small scales. Our interest in this study is to further investigate the non-local nature of these

eddy-mean kinetic energy transfers, leveraging the recent developments of kilometric-scale resolution ensemble simulations to separate mean and eddies based on ensemble statistics.

An emerging concern for the development of turbulent parameterizations for ocean models is placed on the non-locality of energy transfers. In early work on energy-aware parameterizations for mesoscale turbulence, Cessi (2008) has proposed an improved Gent-McWilliams (Gent & McWilliams, 1990) formulation in which the eddy buoyancy diffusivity was defined as a function of the averaged sub-grid scale turbulent kinetic energy through mixing length arguments. Although globally integrated estimates of sub-grid scale kinetic energy offer interesting properties (Marshall & Adcroft, 2010), it obviously *only* provides an averaged estimate. Other studies have provided more elaborated formulations to account for the spatial organization of mesoscale eddy diffusivity (Visbeck et al., 1997; Ferreira et al., 2005; Groeskamp et al., 2020), but at the expense of severely complicating the prognostic equation of sub-grid scale turbulent kinetic energy that needs to be solved (Eden & Greatbatch, 2008; Mak et al., 2018; Jansen et al., 2019). In practice, the several processes involved in this prognostic equation are usually parameterized through isotropic dissipative operators, mostly due to the lack of better theories. However, Grooms (2017) has recently shown that, while this approximation is valid for isotropic barotropic turbulence with no mean flow, idealized advection-diffusion models rapidly fail to accurately represent the transport of EKE when a mean flow is added to the problem (through the  $\beta$  effect in his case). A potential reason to explain this is associated with the non-locality of the eddy energy transfers, as for instance identified in a wind-driven, two-layer QG model by Grooms et al. (2013); in this simulation, the energy lost by eddies in the separated jet is primarily balanced by imports of energy from remote regions. Non-local kinetic energy reported by Grooms et al. (2013) are associated with various processes, such as wave radiation, advection, or eddy-mean flow interactions. The latter relates the dynamics behind energy transfers between the mean and the turbulent flow. The leading order contribution of this latter term has been recently reported by Chen et al. (2014), Kang and Curchitser (2015) and Capó et al. (2019) in realistic simulations, and is thus likely to have important implications for the development of future parameterizations.

There are many ways to define 'mean' and 'eddies', the most traditional approach being to use a time averaging. This definition offers several advantages, such as ease in

implementation and natural interpretation when dealing with observations. Eddies so defined are however associated with all signals that vary in time, which makes the attribution of processes somehow ambiguous (for instance to disentangle processes associated with hydrodynamic instabilities from those associated with time varying forcing). Coarse-graining (e.g. Aluie et al., 2018) offers an alternative approach, which is more intuitive in the context of parameterization. Although the time dimension is retained, this approach induces some subjectivity in the definition of length scale cutoff, thus the size of the eddies, as well as complexities in dealing with solid boundaries, isotropy and inhomogeneities of the flow structure.

Here, we choose to leverage ensemble simulations to define the 'mean' flow as that common to all members (i.e. an ensemble mean), and the 'eddies' as the deviation of each member with its ensemble mean. We will argue in the following that this approach offers an unambiguous definition of 'eddies' through KE budget considerations; it allows to robustly separate the flow in a part that is controlled by the prescribed forcing (the 'mean' flow), and a part that is intrinsically driven by non-linear dynamics (the 'eddies'). This strategy also allows the analysis of the spatio-temporal structure of ocean turbulence and its associated flux of energy. An obvious limitation is associated with the computational resources required to produce such a data set. Here, in order to partially account for the potential effects of submesoscale dynamics in eddy-mean flow interactions, we have used the newly generated kilometric-scale resolution ( $\frac{1}{60}^\circ$ ) MEDWEST60 ensemble simulations of Leroux et al. (2021). It is composed of 20 ensemble members subject to small initial conditions uncertainties (usually referred to as *micro* initial conditions ; Stainforth et al., 2007), run for 120-days from the already spun-up oceanic state of eNATL60 simulation (Brodeau et al., 2020), a numerically identical, single simulation run over the whole North Atlantic basin. Analyzing the decorrelation of each ensemble member in this context informs us on the processes controlling the growth of ensemble spread, thus on the spatio-temporal structure of eddy-mean flow interactions.

The paper is organized as follows. In Section 2, we present the MEDWEST60 ensemble simulations, along with the associated mean and eddy kinetic energy budget decomposition. We then discuss the decorrelation of the turbulent flow from initial conditions, and some aspects of the associated kinetic energy budgets in Section 3. In Section 4, we first diagnose the non-local kinetic energy transfers, and then estimate the scale

dependence of these processes with a view toward parameterization. We finally summarize our results and discuss their implications in Section 5.

## 2 Methods

### 2.1 Kinetic Energy Budget of Ensemble Simulations

Our primary interest is to investigate the kinetic energy budget of the MEDWEST60 submesoscale-permitting ensemble simulations, described in Section 2.2, with a focus on energy transfers between the ensemble mean and the turbulent flow. The momentum equations solved by these simulations are the Boussinesq, hydrostatic equations written in flux form:

$$\partial_t u = -\nabla \cdot \mathbf{u}u + fv - \frac{1}{\rho_0} \partial_x p + \mathbf{D}_u, \quad (1a)$$

$$\partial_t v = -\nabla \cdot \mathbf{u}v - fu - \frac{1}{\rho_0} \partial_y p + \mathbf{D}_v, \quad (1b)$$

with  $\mathbf{u} = (u, v, w)$  the three-dimensional velocity field,  $\nabla = (\partial_x, \partial_y, \partial_z)$  the three-dimensional gradient,  $f = 2\Omega \sin(\phi)$  the Coriolis frequency,  $p = \int_z^\eta \rho g dz$  the (hydrostatic and surface) pressure field, and  $\mathbf{D}_u = \partial_z (\mathbf{A} \partial_z u)$  and  $\mathbf{D}_v = \partial_z (\mathbf{A} \partial_z v)$ , the viscous effects including both surface wind forcing and bottom drag as upper and bottom boundary conditions, respectively, as well as interior ocean dissipation of momentum, with  $\mathbf{A}$  the spatiotemporally varying viscous coefficient computed through the TKE turbulent closure scheme<sup>1</sup>. Following standard practices, an equation for the hydrostatic kinetic energy

$$K = \frac{\rho_0}{2} (\mathbf{u}_h \cdot \mathbf{u}_h), \quad (2)$$

with  $\mathbf{u}_h = (u, v)$  the horizontal component of the velocity field, is obtained by multiplying (1a) by  $\rho_0 u$  and (1b) by  $\rho_0 v$ , and summing the resulting equations, such that:

$$\partial_t K = -\nabla \cdot \mathbf{u}K - \mathbf{u}_h \cdot \nabla_h p + \rho_0 \partial_z (\mathbf{A} \partial_z K) - \epsilon, \quad (3)$$

with  $\nabla_h = (\partial_x, \partial_y)$  the horizontal gradient,  $\rho_0 \partial_z (\mathbf{A} \partial_z K)$  the work done by vertical viscous forces, and  $\epsilon = \rho_0 \mathbf{A} \partial_z \mathbf{u}_h \partial_z \mathbf{u}_h$  the vertical dissipation. Adding and subtracting  $-w \partial_z p = wb$  in (3), and using the continuity equation for Boussinesq fluids  $\nabla \cdot \mathbf{u} = 0$ , allows the

---

<sup>1</sup> Note that horizontal viscous effects are implicitly included in the UBS advective scheme as a biharmonic operator (Shchepetkin & McWilliams, 2005). See Appendix for further details on these operators and their implementation in MEDWEST60.

pressure term to be written as the divergence of a flux, and makes explicit the exchange of kinetic energy with potential energy through  $wb$ . It leads to:

$$\partial_t K = -\nabla \cdot \mathbf{u}K - \nabla \cdot \mathbf{u}p - wb + \rho_0 \partial_z (\mathbf{A} \partial_z K) - \epsilon. \quad (4)$$

In our ensemble simulations, the velocity field simulated by each individual ensemble member obeys this KE equation. It is however possible, from ensemble statistics, to decompose this velocity field as that common to all members, and that specific to each member, and analyze their kinetic energy expression.

For this, we consider the Reynolds decomposition

$$x_n = \langle x \rangle + x'_n, \quad (5)$$

where the mean operator

$$\langle x \rangle = \frac{1}{N} \sum_{n=1}^N x_n. \quad (6)$$

represents the ensemble mean, with  $N$  the size of the ensemble. Following this procedure to decompose the zonal and meridional velocities defining the kinetic energy (2) leads to:

$$K = \tilde{K} + K^* + \rho_0 \langle \mathbf{u}_h \rangle \cdot \mathbf{u}'_h, \quad (7)$$

where  $\tilde{K} = \frac{\rho_0}{2} (\langle \mathbf{u}_h \rangle \cdot \langle \mathbf{u}_h \rangle)$  and  $K^* = \frac{\rho_0}{2} (\mathbf{u}'_h \cdot \mathbf{u}'_h)$ . For reasons explained below, we will refer the former quantity ( $\tilde{K}$ ) as the Forced Kinetic Energy (FKE), and the ensemble mean of the latter quantity ( $\langle K^* \rangle$ ) as the Internal Kinetic Energy (IKE). This refers to the kinetic energy of the ensemble mean flow and that of the perturbations, respectively. The notation used here is somehow different from the more classical Mean and Eddy Kinetic Energy (MKE, EKE) terminology used when working with time averages. While these terms are formally the same, the different terminology used here aims at highlighting differences in their interpretation and properties in the context of ensemble simulations. Such differences are further discussed in the following. Finally, we note that the vector form employed here also emphasizes that, in addition to vanishing identically upon averaging, the cross energy term  $\rho_0 \langle \mathbf{u}_h \rangle \cdot \mathbf{u}'_h$  is also zero for turbulent flow orthogonal to the mean flow.

The kinetic energy equation for the mean flow and that for the perturbations are usually derived based on averaged and residual forms of (1a) and (1b). Formally, multiplying the ensemble mean equations  $\langle (1a) \rangle$  and  $\langle (1b) \rangle$  by the ensemble mean zonal and

meridional velocities  $\rho_0 \langle u \rangle$ ,  $\rho_0 \langle v \rangle$ , respectively, and summing the resulting equations, leads to an equation for the Forced Kinetic Energy (FKE) of the form:

$$\partial_t \tilde{K} = -\nabla \cdot \langle \mathbf{u} \rangle \tilde{K} - \rho_0 \langle \mathbf{u}_h \rangle \cdot \nabla \cdot \langle \mathbf{u}' \mathbf{u}'_h \rangle - \nabla \cdot \langle \mathbf{u} \rangle \langle p \rangle - \langle w \rangle \langle b \rangle + \rho_0 \partial_z \left( \langle \mathbf{A} \rangle \partial_z \tilde{K} \right) - \epsilon_{\tilde{K}}, \quad (8)$$

where the underlined term is associated with eddy-mean flow interactions, and the exchange of FKE with forced potential energy is made explicit through the inclusion of  $\langle w \rangle \langle b \rangle$ .

A similar equation is obtained for the Internal Kinetic Energy (IKE) by multiplying the residual equation for the zonal and meridional momentum (1a)' and (1b)' by the zonal and meridional velocity perturbations  $\rho_0 u'$  and  $\rho_0 v'$ , ensemble averaging and then summing the resulting equations. This leads to:

$$\partial_t \langle K^* \rangle = -\nabla \cdot \langle \mathbf{u} K^* \rangle - \rho_0 \langle \mathbf{u}' \mathbf{u}'_h \rangle \cdot \nabla \cdot \langle \mathbf{u}_h \rangle - \nabla \cdot \langle \mathbf{u}' p' \rangle - \langle w' b' \rangle + \rho_0 \partial_z \langle \mathbf{A}' \partial_z K^* \rangle - \epsilon_{K^*}, \quad (9)$$

where the first term on the RHS of (9) includes advection of IKE by both the ensemble mean and the turbulent flow, and the underlined term is associated with eddy-mean flow interactions. Again, the exchange of IKE with internal potential energy is made explicit through the inclusion of  $\langle w' b' \rangle$ . The sum of (8) and (9) leads to an equation for the ensemble mean kinetic energy of the full flow.

Another, yet equivalent, procedure to derive an equation for the ensemble mean kinetic energy of the full flow consists in expanding the different components of (4) following the Reynolds decomposition in the ensemble dimension (5), then ensemble averaging. This leads to:

$$\begin{aligned} \partial_t \langle K \rangle = & -\nabla \cdot \langle \mathbf{u} \rangle \tilde{K} - \nabla \cdot \langle \mathbf{u} K^* \rangle - \rho_0 \nabla \cdot \langle \mathbf{u}' (\langle \mathbf{u}_h \rangle \cdot \mathbf{u}'_h) \rangle \\ & - \nabla \cdot \langle \mathbf{u} \rangle \langle p \rangle - \nabla \cdot \langle \mathbf{u}' p' \rangle - \langle w \rangle \langle b \rangle - \langle w' b' \rangle + \rho_0 \partial_z \left( \langle \mathbf{A} \rangle \partial_z \tilde{K} \right) + \rho_0 \partial_z \langle \mathbf{A}' \partial_z K^* \rangle - \epsilon_{\tilde{K}} - \epsilon_{K^*}, \end{aligned} \quad (10)$$

where  $\epsilon_{\tilde{K}}$  and  $\epsilon_{K^*}$  represents dissipation of FKE and IKE, respectively. Here, the underlined term emerged from the advection of the cross energy term  $\langle \mathbf{u}_h \rangle \cdot \mathbf{u}'_h$  by the perturbations. This reflects that, although this term vanishes identically upon averaging, its advection by perturbations does not. This is of particular interest because it is associated with kinetic energy transfers between the mean and the turbulent flow, thus plays a critical role in eddy-mean flow interactions. Indeed, following the chain rule, this term can be decomposed as

$$\nabla \cdot \langle \mathbf{u}' (\langle \mathbf{u}_h \rangle \cdot \mathbf{u}'_h) \rangle = \langle \mathbf{u}_h \rangle \cdot \nabla \cdot \langle \mathbf{u}' \mathbf{u}'_h \rangle + \langle \mathbf{u}' \mathbf{u}'_h \rangle \cdot \nabla \langle \mathbf{u}_h \rangle, \quad (11)$$

where the continuity equation has been used to express the last term of the RHS of (11) in a more conventional way. The first term of the RHS of (11) is the covariance of the horizontal mean flow with the divergence of the Reynolds stress tensor associated with the FKE equation, and the second term of the RHS of (11) is the eddy momentum fluxes up or down the gradient of the mean flow associated with the IKE equation. It is then straightforward to show that expanding the underlined term in (10) as (11) leads to an equation for the ensemble mean kinetic energy of the full flow that equates the sum of the FKE and the IKE equation, i.e., Eq. (8) and Eq. (9). In the following, we will refer to the three terms of (11), from left to right, as the DIVERgence of Eddy Fluxes (DIVEF), the Mean-to-Eddy energy Conversion (MEC), and the EDDY momentum FLUX (EDDYFLX).

By volume integration, several components of (10) become statements about fluxes at the boundaries of the volume of integration through the divergence theorem. In ensemble simulations such as those we analyze here, ocean surface and boundary conditions are usually prescribed as ensemble mean conditions, common to all members, such that we can neglect turbulent fluxes at the (surface and open) boundaries. (This, along with bottom turbulent fluxes, are further discussed in Section 2.3). Under such assumptions, several terms of the integrated version of (10) vanish, and the domain integrated equation for the ensemble mean kinetic energy of the full flow simplifies to:

$$\begin{aligned} \partial_t \int_V \langle K \rangle dV = & \partial_t \int_V \tilde{K} dV + \partial_t \int_V \langle K^* \rangle dV = \\ & - \int_S \langle \mathbf{u} \rangle \tilde{K} \cdot \mathbf{n} dS - \int_S \langle \mathbf{u} \rangle \langle p \rangle \cdot \mathbf{n} dS - \int_V (\langle w \rangle \langle b \rangle + \langle w' b' \rangle) dV \\ & + \int_A \langle \mathbf{u}_h \rangle \cdot \langle \boldsymbol{\tau} \rangle dA - \int_B \langle \mathbf{u}_h \rangle \cdot \langle \mathbf{F} \rangle dB - \int_V (\epsilon_{\tilde{K}} + \epsilon_{K^*}) dV, \end{aligned} \quad (12)$$

where  $V$  is the volume of integration,  $S$  the surface bounding  $V$ ,  $A$  and  $B$  its ocean surface and bottom part, respectively, and  $\mathbf{n}$  the normal to the surface  $S$ . Here, the work done by surface wind stress and bottom friction ( $\int_A \langle \mathbf{u}_h \rangle \cdot \langle \boldsymbol{\tau} \rangle dA$  and  $\int_B \langle \mathbf{u}_h \rangle \cdot \langle \mathbf{F} \rangle dB$  with  $\mathbf{F}$  the vertical diffusive flux at the bottom boundary, respectively) comes from the volume integration of viscous forces. The time rate of change of kinetic energy within the domain thus reflects the import/export of FKE and the wave field prescribed at the open boundaries (two first terms), exchanges with potential energy (third term), work associated with prescribed surface forcing (fourth term) and bottom boundary condition (fifth term), and dissipation (last term). We note here that although the transfers of kinetic energy between the mean and the turbulent flow (underlined term in (10)) can

be locally large, they cancel each other when integrated over the entire basin to satisfy the boundary condition of no turbulent flux of the LHS of (11).

The turbulent version of (12) summarizes as:

$$\partial_t \int_V \langle K^* \rangle dV = -\rho_0 \int_V \langle \mathbf{u}' \mathbf{u}'_h \rangle \cdot \nabla \langle \mathbf{u}_h \rangle dV - \int_V \langle w' b' \rangle dV - \int_V \epsilon_{K^*} dV, \quad (13)$$

where the first term of the RHS of (13) comes from the development of (11). In a basin integrated sense, the time rate of change of IKE as diagnosed through ensemble statistics is thus a balance between exchanges with FKE, exchanges with eddy potential energy, and dissipation<sup>2</sup>. It is not directly driven by prescribed forcing, but rather reflects the part of the ocean intrinsic dynamics that develops spontaneously in response to the non-linearity of the system. This provides an energy-budget based rationalization that the ensemble strategy provides an unambiguous definition of the ocean *turbulence*. In the following, we pay a particular attention to the contribution of EDDYFLX for the construction of IKE, and its relation to the mean flow (MEC) through the flux divergence DIVEF.

## 2.2 Model and Simulations

We analyze in this study a subset of the MEDWEST60 ensemble simulations (Leroux et al., 2021). These simulations have been produced to evaluate the predictability of the fine scale dynamics in a typical high-resolution Copernicus Marine Environment Monitoring Service (CMEMS) forecasting model by including the effect of initial and model uncertainties. It is based on a kilometric-scale regional configuration of the Western Mediterranean sea (cf Fig. 1) that uses the same numerical choices as the North Atlantic simulation eNATL60 (Brodeau et al., 2020). Briefly, they are NEMO-v3.6 simulations run at  $\frac{1}{60}^\circ$  and with vertical grid spacing of 1 m at the surface and 24 m at depth, for a total of 212 vertical levels in MEDWEST60. The simulations are forced at the surface with 3-hourly ERA-interim (ECMWF) atmospheric reanalysis through the CORE bulk flux formulation (Large & Yeager, 2004), and they partially account for ocean-atmosphere feedbacks (e.g., Renault, Molemaker, McWilliams, et al., 2016), where only 50% of sur-

---

<sup>2</sup> Note that in our setup, horizontal dissipation is implicitly included in the UBS advective scheme.

As detailed in Section 2.3, such a contribution is neglected when interpreting numerical results. For a theoretical understanding, however, it can be considered as part of the dissipative term of (13).



face currents speed is considered in the computation of the wind stress. Open boundary conditions are applied at the eastern and western boundaries of the domain with a Flow Relaxation Scheme (FRS) for baroclinic velocities and active tracers (Davies, 1976; Engedahl, 1995), and the "Flather" (Flather, 1994) radiation scheme for sea-surface height and barotropic velocities. The former is a simple relaxation of model fields toward hourly, externally-specified values over the 12 grid points adjacent to the boundaries. The relaxation time scale ranges from  $\tau = 0$  seconds at the domain edge and increases exponentially to about 30 days at grid point 12. The latter ("Flather") applies radiation conditions on the normal depth-mean transport across the open boundaries, set as prescribed values plus a correction based on sea surface height anomalies at the boundaries that allows gravity waves generated within the domain to exit through the open boundaries. We note that the prescribed boundary conditions are taken from the eNATL60 North Atlantic experiment run with tidal forcing, such that MEDWEST60 includes tides through boundary conditions in addition to tidal potential forcing.

Among the various ensemble simulations produced in the context of MEDWEST60, we focus here on the 20-member ensemble ENS-CI-GSL19, which has been produced as follows. From the already spun-up (through a 18 months integration) oceanic state of the eNALT60 simulation at February, 5<sup>th</sup> 2010, an ensemble of 20 runs has been produced for 1 day with a stochastic perturbation (Brankart et al., 2015) applied on the horizontal grid of the model to represent uncertainties affecting the smallest scales in the model (for more details, see Leroux et al., 2021). The 20 oceanic states so generated have then been used as initial conditions for the production of a 120-day long, 20-member ensemble where all other components of the simulation (including forcing) are common across all members, and the stochastic perturbations are turned off. Such a procedure is usually referred to as *micro* initial condition uncertainties (Stainforth et al., 2007; Hawkins et al., 2016), and is meant to allow the growth of dynamically consistent small perturbations.

### 2.3 Diagnostic Considerations

During the production of MEDWEST60 ensemble simulations, prognostic variables of the model (T, S, U, V, SSH), as well as vertical velocity (W), have been saved every hour. Based on these hourly averaged model outputs, we have used *offline* diagnostic tools to recompute the kinetic energy budget of MEDWEST60 simulations by closely fol-

lowing the numerical implementations of NEMO<sup>3</sup>. Relevant details for the present analysis are provided in Appendix, along with validation. We note here that these *offline* tools, along with the high frequency of model outputs (hourly), provide us with a reliable procedure to accurately (errors  $\sim \mathcal{O}(10^{-3})$ , see Table A1) compute the kinetic energy trends due to advection, thus the terms associated with eddy-mean kinetic energy transfers.

In our kinetic energy budget considerations derived in Section 2.1, we have assumed zero turbulent fluxes conditions at the boundaries of the domain. In practice, however, the computation of surface wind stress partially (50%) accounts for ocean-atmosphere feedback (Renault, Molemaker, McWilliams, et al., 2016), such that the turbulent wind work  $\langle \mathbf{u}'_h \cdot \boldsymbol{\tau}' \rangle$  is not strictly zero. This contribution is however weak (-0.12 TJ; 1 TJ =  $10^{12}$  J) as compared to mean wind work (+5.10 TJ) over the course of the 120-day long simulation, and is several orders of magnitude smaller than the total IKE production of +2.27 PJ (1 PJ =  $10^{15}$  J) within the domain. Furthermore, this contribution is negative, providing a sink for domain integrated IKE time rate of change, in agreement with the eddy-killing effect (Renault, Molemaker, Gula, et al., 2016). Similar considerations are also relevant for turbulent bottom stress, which damps the production of IKE. Our estimates of surface and bottom velocities ensemble spread suggest the bottom contribution is at least one order of magnitude weaker than the surface contribution. As for the open boundary conditions, the "Flather" scheme allows gravity waves generated within the domain to exit the model through boundaries, thus providing an explicit sink of IKE. In an averaged sense, all members are however expected to exhibit similar levels of energy associated with the development of such waves, such that the spread so induced on model velocities is expected to be weak and can be neglected. We recall that baroclinic velocities are strongly relaxed toward prescribed values at the boundaries. The contribution of surface and boundary turbulent forcing, as well as bottom turbulent stress, for the interpretation of IKE production in our ensemble can then be safely neglected.

Finally, we are primarily interested in diagnosing eddy-mean flow kinetic energy transfers through the DIVEF, MEC and EDDYFLX terms of (11). As detailed above, open boundary conditions ensure that the ensemble spread at the boundaries is controlled, such that the domain integrated eddy fluxes of the cross energy term  $\rho_0 \nabla \cdot \langle \mathbf{u}' (\langle \mathbf{u}_h \rangle \cdot \mathbf{u}'_h) \rangle$  is negligible. This implies that all the energy released by the ensemble mean flow has been

---

<sup>3</sup> These tools are part of the CDFTOOLS (<https://github.com/meom-group/CDFTOOLS.git>)


used to sustain the growth of IKE within the domain. We have tested this by computing the volume integrated MEC and EDDYFLX terms for the full domain, and estimated their divergence DIVEF. We show on top panels of Figure 1 the vertically integrated MEC and EDDYFLX, and their divergence (DIVEF) is obtained by simple summation following (11). Integrated over the full domain, MEC drains -0.53 GW of energy out of the ensemble mean flow at that particular time (day 60), and EDDYFLX supplies +0.58 GW of energy to the turbulent flow. This confirms that our procedure provides reliable estimates of these fluxes, with a  $\sim 10\%$  error. This error, of about 0.05 GW, is relatively constant across the 20 ensemble members ( $\pm 0.01$  GW, Figure 1, lower panel), suggesting this is a systematic error in our estimates. We attribute this error to the implicit dissipation of the UBS advective scheme used in MEDWEST60. As detailed in Appendix, we have performed the eddy-mean flow decomposition of the advective operator based on a  $4^{th}$  order centered scheme, which is the non-dissipative equivalent of the UBS scheme. The error in our estimates being positive and relatively constant across ensemble members, this suggests it is associated with dissipation.

In the following sections, we turn our attention to the analysis of the MEDWEST60-ENS-CI-GSL19 ensemble simulations, where we first diagnose the decorrelation of the turbulent flow from its ensemble mean, then evaluate the respective contribution of MEC and EDDYFLX for the kinetic energy budget of the ensemble mean and the turbulent flow, and then analyze their interactions through DIVEF.

### 3 Results

#### 3.1 Decorrelation of the Turbulent Flow

Figure 2 provides horizontal maps and time evolution of surface kinetic energy, as well as its ensemble statistical decomposition. From left to right, the upper panels show the ensemble mean surface kinetic energy of the full flow  $\langle K \rangle$ , the FKE and the IKE at day 60. Their time evolution over the course of the 120 days, integrated within the green box, are shown on the lower panel. The ensemble mean full kinetic energy  $\langle K \rangle$  exhibits a combination of high and low frequency variations, but remains relatively constant (6-8 TJ; 1 TJ= $10^{12}$  J) over the 120 days, reflecting the already spun-up state of the eNATL60 simulation used to initialize the ensemble. For reference, the level of kinetic energy of a given member is shown in light gray. It exhibits small variations around its ensemble



**Figure 1.** (Top panels) Vertically integrated MEC ( $-\rho_0 \langle \mathbf{u}_h \rangle \cdot \nabla \cdot \langle \mathbf{u}' \mathbf{u}'_h \rangle$ , left panel), EDDYFLX ( $-\rho_0 \langle \mathbf{u}' \mathbf{u}'_h \rangle \cdot \nabla \langle \mathbf{u}_h \rangle$ , center panel), and DIVEF ( $-\rho_0 \nabla \cdot \langle \mathbf{u}' (\langle \mathbf{u}_h \rangle \cdot \mathbf{u}'_h) \rangle$ , right panel) after 60 days of simulation. Their volume integrated values are shown at the bottom right of each panels. (Bottom panel) Basin integrated MEC (black), EDDYFLX (red) and DIVEF (blue) for each individual members.


mean equivalent, illustrating that the ensemble mean kinetic energy of the full flow provides a statistical estimate of the energy level of the ensemble. We note that the deviation of the kinetic energy of a single member from the ensemble mean kinetic energy is not to be confused with the separation between the kinetic energy of the ensemble mean flow and that of the perturbations, which is the primary focus of our study.

The spatial pattern of the FKE ( $\tilde{K}$ ) is representative of the relatively well organized flow within the western Mediterranean basin. In the northern half, the FKE exhibits high levels of energy associated with the southwestward flowing Liguro-Provençal current (Millot, 1999; Waldman, 2016). In the southern half, FKE exhibits a very large import of energy through the strait of Gibraltar (exceeding  $2000 J m^{-3}$ ), the development of standing eddies downstream, and an eastward flowing boundary current along

the southern boundary of the basin (the Algerian Current, Millot, 1985). Around 5°E, this boundary current detaches from the coast, forming a 'loop current', a region of intense meso-scale eddies formation through mixed baroclinic-barotropic instabilities (e.g. Obaton et al., 2000; Poulain et al., 2021). We will focus on the eddy dynamics of this region in the following. Although more pronounced in the southern than in the northern part of the domain, the spatial organization of IKE ( $\langle K^* \rangle$ ) somehow follows the spatial organization of FKE, reflecting the link between the two; turbulent dynamics develop in region of strong currents, which are more prone to instabilities.

The lower panel of Fig. 2 illustrates the time evolution of surface FKE and IKE, integrated within the green box, during the 120 days of simulation. At the beginning all ensemble members are in phase, such that IKE is zero and FKE reflects the energy content of the full flow. The latter diverges from the ensemble mean full KE about one week after initialization as each ensemble member starts to decorrelate. At the end of the 120 days, FKE has dropped to less than 2 TJ, i.e., about one third of its initial energy content. In the same time, the turbulent part of the flow (IKE,  $\langle K^* \rangle$ ) develops and reaches about 5 TJ at the end of the 120 days. This development exhibits several stages until it saturates after about 80 days. It is interesting to note that a first increase in IKE is observed from day 6 to day 20, where it reaches a first plateau. The 6 day time scale for the turbulent flow to start decorrelating from initial conditions is consistent with time scale reported by Fox-Kemper et al. (2008) and Schubert et al. (2020) in their idealized linear study of mixed layer instability and absorption of submesoscale vortices by mesoscale eddies, respectively. In both studies, time scales shorter than one week are associated with the development of submesoscale structures through surface mixed layer instabilities, which then saturate and undergo non-linear interactions to transfer their energy upscale. The 6 days time scale in our ensemble simulations is thus likely associated with similar processes, and suggests that the non-linear interactions of submesoscale instabilities are responsible for the initial growth of IKE. The other stages of IKE increase are associated with further development of turbulent flow. By comparing the IKE patterns at days 30 and 60 for instance (not shown), it appears that initial IKE development mostly takes place along the mean current, while later on, turbulent structures develop more broadly, contributing to the increase in the integrated IKE level within the green box. Further spectral estimates of the decorrelation of ensemble members over the first 60 days can be found in Fig. 6a of Leroux et al. (2021). In what follows, we will fo-

cus our analysis on day 60, which is about 20 days before the saturation of IKE. As shown in the following, this time period exhibits a well organized spatial structure in the eddy-mean flow KE interactions that nicely illustrates non-local processes. Such processes are nonetheless observed all along the 120-day long simulation <sup>4</sup>.



**Figure 2.** (Upper panels) Spatial maps of surface currents ensemble mean kinetic energy of the full flow ( $\langle K \rangle$ ; left), kinetic energy of the ensemble mean flow ( $\tilde{K}$ , FKE; center) and the ensemble mean kinetic energy of the turbulent flow ( $\langle K^* \rangle$ , IKE; right) after 60 days of simulation. (Lower panel) 120-day long time series of these quantities, integrated within the green box. The time series of the kinetic energy of a given member is provided for reference (gray line). Units of the spatial maps are  $\text{J m}^{-3}$  and those of the time series are terrajoules (1 TJ =  $10^{12}$  J). The black box on top left panel is used to validate our recomputation of kinetic energy budgets (cf Appendix).

<sup>4</sup> The interested reader is referred to the following animation: [https://doi.org/10.5281/zenodo](https://doi.org/10.5281/zenodo.6221153)

### 3.2 Kinetic Energy Budget

We now turn our attention to the respective contributions of the advective terms of the FKE and IKE budget, focusing on the 'loop current' region. We recall here that many other processes contribute to these budgets, such as wave radiation, dissipation or exchanges with turbulent potential energy (cf (10)). We briefly discuss the contribution of the latter in what follows, but otherwise postpone the analysis of other contributions for further work. Here, we focus our attention on the terms driving kinetic energy transfers between the mean and the turbulent flow. We first discuss the kinetic energy budget of the mean flow and that of the turbulent flow, and estimate the respective contribution of MEC and EDDYFLX to these budgets.

We show on Fig. 3 the vertically integrated time rate of change of FKE (top left panel), as well as advection of FKE by the mean flow ( $-\nabla \cdot \mathbf{u} \tilde{K}$ ; top right panel) and Mean-to-Eddy Conversion (MEC,  $-\rho_0 \langle \mathbf{u}_h \rangle \cdot \nabla \cdot \langle \mathbf{u}' \mathbf{u}'_h \rangle$ ; bottom left panel) at day 60. Their vertical distributions within the upper 500 meters, horizontally integrated within the green box, appear on the bottom right panel as black, blue and red lines, respectively, and the contribution from other processes (computed as a residual) is shown in green<sup>5</sup>. We first note that the time rate of change of FKE is dominated by a wave-like horizontal structure, which exhibits a strong baroclinic signature. The fast (daily) evolution of this signal (not shown) suggests it is associated with the high frequency signal observed in the FKE time series of surface currents (Fig. 2, bottom panel). As part of the ensemble mean flow, this signal is likely associated with the forcing, such as high frequency winds and, to a smaller extent, tidal forcing. Integrated within the green box, this leads to a time rate of change of FKE of +0.30 GW. In contrast, both advection of FKE by the mean flow and MEC exhibit very different patterns with smaller scale structures. The former exhibits a multipole-like organization, and its contribution tends to be of opposite sign in the upper 50 m (i.e., deeper than the ensemble mean and spatially averaged mixed layer depth of about 30 m) and the rest of the water column, such that when volume integrated, its contribution is two orders of magnitude weaker than the volume integrated time rate of change of FKE. Although MEC exhibits weaker signal lo-

---


<sup>5</sup> Note that all horizontal maps have been integrated down to the ocean floor for consistency, but most of the dynamics is observed within the upper 500 meters

cally, its volume integrated contribution is significant ( $-0.24$  GW), with a maximum at about 40 m depth.

Fig. 4 shows the equivalent of Fig. 3 but for the IKE budget. We first note that the spatial pattern of IKE time rate of change is significantly different from that of FKE, with smaller scale structures. Contribution of advection of IKE by the mean and turbulent flow within the box is weak ( $+0.03$  GW), but exhibits local important contributions for the IKE redistribution. EDDYFLX contributes to  $+0.25$  GW to the budget, which slightly exceeds the time rate of change of IKE of  $+0.21$  GW. The vertical profile of turbulent potential to kinetic energy conversion rate  $-\langle w'b' \rangle$  is also shown, with a net contribution within the green box of about  $+0.20$  GW. It is maximum at about 30 meters depth and tends toward zero at the surface. Although relatively weak when integrated within the green box ( $-0.08$  GW), the large intensification of the residual near the surface is expected to mostly reflect the action of vertical viscous forces and dissipation.


Finally, we quantify the contribution of EDDYFLX for construction of the IKE over the course of the 120 days of simulations, and assess its relation with the loss of energy of the mean flow through MEC. This we address by computing the volume integrated contribution of both EDDYFLX and MEC within the green box of Fig. 2 for the 120 day long simulations. We show on Fig. 5 the time series of these two contributions (left panel), as well as their time integrated estimates (right panel). Starting from zero at the beginning of the simulations where all ensemble members are in phase, EDDYFLX starts to inject energy in the turbulent flow after about 5-6 days, in agreement with surface IKE increase discussed in Section 3.1. The rate at which EDDYFLX inject energy in the turbulent flow is of about  $0.2 \text{ GJ s}^{-1}$  with time variations as large as  $\pm 0.13 \text{ GJ s}^{-1}$ . MEC is draining energy out of the mean flow with similar rate and temporal variations, leading to a small contribution of DIVEF (light blue line). Over the course of the 120 days of simulation, EDDYFLX and MEC have contributed to  $+2.41$  PJ and  $-2.12$  PJ for the IKE and FKE budget, respectively (Fig. 5, right panel). Within this region, the contribution of DIVEF is small, suggesting that eddy-mean energy transfers within this box are mostly local. Also shown on this figure is the contribution of the turbulent potential to kinetic energy conversion rate  $-\langle w'b' \rangle$ . We first note the very large temporal variations in this term as compared to eddy-mean flow interaction processes, suggesting intense exchanges with turbulent potential energy reservoirs on very short time scales. Their





**Figure 3.** Vertically integrated time rate of change of FKE (upper left panel), advection of FKE by the mean flow (upper right panel) and Mean-to-Eddy energy Conversion rate (MEC, lower left panel) in the region of the loop current at day 60, with their volume integrated values within the green box shown at the bottom right of each panels. The vertical distribution of these quantities, within the upper 500 meters and horizontally integrated within the green box, are shown on the bottom right panel. The other components of the FKE budget are shown as a residual (green line).


time integrated contribution, however, is of the same order of magnitude than EDDYFLX but slightly weaker, supporting mixed barotropic-baroclinic instability processes for driv-



**Figure 4.** Same as Fig. 3, but for the IKE budget. The advection of IKE (upper right panel) includes advection by both the mean flow ( $-\nabla \cdot \langle \mathbf{u} \rangle \langle K^* \rangle$ ) and the turbulent flow ( $-\nabla \cdot \langle \mathbf{u}' K^* \rangle$ ). Turbulent potential to kinetic energy conversion rate ( $-\langle w'b' \rangle$ ) is also shown in gray and its net contribution within the green box is of about +0.20 GW. Note the change in amplitude of the colorbar as compared to Fig. 3.

ing the growth of Algerian Eddies as proposed earlier (Obaton et al., 2000; Poulain et al., 2021). It is interesting to compare these estimates to the total IKE and FKE changes. During the 120 days of simulation, the volume integrated IKE within the green box has grown by +0.98 PJ, which is only about a quarter of the total energy injected by ED-

483 DYFLX and  $-\langle w'b' \rangle$ . Similarly, the FKE destruction over the full simulation is -0.91  
 484 PJ, which is about half of the energy drained by MEC. This highlights the leading or-  
 485 der contribution of other processes for balancing kinetic energy budgets of this region.



**Figure 5.** (Left) Time series of volume integrated MEC (green), EDDYFLX (red), DIVEF (light blue) and  $-\langle w'b' \rangle$  (gray) within the green box of Fig. 2, and (right) their time integrated contribution. The 120-day long integrated MEC (EDDYFLX, DIVEF,  $-\langle w'b' \rangle$ ) contribution is -2.12 PJ (+2.41 PJ, +0.30 PJ, +1.38 PJ).

## 486 4 Non-locality of FKE-IKE Energy Transfers

### 487 4.1 Diagnosing Non-Local KE Transfers

488 The patterns and amplitude of MEC and EDDYFLX discussed in the previous sec-  
 489 tion are associated with energy transfers between the mean and the turbulent flow. As  
 490 discussed in Section 2.1, this can reflect either local processes, with a negligible contri-  
 491 bution of DIVEF (LHS of (11)), or non-local processes, with energy transfers with tur-  
 492 bulent processes of remote regions if that term is not negligible. Dynamically, this pro-  
 493 vides an estimate of the level of energy released by the mean flow that *locally* sustains  
 494 the growth of eddies. Or, vice versa, an estimate of the level of energy released by the  
 495 eddies that is *locally* backscattered to energize the mean flow. We further analyze this  
 496 local vs non-local contribution in what follows.

497 Horizontal maps of vertically integrated MEC, EDDYFLX and DIVEF are shown  
 498 in Figure 6 at day 60, and their volume integrated values within the green box appear

at the bottom right of each panel. Averaged over the box, the energy lost by the mean flow (MEC, -0.24 GW) is used to support eddy growth (EDDYFLX, +0.25 GW), and the divergence of eddy fluxes is weak (DIVEF, +0.01 GW). That MEC is draining -2.12 PJ out from FKE and EDDYFLX is injecting +2.41 PJ into IKE during the 120 days of simulation, as diagnosed in Section 3.2, also supports that the turbulence that develops within the green box is largely controlled by local processes. However, the details of these energy transfers is complex, and the radically different spatial structure of MEC and EDDYFLX strongly suggests that eddy-mean flow kinetic energy transfers are non-local at small scales. The spatial scale dependence of these non-local transfers is further analyzed in Section 4.2.


At day 60, the horizontal structure of MEC (Fig. 6, left panel) exhibits alternation of FKE destruction (blue spots) with FKE production (red spot), which tend to organize mostly along the mean flow. In contrast, EDDYFLX (Fig. 6, middle panel) exhibits signals of weaker amplitude, which tend to be more pronounced on the flanks of the flow. This suggests that a significant part of the kinetic energy lost by the mean flow at one location is advected further downstream before being re-injected in the mean flow, but little is used to sustain the growth of eddies locally. The connection between MEC and EDDYFLX involves DIVEF, which is associated with eddy flux divergence of the cross energy term  $\langle \mathbf{u}_h \rangle \cdot \mathbf{u}'_h$ . This term exhibits a rich spatial organization (Fig. 6, right panel), with regions of destruction of FKE associated with a divergence of eddy fluxes, i.e., the cross energy term is fluxed out of the control volume by the turbulent flow, and regions of FKE production associated with a convergence of eddy fluxes, i.e., the cross energy term is fluxed within the controlled volume by the turbulent flow. The region indicated by the black line is of particular interest because it exhibits a region of production of IKE (red spot of EDDYFLX) to the northeast of the region of FKE destruction. MEC, EDDYFLX and DIVEF vertical cross sections along this line are shown in Fig. 7. At the surface, MEC exhibits its largest negative value about 10 km away from the core of the mean current, and follows its tilted vertical structure. In contrast, the EDDYFLX is largest about 20 km northeastward of the minimum of MEC, a region of strong horizontal mean flow gradient, but exhibits a shallower vertical penetration as compared to MEC. As a result, DIVEF is dominated by a divergence of eddy fluxes near the core of the mean flow, and a convergence on its flank. Although a direct interpretation of a turbulent flux of the cross energy term  $\langle \mathbf{u}_h \rangle \cdot \mathbf{u}'_h$  to connect these regions of FKE destruction and IKE

production is tempting, we recall here that this term vanishes identically for turbulent flow orthogonal to the mean flow. This suggests that DIVEF is more efficient at transporting energy in the along stream direction than in the across stream direction, providing a strong horizontal constraint for eddy-mean flow interactions. This may well provide a dynamical rationalization to explain the large variations of MEC observed in the along stream direction, where energy extracted from the mean flow would be transported downstream before to be reinjected into the mean flow, but little would actually be transferred to the turbulent flow through EDDYFLX.


Fig. 8 shows the horizontal and vertical contribution for the three components involved in eddy-mean flow kinetic energy transfers in the upper ocean layer. We first note that, as expected, vertical fluxes are much weaker than horizontal fluxes. However, while weak at each location, vertical turbulent fluxes are predominately positive in the upper layer, such that their horizontally integrated contribution is of the same order of magnitude than the horizontal turbulent fluxes for the three terms (Fig. 9). More interestingly, while the horizontal component of MEC and EDDYFLX tend to oppose each other, the vertical components tend to have the same sign. Indeed, the horizontal contribution of MEC is relatively constant and negative in the upper 100 meters and smoothly decreases further below (left panel), while the horizontal contribution of EDDYFLX is negligible at the surface, reaches its maximum at about 30 meters and smoothly decreases further below (center panel). In contrast, in both MEC and EDDYFLX, vertical turbulent fluxes are upward in the upper 15 meters, reach a maximum downward contribution at the base of the spatially averaged mixed layer (about 30 meters), and decrease further below to reach negligible contribution below about 100 meter. The balanced DIVEF within the green box (right panel) thus results in a balance between horizontal MEC and EDDYFLX below 100 meters, but involves strong contributions from the vertical turbulent fluxes within the upper 100 meters, with a prominent downward turbulent flux across the base of the of the mixed layer. Our results thus highlight the leading order contribution of vertical turbulent fluxes in eddy-mean flow kinetic energy interactions at the base of the mixed layer.

## 4.2 Horizontal Scale Dependence

Finally, we assess the scale-dependence of these non-local energy transfers. Although at small scales, our results suggest that eddy-mean flow interactions are largely non-local,




**Figure 6.** Vertically integrated MEC ( $-\rho_0 \langle \mathbf{u}_h \rangle \cdot \nabla \cdot \langle \mathbf{u}' \mathbf{u}'_h \rangle$ , left panel) EDDYFLX ( $-\rho_0 \langle \mathbf{u}' \mathbf{u}'_h \rangle \cdot \nabla \langle \mathbf{u}_h \rangle$ , middle panel) and DIVEF ( $-\rho_0 \nabla \cdot \langle \mathbf{u}' (\langle \mathbf{u}_h \rangle \cdot \mathbf{u}'_h) \rangle$ , right panel) after 60 days of simulations within the loop current region. Integrated quantities within the green box are shown on the bottom right insert. Ensemble mean surface currents are shown with arrows, and the black line is the section shown in Fig. 7.



**Figure 7.** Associated vertical structure of MEC, EDDYFLX and DIVEF along the cross-stream section of Fig. 6. Gray contours represent the ensemble mean current across the section. Dashed green contours on middle and right panels show the main structure of MEC.


our estimates on larger scales tend toward a local balance (i.e., DIVEF is negligible). This is true for the  $3^\circ \times 3^\circ$  green box of Fig. 6, as well as for other places in the western Mediterranean basin (not shown). This suggests non-local effects are predominantly small scale features, such that we are interested in quantifying their scale dependence. To test this, we have computed the spatial correlation  $r$  between MEC and EDDYFLX as a function





**Figure 8.** Upper layer MEC (left), EDDYFLX (center) and DIVEF (right) at day 60, decomposed into an horizontal (top panels) and a vertical (bottom panels) contribution. Ensemble mean surface currents are shown with arrows.

of coarse grained grid size (Figure 10). Starting from the initial model grid size at  $\frac{1}{60}^\circ$ , a spatial averaging is performed with the adjacent grid points, i.e., a factor 3, up to a grid size of about  $4^\circ$ . This procedure has been performed on four different boxes of  $3^6 \times 3^6$  (i.e.,  $729 \times 729$ ) grid points (colored lines) in order to cover the entire  $883 \times 803$  grid points MEDWEST60 domain. The spatial correlation between MEC and EDDYFLX ranges from -0.12 on average at the model grid size to -0.96 at about  $4^\circ$ . This suggests that although non-local at small scales, kinetic energy transfers can be seen as local processes for scales larger than a few hundreds of kilometers. However, for eddy-resolving ocean models ( $\sim \frac{1}{12}^\circ$ ), such as those that will equip the next generation climate models, non-




**Figure 9.** Vertical profile of horizontally integrated MEC (left), EDDYFLX (center) and DIVEF (right) within the green box of Fig. 6. Three-dimensional estimates (black) are decomposed into an horizontal (blue) and vertical (red) contribution. Positive vertical eddy fluxes are oriented upward, and the dashed gray line represent the spatially averaged mixed layer depth at about 30 meters.

local eddy-mean energy transfers are large ( $r < -0.2$ ). This suggests that the processes associated with this non-locality need to be accounted for in the development of parameterizations for eddy-resolving ocean models.

## 5 Conclusion

In this study, we have investigated the spatio-temporal structure of the kinetic energy transfers between the ensemble mean and the turbulent flow. We have performed our analysis with a kilometric-scale resolution ( $\frac{1}{60}^\circ$ ), 120-day long, 20-member ensemble simulations of the Western Mediterranean basin (Leroux et al., 2021). We have first introduced the Forced and Internal Kinetic Energy equation (FKE and IKE, respectively) in this framework, and discussed the implications for their interpretation. In particular, the prescribed surface and boundary forcings drive the basin integrated time rate of change of FKE, and the basin integrated time rate of change of IKE reflects the energy of the turbulent flow that develops within the domain through the non-linear dynamics sensitive to initial conditions. We have then quantified the respective contributions of Mean-to-Eddy energy Conversion (MEC,  $\langle \mathbf{u}_h \rangle \cdot \nabla \cdot \langle \mathbf{u}' \mathbf{u}'_h \rangle$ ) and the EDDY momentum FLUX (EDDYFLX,  $\langle \mathbf{u}' \mathbf{u}'_h \rangle \cdot \nabla \langle \mathbf{u}_h \rangle$ ) in the FKE and IKE budgets during the 120-day long runs.





**Figure 10.** Spatial correlation of MEC and EDDYFLX as a function of the coarse grained grid size at day 60. The local vs non-local computation is made on four  $3^6 \times 3^6$  regions (color lines and insert) and the results averaged (black line). The associated coarse grained grid size is shown on top axis.

By further analyzing their spatial organization, we have then highlighted the non-locality of the energy transfers between the ensemble mean and the turbulent flow, where non-local processes are associated with energy destruction in one reservoir that does not *locally* sustain the growth of kinetic energy in the other reservoir, in agreement with previous studies (Chen et al., 2014; Kang & Curchitser, 2015; Capó et al., 2019). We have pointed out to the leading contribution of the DIVERgence of Eddy Fluxes (DIVEF,  $\nabla \cdot \langle \mathbf{u}'(\langle \mathbf{u}_h \rangle \cdot \mathbf{u}_h') \rangle$ ) as a key component in this non-local transfers. Our main contribution is to recognize that this term is associated with advection of the cross energy term  $\langle \mathbf{u}_h \rangle \cdot \mathbf{u}_h'$  by the turbulent flow, which provides a strong spatial constraint on these transfers since the cross energy term vanishes identically for turbulent flow orthogonal to the mean flow. Finally, we have shown that although weaker than the horizontal component at the model grid size, the vertical eddy fluxes become leading order when horizontally integrated over sufficiently large scales. On average, their contribution is to flux energy (mean, eddy and cross energy term) downward across the base of the mixed layer.

Analyzing the scale dependence of these non-local KE transfers, we have shown that, although prevalent at eddy scales, they tend toward a local balance at non-eddy scale

(i.e.,  $> 1^\circ$ ). Thus, while our results support approximations usually made in the development of energy-aware parameterizations of meso-scale turbulence (Eden & Greatbatch, 2008; Mak et al., 2018; Jansen et al., 2019), i.e., that the growth of sub-grid scale turbulent kinetic energy is locally sustained by a destruction of kinetic energy of the resolved flow, they point out to the necessity of accounting for this non-local dynamics for the development of parameterizations for eddy-resolving ocean models, such as those that will equip next generation climate models. In this direction, the emerging approach of *transport under location uncertainty* for the representation of small scale, stochastic dynamics and its effect on the large scale flow (e.g., Mémin, 2014; Resseguier et al., 2017; Chapron et al., 2018) is an attractive alternative to the mixing length approach.

Finally, we want to discuss the implications of our results for the interpretation of the dynamics of western boundary currents jet extension such as the Gulf Stream. Jamet et al. (2021) have recently shown the leading order contribution of MEC for the energetic balance of the North Atlantic subtropical, wind driven gyre. They concluded that MEC in the Gulf Stream extension region is the primary sink of 26-year mean kinetic energy within the gyre, balancing the energy inputted by the wind in the westerly wind region of the North Atlantic subtropical gyre. However, how this loss of mean kinetic energy interacts with the turbulent flow remains an open question. Some indications of spatial organization of EDDYFLX can be found in previous in-situ and satellite observation analyzes. In their earlier work on Gulf Stream energetics based on in-situ observations, Webster (1961, 1965), Rossby (1987) and Dewar and Bane (1989) have reported on eddy fluxes that are more pronounced on the inshore flank of the Gulf Stream, both along the US coastline and downstream of Cap Hatteras. Based on satellite observations, Ducet and Le Traon (2001) and Greatbatch et al. (2010) have highlighted a prominent feature of the Gulf Stream, so-called the 'double-blade' structure, associated with the turbulent dynamics just downstream of Cape Hatteras. There, the Reynolds stress cross-covariance was found to be maximum on both flanks on the stream, and to exhibit alternation of highs and lows further downstream. This 'double-blade' structure suggests that eddy fluxes (EDDYFLX) are more pronounced on the flank of the jet, where large Reynolds stresses  $\overline{u'v'}$  are colocalized with a strong horizontal shear of the mean flow  $\partial_y \bar{u}$ , while mean-to-eddy conversion rates (MEC) would be more pronounced toward the core of the jet, where the cross-stream gradient of Reynolds stresses  $\partial_y \overline{u'v'}$  are colocalized with maximum of the mean zonal current  $\bar{u}$ . We can also find some indications of

such a spatial organization of eddy-mean flow interactions in the Lorenz energy cycle based on eddy-resolving numerical simulations of Kang and Curchitser (2015), although further analyses are needed to conclude on this.

## Appendix A Offline Recomputation of Kinetic Energy Budget

We are interested in analyzing the energetic of the MEDWEST60 ensemble simulations, which have been recently produced (Leroux et al., 2021). We thus developed diagnostic tools to recompute the momentum budget, which kinetic energy builds upon, of these simulations based on the variables saved during the production of these simulations, i.e. three-dimensional temperature (T), salinity (S) and velocity (U, V, W), as well as two-dimensional free-surface elevation (SSH). These *offline* tools are developed as part of the CDFTOOLS diagnostic package for the analysis of NEMO model output (<https://github.com/meom-group/CDFTOOLS.git>), which are written in FORTRAN 90 and follow the numerical implementation of the NEMO General Circulation Model (Madec et al., 2017).

As all GCM, NEMO offers different numerical schemes to integrate the Primitive Equations with various levels of approximation. The numerical schemes that have currently been implemented in these tools are those relevant for the analysis of the energetic of the MEDWEST60 ensemble simulations, which are based on the version 3.6 of the NEMO model. This includes: A dynamical vertical coordinate following the free surface elevation, with partial stepping along the ocean floor; the third order upstream biased scheme (UBS, Shchepetkin & McWilliams, 2005) to advect momentum; the TEOS-10 equation of state (Roquet et al., 2015) to compute density; a split-explicit formulation to compute surface pressure gradients (Shchepetkin & McWilliams, 2005), which also accounts for atmospheric surface pressure loading and freshwater air-land-sea fluxes; and an implicit time differencing scheme to compute vertical viscous effects, which include surface wind stress forcing following the CORE bulk flux formulation (Large & Yeager, 2004), bottom friction due to bottom boundary condition, tides, internal waves breaking and other short time scale currents, as well as vertical dissipation of momentum within the water column based on the Turbulent Kinetic Energy (TKE) turbulent closure scheme (Mellor & Yamada, 1982; Gaspar et al., 1990; Blanke & Delecluse, 1993). A full description of these schemes is available online ([https://github.com/quentinjamet/CDFTOOLS/blob/cdfdyn/note\\_KE.bgt\\_cdftools.pdf](https://github.com/quentinjamet/CDFTOOLS/blob/cdfdyn/note_KE.bgt_cdftools.pdf)). With shorthands, the full kinetic energy bud-

get can be represented as:

$$NXT = ADV + (HPG + SPG_{1st\ guess}) + SPG_{correction} + ZDF, \quad (A1)$$

where  $NXT$  refers to the time rate of change  $\partial_t$  (before application of the Asselin filter),  $ADV$  to three-dimensional advection,  $HPG$  to hydrostatic pressure work,  $SPG_{1st\ guess}$  to surface pressure work computed at baroclinic time step due to the rescaled vertical coordinate following free surface elevation,  $SPG_{correction}$  to surface pressure work correction associated with the time-splitting scheme of Shchepetkin and McWilliams (2005) which includes atmospheric pressure loading and freshwater fluxes, and  $ZDF$  to vertical viscous effects.

## A1 Validation at Model Time Step

In order to insure that our offline recomputation lines up with the online estimates computed by the NEMO model, we have re-run for a short period of time one member of the ensemble and outputted, at the model time step ( $\Delta t = 80s$ ), momentum and kinetic energy trends, as well as required prognostic variables necessary for their offline recomputation, within the 150x150 grid point sub-region (black box on Fig. 2). Comparing our offline recomputation with the online estimates provides an robust estimate of the errors. An example is provided on Fig. A1 for the three-dimensional advection of kinetic energy within the model upper layer. The errors are relatively small (locally four order of magnitude, but five order of magnitude when horizontally averaged within the sub-domain, cf Table A1), providing strong confidence in the accuracy of these tools. Tests for the other terms of the KE budget have been conducted, providing similar level of accuracy for time rate of change and pressure work (cf Table A1). Offline estimates of vertical viscous effects are associated with much larger errors, of the order of 10%, and we currently have no estimates for the surface pressure correction associated with the split-explicit scheme.

## A2 Estimation of Errors Due to Time Discretization and Averaging

Based on model time step accuracy estimates, we have quantified the errors associated with time discretization of the different operators, as well as the use of time averaged quantities. We discuss here these implications for the estimates of the advective component of the budget.

The advective operator used in the MEDWEST60 is an upstream biased third order scheme (UBS, Shchepetkin & McWilliams, 2005). This scheme has two component, a second order scheme and a third order biased scheme. While the former is centered in time, the latter is implemented forward in time, i.e. it is evaluated with *before* velocities. While this numerical detail provides stability for a GCM, it is not required in the context of *offline* computations and introduces ambiguities about how this should be evaluated when working with time averaged quantities. We thus decided to evaluate the third order biased scheme of the advective operator as centered in time instead. This leads to a growth of the errors made in the recomputation by one order of magnitude (cf Table A1). When computed based on hourly model outputs, as available from MEDWEST60, the error increases by another order of magnitude to reach  $10^{-3}$ . Also increased from model time step to hourly model outputs, the accuracy of these *offline* diagnostic tools remains high, providing reliable estimates of the advective operator of the model. Similar considerations are applied for the vertical viscous effects (i.e. time discretization, hourly model outputs), but the already large error of  $10^{-1}$  is found to be unchanged.


Finally, we estimate the evolution in time of these errors by comparing the recomputation made with hourly model outputs with estimates outputted by the model over a time period of 10 days (Figure A2). From these tests, no systematic errors emerged for both time rate of change (upper left panel) and hydrostatic pressure work (bottom left panel). We observe, however, a steady growth in the error made in the recomputation of the advective term (top right panel), reaching about  $-20 \times 10^{-3}$  GW h $^{-1}$  at the end of the 10 days of simulation. Finally, the largest errors are observed in the recomputation of the vertical viscous effects (bottom right panel), in agreement with errors reported earlier. We are currently working on improving this recomputation.

### A3 Eddy-mean Separation

Based on these *offline* estimates, we explicitly decompose the full equation into mean and eddy contributions. For the zonal momentum advection, it leads to:

$$\nabla \cdot \mathbf{u}u = \nabla \cdot \langle \mathbf{u} \rangle \langle u \rangle + \nabla \cdot \langle \mathbf{u} \rangle u' + \nabla \cdot \mathbf{u}' \langle u \rangle + \nabla \cdot \mathbf{u}' u' \quad (\text{A2})$$

where  $\langle \cdot \rangle$  and  $\cdot'$  denotes averaging and perturbation, respectively (cf Section 2.1 for details on the decomposition used in this study). Performing a similar procedure for the advection of meridional momentum, multiplying the former by  $\rho(\langle u \rangle + u')$  and the lat-




**Figure A1.** Upper layer Kinetic Energy trends associated with three-dimensional advection based on the model outputs (left), its offline recomputation (center), and associated errors (right). The *offline* recomputation is performed at model time step and accounts for the forward time discretization of the third order upstream biased part of UBS advective scheme. Note the different scale factor used for errors.

ter by  $\rho(\langle v \rangle + v')$  and summing the resulting equations leads to a decomposition of the advection of kinetic energy that accounts for the different contributions that compose the FKE and IKE budgets (equations (8) and (9), respectively). We note here that in MEDWEST60, the advection of momentum is achieved by the upstream biased third order scheme (UBS, Shchepetkin & McWilliams, 2005). This scheme accounts for the horizontal dissipation of momentum through an implicit formulation which takes the form of a biharmonic operator with an eddy coefficient proportional to the velocity  $A_h = -|u|\Delta x^3/12$ . The formulation of this implicit dissipation introduces complexities in the eddy-mean decomposition. We thus decided to evaluate the horizontal advection terms using a 4<sup>th</sup> order finite differencing centered scheme instead, which is the non-dissipative equivalent of the UBS scheme (Jouanno et al., 2016; Madec et al., 2017).

## Acknowledgments

This work was supported through the French ‘Make Our Planet Great Again’ program managed by the Agence Nationale de la Recherche under the Programme d’Investissement d’Avenir, with the reference ANR-18-MPGA-0002. MEDWEST60 ensemble simulations and their analyses have been performed using HPC resources from GENCI-IDRIS through



**Figure A2.** Time integrated KE trends of the full sub-domain, volume integrated time rate of change (upper left), three-dimensional advection (upper right), pressure work (bottom left) and vertical dissipation (bottom right) based on hourly averaged model outputs (black lines), recomputation based on hourly averaged T, S, U, V, W,  $\eta$  (red dots), and the associated errors (blue lines). Note the scale factor used for errors in the legend panels, which differs for each quantities.

**Table A1.** Order of magnitude of the errors of the offline estimates for the different terms of the kinetic energy budget, computed as the spatial root-mean-square error normalized by the spatial standard deviation of the reference, NEMO outputs. The third line stands for the sensitivity of the error associated with the forward time discretization of the third order upstream biased part of UBS advective scheme and in the TKE turbulent closure scheme. We currently have estimates for the surface pressure work correction associated with the split-explicit scheme (third term of the RHS), such that no values are reported on here.

	$\partial_t K$	$= - \nabla \cdot \mathbf{u} K$	$- \mathbf{u}_h \cdot \nabla_h \phi_{hyd}$	$- \mathbf{u}_h \cdot \nabla_h \phi_{surf}$	$+ \rho_0 \mathbf{u}_h \cdot \mathbf{D}^m$
Model time step	$10^{-3}$	$10^{-5}$	$10^{-5}$	–	$10^{-1}$
Time discretization	–	$10^{-4}$	–	–	$10^{-1}$
Hourly average	$10^{-2}$	$10^{-3}$	$10^{-3}$	–	$10^{-1}$

the Grant A008-0111279 allocated in the context of the H2020-IMMERSE projet (European Union Horizon 2020 research and innovation programme, grant No 821926).

## References

- Aluie, H., Hecht, M., & Vallis, G. K. (2018). Mapping the energy cascade in the North Atlantic Ocean: The coarse-graining approach. *J. Phys. Oceanogr.*, *48*(2), 225–244.
- Blanke, B., & Delecluse, P. (1993). Variability of the tropical Atlantic Ocean simulated by a general circulation model with two different mixed-layer physics. *J. Phys. Oceanogr.*, *23*(7), 1363–1388.
- Brankart, J.-M., Candille, G., Garnier, F., Calone, C., Melet, A., Bouttier, P.-A., Brasseur, P., and Verron, J. (2015). A generic approach to explicit simulation of uncertainty in the nemo ocean model. *Geoscientific Model Development*, *8*(5), 1285–1297.
- Brodeau, L., Le Sommer, J., & Albert, A. (2020). Ocean-next/eNATL60: Material describing the set-up and the assessment of NEMO-eNATL60 simulations (Version v1). [available online at :<http://doi.org/10.5281/zenodo.4032732>].
- Capó, E., Orfila, A., Mason, E., & Ruiz, S. (2019). Energy conversion routes in



- 767 the western mediterranean sea estimated from eddy-mean flow interactions. *J.*  
768 *Phys. Oceanogr.*, *49*(1), 247–267.
- 769 Cessi, P. (2008). An energy-constrained parameterization of eddy buoyancy flux. *J.*  
770 *Phys. Oceanogr.*, *38*(8), 1807–1819.
- 771 Chapron, B., Dérian, P., Mémin, E., & Resseguier, V. (2018). Large-scale flows un-  
772 der location uncertainty: a consistent stochastic framework. *Quarterly Journal*  
773 *of the Royal Meteorological Society*, *144*(710), 251–260.
- 774 Chen, R., Flierl, G. R., & Wunsch, C. (2014). A description of local and nonlo-  
775 cal eddy-mean flow interaction in a global eddy-permitting state estimate. *J.*  
776 *Phys. Oceanogr.*, *44*(9), 2336–2352.
- 777 Chen, R., Thompson, A. F., & Flierl, G. R. (2016). Time-dependent eddy-mean  
778 energy diagrams and their application to the ocean. *J. Phys. Oceanogr.*, *46*(9),  
779 2827–2850.
- 780 Davies, H. (1976). A lateral boundary formulation for multi-level prediction models.  
781 *Q. J. R. Meteorolog. Soc.*, *102*(432), 405–418.
- 782 Dewar, W. K., & Bane, J. M. (1989). Gulf stream dynamics. pad ii: Eddy energetics  
783 at 73° w. *Journal of Physical Oceanography*, *19*(10), 1574–1587.
- 784 Ducet, N., & Le Traon, P.-Y. (2001). A comparison of surface eddy kinetic energy  
785 and Reynolds stresses in the Gulf Stream and the Kuroshio Current systems  
786 from merged TOPEX/Poseidon and ERS-1/2 altimetric data. *Journal of*  
787 *Geophysical Research: Oceans*, *106*(C8), 16603–16622.
- 788 Eden, C., & Greatbatch, R. J. (2008). Towards a mesoscale eddy closure.  
789 *Ocean Model.*, *20*(3), 223–239.
- 790 Engedahl, H. (1995). Use of the flow relaxation scheme in a three-dimensional baro-  
791 clinic ocean model with realistic topography. *Tellus A*, *47*(3), 365–382.
- 792 Ferreira, D., Marshall, J., & Heimbach, P. (2005). Estimating eddy stresses by fit-  
793 ting dynamics to observations using a residual-mean ocean circulation model  
794 and its adjoint. *J. Phys. Oceanogr.*, *35*(10), 1891–1910.
- 795 Flather, R. A. (1994). A storm surge prediction model for the northern bay of ben-  
796 gal with application to the cyclone disaster in april 1991. *J. Phys. Oceanogr.*,  
797 *24*(1), 172–190.
- 798 Fox-Kemper, B., Ferrari, R., & Hallberg, R. (2008). Parameterization of mixed layer  
799 eddies. part i: Theory and diagnosis. *J. Phys. Oceanogr.*, *38*(6), 1145–1165.

- 800 Gaspar, P., Grégoris, Y., & Lefevre, J.-M. (1990). A simple eddy kinetic energy  
801 model for simulations of the oceanic vertical mixing: Tests at station papa and  
802 long-term upper ocean study site. *Journal of Geophysical Research: Oceans*,  
803 *95*(C9), 16179–16193.
- 804 Gent, P. R., & McWilliams, J. C. (1990). Isopycnal mixing in ocean circulation  
805 models. *J. Phys. Oceanogr.*, *20*(1), 150–155.
- 806 Greatbatch, R. J., Zhai, X., Kohlmann, J.-D., & Czeschel, L. (2010). Ocean eddy  
807 momentum fluxes at the latitudes of the gulf stream and the kuroshio exten-  
808 sions as revealed by satellite data. *Ocean Model.*, *60*(3), 617–628.
- 809 Groeskamp, S., LaCasce, J. H., McDougall, T. J., & Rogé, M. (2020). Full-depth  
810 global estimates of ocean mesoscale eddy mixing from observations and theory.  
811 *Geophys. Res. Lett.*, *47*(18), e2020GL089425.
- 812 Grooms, I. (2017). Simulations of eddy kinetic energy transport in barotropic turbu-  
813 lence. *Physical Review Fluids*, *2*(11), 113801.
- 814 Grooms, I., Nadeau, L.-P., & Smith, K. S. (2013). Mesoscale eddy energy locality in  
815 an idealized ocean model. *J. Phys. Oceanogr.*, *43*(9), 1911–1923.
- 816 Hawkins, E., Smith, R. S., Gregory, J. M., & Stainforth, D. A. (2016). Irreducible  
817 uncertainty in near-term climate projections. *Clim. Dyn.*, *46*(11-12), 3807–  
818 3819.
- 819 Jamet, Q., Deremble, B., Wienders, N., Uchida, T., & Dewar, W. (2021). On Wind-  
820 driven Energetics of Subtropical Gyres. *Journal of Advances in Modeling Earth*  
821 *Systems*, e2020MS002329.
- 822 Jansen, M. F., Adcroft, A., Khani, S., & Kong, H. (2019). Toward an energetically  
823 consistent, resolution aware parameterization of ocean mesoscale eddies. *Jour-*  
824 *nal of Advances in Modeling Earth Systems*, *11*(8), 2844–2860.
- 825 Jouanno, J., Capet, X., Madec, G., Roulet, G., & Klein, P. (2016). Dissipation of  
826 the energy imparted by mid-latitude storms in the southern ocean. *Ocean Sci.*,  
827 *12*(3), 743–769.
- 828 Kang, D., & Curchitser, E. N. (2015). Energetics of eddy–mean flow interactions in  
829 the Gulf Stream region. *J. Phys. Oceanogr.*, *45*(4), 1103–1120.
- 830 Large, W. G., & Yeager, S. G. (2004). Diurnal to decadal global forcing for ocean  
831 and sea-ice models: the data sets and flux climatologies.
- 832 Leroux, S., Brankart, J.-M., Albert, A., Brodeau, L., Molines, J.-M., Jamet, Q.,

- 833 Le Sommer, J., Penduff, T., and Brasseur, P. (2021). Ensemble quantifica-  
834 tion of short-term predictability of the ocean fine-scale dynamics: A western  
835 mediterranean test-case at kilometric-scale resolution. *Manuscript submitted to*  
836 *Ocean Sci.*,  $-( - )$ ,  $-$ .
- 837 Madec, G., Bourdallé-Badie, R., Bouttier, P.-A., Bricaud, C., Bruciaferri, D.,  
838 Calvert, D., Chanut, J., Clementi, E., Coward, A., Delrosso, D., and others  
839 (2017). Nemo ocean engine.
- 840 Mak, J., Maddison, J. R., Marshall, D. P., & Munday, D. R. (2018). Implementa-  
841 tion of a geometrically informed and energetically constrained mesoscale eddy  
842 parameterization in an ocean circulation model. *J. Phys. Oceanogr.*, *48*(10),  
843 2363–2382.
- 844 Marshall, D. P., & Adcroft, A. J. (2010). Parameterization of ocean eddies:  
845 Potential vorticity mixing, energetics and arnold’s first stability theorem.  
846 *Ocean Model.*, *32*(3-4), 188–204.
- 847 Mellor, G. L., & Yamada, T. (1982). Development of a turbulence closure model for  
848 geophysical fluid problems. *Reviews of Geophysics*, *20*(4), 851–875.
- 849 Mémin, E. (2014). Fluid flow dynamics under location uncertainty. *Geophys. Astro-*  
850 *phys. Fluid Dynamics*, *108*(2), 119–146.
- 851 Millot, C. (1985). Some features of the algerian current. *Journal of Geophysical Re-*  
852 *search: Oceans*, *90*(C4), 7169–7176.
- 853 Millot, C. (1999). Circulation in the western mediterranean sea. *Journal of Marine*  
854 *Systems*, *20*(1-4), 423–442.
- 855 Obaton, D., Millot, C., d’Hières, G. C., & Taupier-Letage, I. (2000). The Algerian  
856 Current: Comparisons between in situ and laboratory data sets. *Deep Sea Re-*  
857 *search Part I: Oceanographic Research Papers*, *47*(11), 2159–2190.
- 858 Poulain, P.-M., Centurioni, L., Özgökmen, T., Tarry, D., Pascual, A., Ruiz, S.,  
859 Mauri, E., Menna, M., and Notarstefano, G. (2021). On the structure and  
860 kinematics of an algerian eddy in the southwestern mediterranean sea. *Remote*  
861 *Sensing*, *13*(15), 3039.
- 862 Renault, L., Molemaker, M. J., Gula, J., Masson, S., and McWilliams, J. C. (2016).  
863 Control and stabilization of the gulf stream by oceanic current interaction with  
864 the atmosphere. *J. Phys. Oceanogr.*, *46*(11), 3439–3453.
- 865 Renault, L., Molemaker, M. J., McWilliams, J. C., Shchepetkin, A. F., Lemarié,

- 866 F., Chelton, D., Illig, S., and Hall, A. (2016). Modulation of wind work by  
867 oceanic current interaction with the atmosphere. *J. Phys. Oceanogr.*, 46(6),  
868 1685–1704.
- 869 Resseguier, V., Mémin, E., & Chapron, B. (2017). Geophysical flows under location  
870 uncertainty, part i random transport and general models. *Geophysical & Astro-*  
871 *physical Fluid Dynamics*, 111(3), 149–176.
- 872 Roquet, F., Madec, G., McDougall, T. J., & Barker, P. M. (2015). Accurate poly-  
873 nomial expressions for the density and specific volume of seawater using the  
874 TEOS-10 standard. *Ocean Modelling*, 90, 29–43.
- 875 Rossby, T. (1987). On the energetics of the gulf stream at 73w. *Journal of Marine*  
876 *Research*, 45(1), 59–82.
- 877 Schubert, R., Gula, J., Greatbatch, R. J., Baschek, B., & Biastoch, A. (2020). The  
878 Submesoscale Kinetic Energy Cascade: Mesoscale Absorption of Submesoscale  
879 Mixed Layer Eddies and Frontal Downscale Fluxes. *J. Phys. Oceanogr.*, 50(9),  
880 2573–2589.
- 881 Shchepetkin, A. F., & McWilliams, J. C. (2005). The regional oceanic modeling  
882 system (ROMS): a split-explicit, free-surface, topography-following-coordinate  
883 oceanic model. *Ocean modelling*, 9(4), 347–404.
- 884 Stainforth, D. A., Allen, M. R., Tredger, E. R., & Smith, L. A. (2007). Confidence,  
885 uncertainty and decision-support relevance in climate predictions. *Philosophical*  
886 *Transactions of the Royal Society A: Mathematical, Physical and Engineering*  
887 *Sciences*, 365(1857), 2145–2161.
- 888 Visbeck, M., Marshall, J., Haine, T., & Spall, M. (1997). Specification of eddy trans-  
889 fer coefficients in coarse-resolution ocean circulation models. *Journal of physi-*  
890 *cal oceanography*, 27(3), 381–402.
- 891 Waldman, R. (2016). *Etude multi-échelle de la convection océanique profonde en*  
892 *mer méditerranée: de l’observation à la modélisation climatique* (Unpublished  
893 doctoral dissertation). Université Paul Sabatier-Toulouse III.
- 894 Webster, F. (1961). The effect of meanders on the kinetic energy balance of the gulf  
895 stream. *Tellus*, 13(3), 392–401.
- 896 Webster, F. (1965). Measurements of eddy fluxes of momentum in the surface layer  
897 of the gulf stream. *Tellus*, 17(2), 240–245.
- 898 Wunsch, C., & Ferrari, R. (2004). Vertical mixing, energy, and the general circula-

tion of the oceans. *Annu. Rev. Fluid Mech.*, *36*, 281–314.