


An empirical orthogonal function analysis of ocean shoreline location on the Virginia barrier islands

James D. Haluska, Old Dominion University, Ocean Earth and Atmospheric Sciences Department, Center for Coastal Physical Oceanography

PROJECT LOCATION



Contact:
James Haluska
jdhaluska@gmail.com
757-635-4726 (cell and SMS)

Background

- Study location is along the East Coast of the US in the mid-Atlantic bight.
- Atlantic shorelines of eleven Virginia barrier islands from 1990 to 2014 digitized using LANDSAT 5, 7, and 8 satellite images and USDA NAIP aerial photos.
- Shoreline locations were combined into a 338 by 250 matrix for empirical orthogonal function (EOF) analysis.
- Resulting eigenvalues (temporal component) and eigenvectors (spatial component) were used for further analysis.
- Principal components (PC) calculated from eigenvalues correlated to sea level, wave height, the north Atlantic oscillation index (NAO), Arctic oscillation (AO), and the multivariate ENSO index (MEI).

Methods and Materials

- LANDSAT and aerial photos were assembled.
- All images were converted to the same geographical projection.
- The eastern shoreline of each island in each useable image was digitized.
- The digitized shorelines were used to produce a shore movement time series at 300m intervals using the Digital Shoreline Analysis System.
- All island shoreline change data combined and analyzed..
- First four eigenvalues and eigenvectors (EV) used for further analysis.
- Correlation for the various indices, cycles, and environmental factors with PCs 1 to 4 determined.

RESULTS





- Table 1 is the fraction of EOF variance explained for both detrended and non-detrended data.
- The years 2004 to 2005 and 2010 to 2011, were years of large shoreline changes to the barrier islands.
- Table 2 summarizes the correlation coefficients and coefficients of determination for the indices and environmental factors used for comparison to principal components 1 to 4 for the non-detrended data
- PC1 correlated to sea level ($r^2 = 0.41$). The NAO ($r^2 = 0.05$) and MEI ($r^2 = 0.12$) also correlated with PC1 at the 0.99 confidence level.
- The NAO correlated with PC4 ($r^2 = 0.05$) .
- Eigenvector plots for EVs 1 to 4 indicated that most of the spatial changes to the islands were occurring at the inlets.

Table 1. Fraction of the EOF variance explained for non-detrended and detrended shoreline location measurements. A linear trend was removed for the detrended data analysis.		
EOF	Non Detrended	Detrended
1	0.59	0.33
2	0.13	0.15
3	0.08	0.08
4	0.06	0.06
5	0.03	0.04
6	0.02	0.03
7	0.02	0.02
8	0.01	0.02
9	0.01	0.02
10	0.01	0.02





Table 2. Correlation coefficients (RHO), significance value (PVAL), and coefficients of determination (Coef_Det) for principal components 1 to 4. PVAL >0.99% significance highlighted												
Factor	PC1			PC2			PC3			PC4		
	PVAL	RHO	Coef_Det	PVAL	RHO	Coef_Det	PVAL	RHO	Coef_Det	PVAL	RHO	Coef_Det
AO	0.0137	-0.142	0.020	0.030	-0.100	0.010	0.011	0.146	0.020	0.0324	-0.010	
NAO	1.15E-09	-0.217	0.047	0.182	-0.030	0.631	0.146			0.00115	-0.221	0.048
MEI	1.15E-09	-0.348	0.117	0.030	-0.130	0.017	0.030	0.005	0.0286	-0.027	0.016	
Waves	0.0116	0.146	0.021	0.057	0.110	0.739	0.015			0.075	0.013	
SeaLevel	1.7E-35	0.637	0.405	0.634	-0.029	0.019	-0.135	0.018	0.2813	0.038		

Plots of the first four principal components (PC) and eigenvectors calculated from non-detrended data of shoreline movement for the eleven Virginia barrier islands. The first four eigenvalues and eigenvectors explain 86 percent of the total data set variance. The PCs are calculated from the eigenvalues. The eigenvectors are the spatial component of the data set. The island names are added to the eigenvector plots for spatial reference.

Principle Components



Eigenvectors



CONCLUSIONS

- Sea level rise is the largest environmental factor affecting the eleven islands of the five factors evaluated in this study. .
- The NAO and MEI indices are important in describing episodic shoreline events on these islands.
- Steady shoreline retreat narrows parts of the islands to the point that they are sensitive to episodes of sudden shoreline loss episodic events.
- Large shoreline change events are episodic and are associated with extra-tropical storms, hurricanes, and high water tidal events.