Spatially Varying Covariate Model: A Hierarchical Bayesian Framework for Precipitation Frequency Analysis in the Gulf Coast

Yuchen Lu*[1], Benjamin Seiyon Lee[2], James Doss-Gollin[1]

*yl238@rice.edu, [1]Department of Civil and Environmental Engineering, Rice University, [2]Department of Statistics, George Mason University

Motivation

Estimates of precipitation frequency are widely used in risk assessment and management. Yet despite recognition that interannual variability and **climate change** affect hazards, most current guidance (e.g. Atlas 14) assumes **stationarity**.

Knowledge Gap

Incorporating nonstationarity into precipitation frequency estimates can dramatically amplify **parametric uncertainty**. Here, we demonstrate how hierarchical **spatial pooling** can enhance inference for nonstationary extreme value models.



Fagnant et. al (2020): moving window analysis on stations in SE Texas and W Louisiana. Estimated precipitation frequencies and trends vary dramatically between nearby stations, **motivating more robust estimation strategies.**

Methodology

We assume that the impacts of climate on extreme precipitation characteristics are spatially coherent, represented by a **latent spatial field** describing the smoothly varying GEV parameters (details provided below).

Bayesian Hierarchical Model

Annual maximum rainfall at location s and year t follows the Generalized Extreme Value (GEV) distribution

$$y(s,t) \sim GEV(\mu(s,t), \sigma(s,t), \xi)$$

Process-Informed Nonstationary Model

We condition the GEV parameters on climate time series $x_j(t)$ (log of global CO_2 concentration)

$$\mu(s,t) = \mu_0(s) + \sum_{j=1}^{J} \beta_j^{\mu}(s) x_j(t) \quad \sigma(s,t) = \sigma_0(s) + \sum_{j=1}^{J} \beta_j^{\sigma}(s) x_j(t)$$

Latent parameters are smooth in space, implemented with a **Gaussian Process hierarchical prior.** We conduct **Bayesian** Inference via Markov Chain Monte Carlo.

Robust Trends in Extreme Precipitation



Posterior mean of coefficients of the anomalies of log of CO2 concentration on (T) the location and (B) the scale parameters estimated from (L) nonstatiaonry model at separate stations and (R) spatially varying covariate model

Spatially Consistent Increases in Heavy Rainfall Probabilities



Posterior mean of return levels estimates (T) in 2022 (B) difference between 2022 and 1940 for the (L) 10 year and (R) 100 year return periods

Well Calibrated and Reduced Uncertainties

Our Spatially Varying Covariate Model improves estimates by (1) reducing uncertainty compared to conventional nonstationary model simulated at separate stations and (2) achieving statistical calibration.



(L) Quantiles of the observation records given the simulated posterior GEV distributions. An ideal model would have a uniform distribution.
Return level estimates in Houston with uncertainty boundary using (M) nonstatiaonry model at separate stations (R) Spatially Varying Covariate Model

Comparison with Atlas 14

	Atlas 14	Spatially Varying Covariate Model
Stationarity	Stationary	Process-Informed Nonstationarity
Regionalization	Region of Influence	Hierarchical Gaussian Process
Inference	L-moments	Bayesian Inference

We estimate **higher current (2022) hazard** than Atlas 14 in most of the domain. Our current estimates are lower in the area directly impacted by Hurricane Harvey (SE Texas; 2017).



Return level estimates from the spatially varying covariate model minus that from Atlas 14

Conclusions

Our **Spatially Varying Covariates** model:

1. Identifies robust and spatially coherent trends
2. Improves estimation

We find **increasing risk of 24-hour precipitation** over the study area driven by climate change. This model can be used for IDF curves and to other spatially and temporally varying climate hazards.

References

Perica, S., Pavlovic, S., St Laurent, M., Trypaluk, C., Unruh, D., & Wilhite, O. (2018). Precipitation-Frequency Atlas of the United States. Volume 11, Version 2.0. Texas.

Nielsen-Gammon, J. W. (2020). Observation-Based Estimates of Present-Day and Future Climate Change Impacts on Heavy Rainfall in Harris County.

Ossandón, Á., Rajagopalan, B., & Kleiber, W. (2021). Spatial-temporal multivariate semi-Bayesian hierarchical framework for extreme precipitation frequency analysis. Journal of Hydrology, 600, 126499.

Fagnant, C., Gori, A., Sebastian, A., Bedient, P. B., & Ensor, K. B. (2020). Characterizing spatiotemporal trends in extreme precipitation in Southeast Texas. *Natural Hazards*, *104*, 1597-1621.