

## Introduction

- **Subtropical Mode Waters** (STMW) are extensive volume of water in the oceanic upper layers. In the South Atlantic it forms from winter to mid-spring (from July to October) and persists throughout the year.
- Temperature (T) and salinity (S) are homogeneous horizontally and vertically. The STMW is usually formed in the warm side of west boundary currents in the subtropical gyres and in oceanic fronts (Figure 1).
- The STMW formation and its advection can influence biogeochemical processes in the subtropical gyre by introducing variability in the subsuperficial nutrients reservoir (Palter et al., 2005).




Figure 1. Global distribution of the STMWs. Talley et al., 2011.

## Objetives

We aim is to determine how STMW in the southwest South Atlantic influences the concentration of nutrients in the region. Our specific goals are to:

- Detect the STMW in the South Atlantic;
- Correlate nitrate concentration during the formation of STMW in vertical profiles;
- Determine how nutrients and chlorophyll-a (Chl-a) concentrations vary in interannual scale due to the formation of STMW.

## Metodology

The study area extends from 60°W to 10°W and 25°S to 50°S. Three monthly resolution datasets were used:

- T and S from WOA13 (Garcia et al 2014; Locarnini et al. 2013) with a spatial resolution of  $1^\circ \times 1^\circ$ ;
- $\text{NO}_3$ ,  $\text{SiO}_2$  e  $\text{PO}_4$  concentrations from Bio Analysis 018 reanalysis model from CMEMS (Paul et al., 2015) with a spatial resolution of  $0.5^\circ \times 0.5^\circ$ ;
- T and S from the In Situ Analysis System (ISAS) (Gaillard et al., 2009) from 2002 to 2012 with a resolution in longitude of  $0.5^\circ$ .

The data were selected for the study area and interpolated in the same spatial grid. The STMW was identified through T, S and potential vorticity (q) values determined by Sato and Polito (2014):

T(°C)	S	q ( $m^{-1}s^{-1}$ )
14.1 - 15.9	35.4 - 35.8	$1.5 \times 10^{-10}$

The  $\text{NO}_3$  vertical profiles in regions with and without STMW formation were compared (Figure 3) based on Palter et al. (2005). The mean concentration of  $\text{NO}_3$ ,  $\text{SiO}_2$ ,  $\text{PO}_4$  and Chl-a were computed between 0 and 200 meters in the water column from 2002 to 2012 (Figure 4), for the STMW formation area: 42.5°W to 47°W and 35°S to 37°S.

## Finding the STMW in the climatology

- STMW mean thickness is  $112 \pm 5$  m in the summer and  $155 \pm 20$  m in the winter;
- The STMW can only be detected in the subsurface in summer and autumn. From winter to mid-spring, the STMW outcrops at the surface due to its formation.




Figure 2. Seasonal mean thickness of STMW (WOA13). Rectangle: area where the mean concentrations of nutrients and Chl-a were calculated (Figure 4). The X position indicates the vertical profiles on Figure 3.

## Deepening of the nutricline

- Analysis of the mean vertical profiles of T (ISAS) and  $\text{NO}_3$  (CMEMS) indicates that the  $\text{NO}_3$  nutricline deepens in regions where a thicker STMW was found (Figure 3a) compared to regions with thinner STMW (Figure 3b) or regions without STMW (Figure 3c);
- The nutricline mean depth is of the order of  $230 \pm 8$  m in regions with formation of STMW (Figures 3a and 3b) and  $150 \pm 8$  m outside.




Figure 3. Vertical profiles of T (ISAS) and  $\text{NO}_3$  ( $\mu\text{mol/L}$ ) (CMEMS), at 36.5°S, at different longitudes: 36.5°W (left), 24°W (center) and 20.5°W (right), in September 2012.

## Relation between STMW thickness, nutrients and chlorophyll-a

- The concentration of  $\text{NO}_3$ ,  $\text{SiO}_2$  and  $\text{PO}_4$  decrease during the formation of STMW. The Chl-a concentration increases from  $0.2 \mu\text{mol/L}$  in summer to  $0.35 \mu\text{mol/L}$  during the formation period (Figure 4);
- The formation of STMW likely enhances water mixing (Figure 4 and 5), taking phytoplankton to deeper waters with low-light conditions and leading to its photoacclimation. This process can be noticed by high Chl-a during formation.
- Significant correlation is found between the nutricline depth and thickness of STMW (Figure 5).




Figure 4. Interannual mean thickness of STMW (m) (ISAS), nutrients and Chl-a concentration ( $\mu\text{mol/L}$ ) (CMEMS):  $\text{SiO}_2$ ,  $\text{NO}_3$ ,  $\text{PO}_4$ , Chl-a, between 0 and 200 m, from 2002 to 2012.




Figure 5. Monthly mean nutricline depth (m) based on the  $\text{NO}_3$  maximum gradient (Palter et al. 2005) (in green) and monthly mean thickness of the superficial STMW (m) (in blue) for the region A.

## Conclusion

- The STMW main formation occurs during the winter until the mid-spring;
- In the formation region, the deepening of the nutricline coincides with the occurrence of the STMW which is responsible for the superficial layer mixing in the oceans and therefore changes the nutrients concentration on surface;
- The formation of STMW in the southwest Atlantic influences the concentration of nutrients and Chl-a due to convective mixing between layers below the nutricline and superficial waters.
- This phenomenon is key for understanding the relationship between physical and biological processes in the subtropical gyre.

## References

- Gaillard, F., E. Brion, and R. Charraudeau (2009). Isas-v5: Description of the method and user manual. IFREMER Rapport LPO, 09-04.
- Garcia, H., R. Locarnini, T. Boyer, J. Antonov, O. Baranova, M. Zweng, J. Reagan, and D. Johnson (2014). World ocean atlas 2013, volume 4: Dissolved inorganic nutrients (phosphate, nitrate, silicate), s. A. Mishonov Technical Ed, 1-25.
- Locarnini, R., A. Mishonov, J. Antonov, T. Boyer, H. Garcia, O. Baranova, M. Zweng, C. Paver, J. Reagan, D. Johnson, et al. (2013). World Ocean Atlas 2013. Vol. 1: Temperature. A. Mishonov, Technical Ed. NOAA Atlas NESDIS 73, 40.
- Palter, J. B., M. S. Lozier, and R. T. Barber (2005). The effect of advection on the nutrient reservoir in the North Atlantic subtropical gyre. Nature 437(7059), 687-692.
- Paul, J., C. Perruche, and A. Hameau (2015). Global ocean biogeochemistry analysis and weekly forecast: Product user manual.
- Sato, O. and P. Polito (2014). Observation of South Atlantic subtropical mode waters with Argo profiling float data. JGR: Oceans 119(5), 2860-2881.
- Talley, L. D. (2011). Descriptive physical oceanography: an introduction. Academic press.